



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 10:16 PM UTC

PDB ID : 4LNV / pdb_00004lnv
Title : Crystal Structure of TEP1s
Authors : Le, B.V.; Williams, M.; Logarajah, S.; Baxter, R.H.G.
Deposited on : 2013-07-12
Resolution : 3.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 3.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1131 (3.80-3.60)
Clashscore	190562	1171 (3.80-3.60)
Ramachandran outliers	187476	1129 (3.80-3.60)
Sidechain outliers	187428	1126 (3.80-3.60)
RSRZ outliers	180081	1130 (3.80-3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1323	 78% 17% . .
1	B	1323	 63% 16% . 20%
1	C	1323	 78% 17% . .
2	D	2	 100%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 27034 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Thioester-containing protein I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1276	Total	C	N	O	S	0	0	0
			9704	6210	1583	1866	45			
1	B	1063	Total	C	N	O	S	0	0	0
			7461	4750	1225	1454	32			
1	C	1276	Total	C	N	O	S	0	0	0
			9631	6163	1573	1850	45			

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1339	HIS	-	expression tag	UNP Q9GYW4
A	1340	HIS	-	expression tag	UNP Q9GYW4
A	1341	HIS	-	expression tag	UNP Q9GYW4
A	1342	HIS	-	expression tag	UNP Q9GYW4
A	1343	HIS	-	expression tag	UNP Q9GYW4
A	1344	HIS	-	expression tag	UNP Q9GYW4
B	1339	HIS	-	expression tag	UNP Q9GYW4
B	1340	HIS	-	expression tag	UNP Q9GYW4
B	1341	HIS	-	expression tag	UNP Q9GYW4
B	1342	HIS	-	expression tag	UNP Q9GYW4
B	1343	HIS	-	expression tag	UNP Q9GYW4
B	1344	HIS	-	expression tag	UNP Q9GYW4
C	1339	HIS	-	expression tag	UNP Q9GYW4
C	1340	HIS	-	expression tag	UNP Q9GYW4
C	1341	HIS	-	expression tag	UNP Q9GYW4
C	1342	HIS	-	expression tag	UNP Q9GYW4
C	1343	HIS	-	expression tag	UNP Q9GYW4
C	1344	HIS	-	expression tag	UNP Q9GYW4

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	D	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...


Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		



• Molecule 1: Thioester-containing protein I

Chain C: 78% 17%





● Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D: 100%

NAG1
NAG2

4 Data and refinement statistics

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	196.47Å 196.47Å 225.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.12 – 3.70 49.12 – 3.70	Depositor EDS
% Data completeness (in resolution range)	99.7 (49.12-3.70) 99.6 (49.12-3.70)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.89	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.95 (at 3.67Å)	Xtriage
Refinement program	REFMAC refmac _5.7.0025	Depositor
R, R_{free}	0.237 , 0.276 0.211 , 0.250	Depositor DCC
R_{free} test set	4836 reflections (5.31%)	wwPDB-VP
Wilson B-factor (Å ²)	147.0	Xtriage
Anisotropy	0.003	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 114.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.196 for h,-k,-l	Xtriage
Reported twinning fraction	0.504 for H, K, L 0.496 for K, H, -L	Depositor
Outliers	2 of 90746 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	27034	wwPDB-VP
Average B, all atoms (Å ²)	157.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 23.99 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.1834e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	0/9902	0.80	4/13515 (0.0%)
1	B	0.55	0/7592	0.85	6/10410 (0.1%)
1	C	0.51	0/9825	0.79	5/13417 (0.0%)
All	All	0.52	0/27319	0.81	15/37342 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1067	THR	N-CA-C	7.79	119.40	111.07
1	C	1065	GLY	N-CA-C	6.37	121.72	112.55
1	A	1065	GLY	N-CA-C	6.11	121.35	112.55
1	B	1065	GLY	N-CA-C	6.07	121.28	112.60
1	A	190	VAL	CA-C-N	6.04	127.39	119.84

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1064	TRP	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	9704	0	9100	143	0
1	B	7461	0	6369	103	0
1	C	9631	0	8983	135	0
2	D	28	0	25	0	0
3	A	98	0	91	0	0
3	B	14	0	13	0	0
3	C	98	0	91	3	0
All	All	27034	0	24672	376	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 376 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:1402:NAG:H4	3:C:1403:NAG:H5	1.59	0.84
1:A:62:ASN:HB3	1:B:970:VAL:HG23	1.63	0.79
1:C:862:GLN:HA	1:C:1109:THR:HG21	1.64	0.79
1:A:1100:ARG:HG2	1:A:1106:PHE:HE1	1.48	0.77
1:A:1100:ARG:HG2	1:A:1106:PHE:CE1	2.20	0.77

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1266/1323 (96%)	1154 (91%)	101 (8%)	11 (1%)	14	45
1	B	1011/1323 (76%)	898 (89%)	94 (9%)	19 (2%)	6	33
1	C	1266/1323 (96%)	1157 (91%)	97 (8%)	12 (1%)	14	45
All	All	3543/3969 (89%)	3209 (91%)	292 (8%)	42 (1%)	10	40

5 of 42 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	642	ARG
1	A	1160	GLU
1	C	642	ARG
1	C	1160	GLU
1	A	418	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	997/1177 (85%)	940 (94%)	57 (6%)	18	45
1	B	670/1177 (57%)	624 (93%)	46 (7%)	14	41
1	C	979/1177 (83%)	928 (95%)	51 (5%)	21	47
All	All	2646/3531 (75%)	2492 (94%)	154 (6%)	18	45

5 of 154 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	370	ASP
1	C	1143	THR
1	C	457	MET
1	C	803	THR
1	C	1279	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 38 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	605	ASN
1	C	1256	ASN
1	C	907	GLN
1	C	1022	ASN
1	C	1324	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	D	1	2,1	14,14,15	0.71	0	17,19,21	1.89	4 (23%)
2	NAG	D	2	2	14,14,15	0.44	0	17,19,21	0.94	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	D	1	2,1	-	3/6/23/26	0/1/1/1
2	NAG	D	2	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1	NAG	C2-N2-C7	4.82	129.35	122.90
2	D	1	NAG	O5-C1-C2	-3.33	106.13	111.29
2	D	1	NAG	C4-C3-C2	3.03	115.46	111.02
2	D	2	NAG	C1-O5-C5	2.46	115.48	112.19
2	D	1	NAG	C3-C4-C5	2.40	114.59	110.23

There are no chirality outliers.


All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	1	NAG	C4-C5-C6-O6
2	D	1	NAG	O5-C5-C6-O6
2	D	1	NAG	C3-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	2003	1	14,14,15	0.49	0	17,19,21	0.82	0
3	NAG	A	2002	1	14,14,15	0.49	0	17,19,21	0.85	0
3	NAG	C	1405	1	14,14,15	0.51	0	17,19,21	1.14	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	A	2009	1	14,14,15	0.84	1 (7%)	17,19,21	1.63	3 (17%)
3	NAG	C	1406	1	14,14,15	0.29	0	17,19,21	0.56	0
3	NAG	C	1407	1	14,14,15	0.59	0	17,19,21	1.38	2 (11%)
3	NAG	A	2008	1	14,14,15	0.44	0	17,19,21	2.57	3 (17%)
3	NAG	C	1401	1	14,14,15	0.52	0	17,19,21	1.15	1 (5%)
3	NAG	C	1402	1	14,14,15	0.47	0	17,19,21	0.66	0
3	NAG	A	2001	1	14,14,15	0.55	0	17,19,21	0.98	1 (5%)
3	NAG	B	1401	1	14,14,15	0.45	0	17,19,21	0.99	1 (5%)
3	NAG	C	1404	1	14,14,15	0.49	0	17,19,21	0.90	1 (5%)
3	NAG	A	2005	1	14,14,15	0.65	0	17,19,21	2.33	5 (29%)
3	NAG	A	2004	-	14,14,15	0.45	0	17,19,21	0.90	1 (5%)
3	NAG	C	1403	-	14,14,15	0.47	0	17,19,21	1.62	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	2003	1	-	1/6/23/26	0/1/1/1
3	NAG	A	2002	1	-	2/6/23/26	0/1/1/1
3	NAG	C	1405	1	-	2/6/23/26	0/1/1/1
3	NAG	A	2009	1	-	2/6/23/26	0/1/1/1
3	NAG	C	1406	1	-	4/6/23/26	0/1/1/1
3	NAG	C	1407	1	-	2/6/23/26	0/1/1/1
3	NAG	A	2008	1	-	2/6/23/26	0/1/1/1
3	NAG	C	1401	1	-	0/6/23/26	0/1/1/1
3	NAG	C	1402	1	-	2/6/23/26	0/1/1/1
3	NAG	A	2001	1	-	2/6/23/26	0/1/1/1
3	NAG	B	1401	1	-	2/6/23/26	0/1/1/1
3	NAG	C	1404	1	-	0/6/23/26	0/1/1/1
3	NAG	A	2005	1	-	3/6/23/26	0/1/1/1
3	NAG	A	2004	-	-	1/6/23/26	0/1/1/1
3	NAG	C	1403	-	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	2009	NAG	C1-C2	2.40	1.55	1.52

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	2008	NAG	C1-O5-C5	9.71	125.20	112.19
3	A	2005	NAG	C1-O5-C5	6.28	120.61	112.19
3	A	2005	NAG	C2-N2-C7	4.75	129.27	122.90
3	A	2009	NAG	C4-C3-C2	4.56	117.70	111.02
3	C	1403	NAG	C1-O5-C5	4.17	117.77	112.19

There are no chirality outliers.

5 of 27 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	2005	NAG	C1-C2-N2-C7
3	B	1401	NAG	C4-C5-C6-O6
3	C	1402	NAG	C4-C5-C6-O6
3	A	2005	NAG	O5-C5-C6-O6
3	A	2008	NAG	O5-C5-C6-O6

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	1406	NAG	1	0
3	C	1402	NAG	2	0
3	C	1403	NAG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	1276/1323 (96%)	-0.29	4 (0%)	90 76	113, 151, 175, 200	0
1	B	1063/1323 (80%)	-0.38	4 (0%)	88 72	125, 165, 200, 200	0
1	C	1276/1323 (96%)	-0.34	1 (0%)	92 86	123, 160, 191, 200	0
All	All	3615/3969 (91%)	-0.34	9 (0%)	91 80	113, 157, 194, 200	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	563	LEU	3.3
1	B	457	MET	3.1
1	B	121	ILE	3.0
1	C	563	LEU	2.6
1	B	264	TYR	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q < 0.9
2	NAG	D	2	14/15	0.87	0.07	169,171,172,173	0
2	NAG	D	1	14/15	0.92	0.08	167,168,169,170	0

The following is a graphical depiction of the model fit to experimental electron density for oligosac-

charide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	A	2009	14/15	0.70	0.10	166,168,168,168	0
3	NAG	C	1404	14/15	0.74	0.08	173,174,175,175	0
3	NAG	A	2005	14/15	0.75	0.11	170,172,173,173	0
3	NAG	B	1401	14/15	0.82	0.08	175,176,176,176	0
3	NAG	A	2008	14/15	0.82	0.07	152,152,152,153	0
3	NAG	C	1407	14/15	0.82	0.09	168,169,169,169	0
3	NAG	A	2001	14/15	0.83	0.10	165,168,168,169	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	C	1401	14/15	0.83	0.07	166,167,168,168	0
3	NAG	C	1405	14/15	0.84	0.07	162,163,163,164	0
3	NAG	C	1406	14/15	0.84	0.08	174,175,176,176	0
3	NAG	A	2004	14/15	0.84	0.08	177,177,178,178	0
3	NAG	A	2003	14/15	0.88	0.08	160,160,161,162	0
3	NAG	A	2002	14/15	0.91	0.06	156,157,158,158	0
3	NAG	C	1403	14/15	0.92	0.07	183,185,185,185	0
3	NAG	C	1402	14/15	0.94	0.06	166,166,167,168	0

6.5 Other polymers [i](#)

There are no such residues in this entry.