



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 02:52 AM UTC

PDB ID : 6LN2 / pdb_00006ln2
Title : Crystal structure of full length human GLP1 receptor in complex with Fab fragment (Fab7F38)
Authors : Wu, F.; Yang, L.; Hang, K.; Laursen, M.; Wu, L.; Han, G.W.; Ren, Q.; Roed, N.K.; Lin, G.; Hanson, M.; Jiang, H.; Wang, M.; Reedtz-Runge, S.; Song, G.; Stevens, R.C.
Deposited on : 2019-12-28
Resolution : 3.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	469	
2	B	213	
3	C	226	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6524 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glucagon-like peptide 1 receptor,Rubredoxin,Glucagon-like peptide 1 receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	437	Total	C	N	O	S	0	0	0
			3178	2083	508	564	23			

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	193	CYS	SER	engineered mutation	UNP P43220
A	196	PHE	ILE	engineered mutation	UNP P43220
A	225	ALA	SER	engineered mutation	UNP P43220
A	233	CYS	MET	engineered mutation	UNP P43220
A	271	ALA	SER	engineered mutation	UNP P43220
A	317	CYS	ILE	engineered mutation	UNP P43220
A	318	ILE	GLY	engineered mutation	UNP P43220
A	346	ALA	LYS	engineered mutation	UNP P43220
A	347	PHE	CYS	engineered mutation	UNP P43220
A	361	CYS	GLY	engineered mutation	UNP P43220
A	387	ASP	GLU	engineered mutation	UNP P43220

- Molecule 2 is a protein called Fab7F38_light chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	213	Total	C	N	O	S	0	0	0
			1628	1025	266	329	8			

- Molecule 3 is a protein called Fab7F38_heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	226	Total	C	N	O	S	0	0	0
			1666	1056	266	337	7			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula:

C₈H₁₅NO₆) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Zn	0	0
			1	1		

- Molecule 6 is N-{4-[(R)-(3,3-dimethylcyclobutyl)({6-[4-(trifluoromethyl)-1H-imidazol-1-yl]pyridin-3-yl}amino)methyl]benzene-1-carbonyl}-beta-alanine (CCD ID: 97Y) (formula: C₂₆H₂₈F₃N₅O₃) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	F	N	O	0	0
			37	26	3	5	3		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glucagon-like peptide 1 receptor,Rubredoxin,Glucagon-like peptide 1 receptor



- Molecule 2: Fab7F38_light chain



- Molecule 3: Fab7F38_heavy chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	63.04Å 64.66Å 322.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 3.20 30.00 – 3.20	Depositor EDS
% Data completeness (in resolution range)	97.0 (30.00-3.20) 96.8 (30.00-3.20)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.71 (at 3.18Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.213 , 0.262 (Not available) , 0.284	Depositor DCC
R_{free} test set	1098 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	112.7	Xtriage
Anisotropy	0.314	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 107.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	0.054 for k,h,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6524	wwPDB-VP
Average B, all atoms (Å ²)	136.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, 97Y, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.69	0/3276	0.92	2/4518 (0.0%)
2	B	1.00	3/1669 (0.2%)	1.00	0/2270
3	C	0.86	0/1710	0.91	0/2340
All	All	0.82	3/6655 (0.0%)	0.94	2/9128 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	77	MET	SD-CE	-8.57	1.58	1.79
2	B	11	MET	SD-CE	5.85	1.94	1.79
2	B	85	TYR	CA-C	-5.37	1.46	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	390	PHE	CA-CB-CG	9.23	123.03	113.80
1	A	377	GLY	N-CA-C	-5.64	108.47	114.67

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3178	0	2715	44	0
2	B	1628	0	1558	26	0
3	C	1666	0	1556	27	0
4	A	14	0	13	0	0
5	A	1	0	0	0	0
6	A	37	0	0	0	0
All	All	6524	0	5842	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (95) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:360:LEU:HD21	1:A:394:GLN:HG3	1.48	0.93
1:A:82:ASN:OD1	1:A:101:TYR:CD1	2.32	0.83
2:B:31:TYR:CD2	2:B:49:ARG:HG2	2.20	0.76
2:B:157:ASN:ND2	2:B:178:LEU:HD12	2.06	0.70
1:A:224:LEU:HA	1:A:227:ARG:HE	1.62	0.64
1:A:1011:TYR:HB2	1:A:1041:LEU:HD12	1.84	0.60
3:C:136:LEU:O	3:C:150:LEU:HA	2.02	0.60
3:C:171:LEU:HD21	3:C:194:VAL:HG21	1.83	0.59
3:C:154:VAL:HG11	3:C:162:VAL:HG11	1.85	0.57
1:A:289:TYR:HA	1:A:293:ASP:HB3	1.88	0.56
2:B:77:MET:HE1	2:B:103:LEU:HD21	1.87	0.56
2:B:123:GLN:C	2:B:125:LYS:H	2.15	0.55
3:C:151:GLY:HA2	3:C:166:TRP:CZ2	2.42	0.55
1:A:384:LEU:O	1:A:388:LEU:HD22	2.07	0.55
3:C:222:ARG:NH1	3:C:224:GLU:HB3	2.22	0.54
1:A:317:CYS:HB3	1:A:365:VAL:HG13	1.91	0.53
2:B:119:PRO:HD3	2:B:131:VAL:HG22	1.91	0.52
1:A:1013:TYR:CZ	1:A:1033:ILE:HD11	2.45	0.52
2:B:3:VAL:H	2:B:26:SER:HB3	1.74	0.52
3:C:160:GLU:HG3	3:C:161:PRO:HA	1.91	0.51
3:C:171:LEU:CD2	3:C:194:VAL:HG21	2.41	0.51
1:A:82:ASN:OD1	1:A:101:TYR:CE1	2.64	0.50
1:A:235:TYR:HA	1:A:281:VAL:HG21	1.93	0.50
3:C:210:VAL:HB	3:C:219:VAL:HG23	1.93	0.50
2:B:31:TYR:CE2	2:B:49:ARG:HG2	2.45	0.50
1:A:345:ILE:HD12	1:A:346:ALA:H	1.76	0.50
1:A:264:TRP:HZ3	1:A:268:LEU:HD11	1.76	0.50
2:B:134:LEU:HD22	3:C:193:VAL:HG21	1.94	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:98:ARG:HB3	3:C:114:TYR:HB2	1.92	0.50
1:A:190:ARG:HH12	1:A:394:GLN:CD	2.21	0.49
1:A:297:TRP:N	1:A:297:TRP:CD1	2.81	0.49
1:A:83:VAL:HG23	1:A:100:VAL:HG13	1.94	0.48
2:B:15:PRO:HG3	2:B:105:ILE:HD11	1.95	0.48
1:A:102:ARG:HH11	1:A:119:PRO:HG2	1.79	0.48
2:B:89:GLN:HG2	2:B:91:HIS:H	1.79	0.48
3:C:164:VAL:HG21	3:C:190:LEU:HD11	1.96	0.48
1:A:327:VAL:O	1:A:331:VAL:HG23	2.15	0.47
3:C:40:LYS:HB3	3:C:43:GLN:HB2	1.96	0.47
3:C:68:ALA:HA	3:C:82:GLU:O	2.14	0.47
1:A:360:LEU:HD21	1:A:394:GLN:CG	2.33	0.47
1:A:150:VAL:O	1:A:154:LEU:HB2	2.15	0.47
1:A:385:PHE:HA	1:A:388:LEU:HD22	1.97	0.47
2:B:89:GLN:NE2	2:B:92:THR:O	2.47	0.47
1:A:166:LEU:HB2	1:A:181:LEU:HD21	1.97	0.47
1:A:162:ALA:HA	1:A:165:ILE:HD12	1.98	0.46
2:B:210:ARG:HH21	2:B:210:ARG:HG3	1.78	0.46
1:A:280:PHE:C	1:A:283:PRO:HD2	2.40	0.46
1:A:85:CYS:H	1:A:98:GLY:HA3	1.80	0.46
1:A:282:VAL:CG2	1:A:283:PRO:HD3	2.46	0.46
2:B:77:MET:HE1	2:B:103:LEU:CD2	2.46	0.45
1:A:345:ILE:HD12	1:A:346:ALA:N	2.31	0.45
3:C:50:TYR:CD1	3:C:50:TYR:C	2.93	0.45
3:C:151:GLY:HA2	3:C:166:TRP:HZ2	1.81	0.45
1:A:235:TYR:HB2	1:A:281:VAL:HG11	1.99	0.45
3:C:7:SER:HB3	3:C:21:SER:H	1.82	0.45
1:A:277:PRO:HA	1:A:280:PHE:CE2	2.52	0.44
1:A:146:ILE:O	1:A:150:VAL:HG22	2.16	0.44
2:B:53:LEU:HD12	2:B:57:VAL:CG1	2.47	0.44
2:B:13:ALA:O	2:B:105:ILE:HA	2.18	0.44
1:A:406:ASN:HD22	1:A:407:ASN:N	2.15	0.44
1:A:255:LEU:HD21	1:A:331:VAL:HG22	1.99	0.44
1:A:86:PRO:HB3	1:A:88:TYR:CE2	2.52	0.44
2:B:10:ILE:HG22	2:B:102:LYS:HB3	1.99	0.44
3:C:38:LYS:HB2	3:C:48:ILE:HD11	2.00	0.44
2:B:92:THR:O	2:B:95:TRP:HD1	2.01	0.43
1:A:391:THR:O	1:A:391:THR:HG22	2.19	0.43
2:B:14:SER:OG	2:B:17:GLU:HG3	2.19	0.43
2:B:157:ASN:ND2	2:B:178:LEU:CD1	2.80	0.43
2:B:188:HIS:O	2:B:210:ARG:NH2	2.52	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:157:TYR:CE1	3:C:188:TYR:HB2	2.54	0.42
1:A:394:GLN:O	1:A:394:GLN:HG2	2.20	0.42
2:B:88:GLN:HG2	2:B:89:GLN:N	2.33	0.42
3:C:171:LEU:HD21	3:C:194:VAL:CG2	2.49	0.42
1:A:81:VAL:O	1:A:81:VAL:HG13	2.19	0.42
1:A:356:LEU:HD21	1:A:398:VAL:HG22	2.02	0.42
3:C:166:TRP:CH2	3:C:208:CYS:HB2	2.55	0.42
2:B:1:GLN:HB2	2:B:94:PRO:HD2	2.02	0.42
3:C:196:VAL:HG11	3:C:206:TYR:CE1	2.55	0.42
1:A:270:VAL:HG13	1:A:274:TRP:HD1	1.85	0.41
3:C:190:LEU:HD12	3:C:190:LEU:C	2.44	0.41
1:A:156:PHE:CE1	1:A:188:ILE:HG23	2.55	0.41
3:C:196:VAL:HB	3:C:197:PRO:HD2	2.02	0.41
3:C:8:GLY:O	3:C:119:THR:HG23	2.19	0.41
1:A:59:ASP:HB3	2:B:92:THR:HG22	2.03	0.41
1:A:356:LEU:CD2	1:A:398:VAL:HG22	2.51	0.41
2:B:77:MET:CE	2:B:103:LEU:HD21	2.49	0.41
2:B:182:LYS:HE3	2:B:186:GLU:OE1	2.20	0.41
1:A:276:VAL:HA	1:A:279:LEU:HD12	2.03	0.40
2:B:103:LEU:HD23	2:B:103:LEU:C	2.46	0.40
3:C:156:ASP:HB3	3:C:187:LEU:HD13	2.03	0.40
1:A:247:GLU:HG2	1:A:402:TYR:OH	2.20	0.40
3:C:108:GLY:HA2	3:C:110:TYR:CE2	2.56	0.40
3:C:171:LEU:CD2	3:C:175:VAL:HG22	2.51	0.40
1:A:189:LEU:O	1:A:193:CYS:HB2	2.22	0.40
1:A:235:TYR:CE1	1:A:278:LEU:HG	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	431/469 (92%)	396 (92%)	35 (8%)	0	100	100
2	B	211/213 (99%)	192 (91%)	19 (9%)	0	100	100
3	C	224/226 (99%)	202 (90%)	22 (10%)	0	100	100
All	All	866/908 (95%)	790 (91%)	76 (9%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	282/420 (67%)	269 (95%)	13 (5%)	24	58
2	B	183/186 (98%)	172 (94%)	11 (6%)	17	50
3	C	179/196 (91%)	174 (97%)	5 (3%)	38	68
All	All	644/802 (80%)	615 (96%)	29 (4%)	24	58

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	100	VAL
1	A	128	GLU
1	A	304	ASN
1	A	340	MET
1	A	345	ILE
1	A	361	CYS
1	A	369	PHE
1	A	376	ARG
1	A	382	ILE
1	A	384	LEU
1	A	388	LEU
1	A	390	PHE
1	A	394	GLN
2	B	2	ILE
2	B	7	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	12	SER
2	B	14	SER
2	B	59	VAL
2	B	113	SER
2	B	121	ASP
2	B	157	ASN
2	B	179	THR
2	B	182	LYS
2	B	201	SER
3	C	56	ASP
3	C	125	SER
3	C	128	THR
3	C	213	LYS
3	C	218	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	112	GLN
1	A	240	ASN
1	A	1025	ASN
1	A	1048	GLN
1	A	410	GLN
2	B	91	HIS
3	C	183	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	97Y	A	503	-	39,40,40	1.98	9 (23%)	52,59,59	1.83	8 (15%)
4	NAG	A	501	1	14,14,15	0.44	0	17,19,21	0.69	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	97Y	A	503	-	-	4/30/42/42	0/4/4/4
4	NAG	A	501	1	-	2/6/23/26	0/1/1/1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	503	97Y	C34-N11	6.03	1.47	1.33
6	A	503	97Y	C22-N07	-4.52	1.32	1.38
6	A	503	97Y	C20-N08	3.52	1.46	1.39
6	A	503	97Y	C16-C14	-3.19	1.51	1.55
6	A	503	97Y	C15-C14	-3.04	1.51	1.55
6	A	503	97Y	C15-C12	-2.96	1.51	1.55
6	A	503	97Y	C12-C13	-2.79	1.48	1.53
6	A	503	97Y	C17-C13	2.77	1.55	1.52
6	A	503	97Y	C16-C12	-2.49	1.51	1.55

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	503	97Y	C17-C13-C12	-5.63	106.59	112.15
6	A	503	97Y	N10-C19-N07	4.96	121.57	114.75

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	503	97Y	N07-C23-N09	-4.95	110.33	112.45
6	A	503	97Y	C20-C27-N10	-4.49	120.36	124.07
6	A	503	97Y	C30-C14-C29	3.00	115.66	110.17
6	A	503	97Y	C27-N10-C19	2.86	122.09	117.41
6	A	503	97Y	C22-N07-C23	2.55	107.74	106.31
4	A	501	NAG	O5-C1-C2	-2.32	107.69	111.29
6	A	503	97Y	F02-C21-C18	-2.07	108.61	112.86

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	A	503	97Y	C15-C12-C13-C17
4	A	501	NAG	C8-C7-N2-C2
4	A	501	NAG	O7-C7-N2-C2
6	A	503	97Y	N11-C35-C36-C37
6	A	503	97Y	C22-C18-C21-F01
6	A	503	97Y	C22-C18-C21-F03

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand 97Y A 503



Ligand NAG A 501



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	437/469 (93%)	0.09	15 (3%) 48 30	124, 161, 211, 244	0
2	B	213/213 (100%)	-0.36	3 (1%) 73 54	80, 101, 122, 161	0
3	C	226/226 (100%)	-0.22	2 (0%) 81 64	78, 114, 161, 180	0
All	All	876/908 (96%)	-0.10	20 (2%) 61 41	78, 137, 198, 244	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1010	GLY	6.4
1	A	192	LEU	4.3
1	A	371	MET	3.7
1	A	1012	ILE	3.6
1	A	1007	THR	3.1
3	C	110	TYR	3.1
1	A	394	GLN	2.6
1	A	68	GLU	2.6
2	B	28	SER	2.4
1	A	257	PHE	2.3
1	A	370	VAL	2.3
1	A	1008	VAL	2.3
2	B	103	LEU	2.3
1	A	92	ALA	2.3
2	B	62	SER	2.2
1	A	97	GLN	2.2
3	C	207	THR	2.2
1	A	267	ARG	2.1
1	A	377	GLY	2.1
1	A	398	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	A	501	14/15	0.71	0.11	170,177,181,182	0
6	97Y	A	503	37/37	0.89	0.10	126,149,185,186	0
5	ZN	A	502	1/1	0.98	0.03	175,175,175,175	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around NAG A 501:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 97Y A 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.