



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:43 PM UTC

PDB ID : 4LO1 / pdb_00004lo1
Title : HA17-HA33-Gal
Authors : Lee, K.; Gu, S.; Jin, L.; Le, T.T.; Cheng, L.W.; Strotmeier, J.; Kruel, A.M.;
Yao, G.; Perry, K.; Rummel, A.; Jin, R.
Deposited on : 2013-07-12
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1898 (2.26-2.26)
Clashscore	190562	2005 (2.26-2.26)
Ramachandran outliers	187476	1965 (2.26-2.26)
Sidechain outliers	187428	1966 (2.26-2.26)
RSRZ outliers	180081	1898 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	296	4% 91% 6% .
1	B	296	9% 90% 5% .
2	C	147	3% 87% 10% ..

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6107 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HA-33.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	286	Total	C	N	O	S	5	1	0
			2334	1477	399	455	3			
1	B	285	Total	C	N	O	S	7	1	0
			2326	1473	397	453	3			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	294	PRO	-	expression tag	UNP Q45871
A	295	GLY	-	expression tag	UNP Q45871
A	296	SER	-	expression tag	UNP Q45871
A	297	ALA	-	expression tag	UNP Q45871
B	294	PRO	-	expression tag	UNP Q45871
B	295	GLY	-	expression tag	UNP Q45871
B	296	SER	-	expression tag	UNP Q45871
B	297	ALA	-	expression tag	UNP Q45871

- Molecule 2 is a protein called HA-17.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	144	Total	C	N	O	S	3	0	0
			1187	767	189	226	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	GLY	-	expression tag	UNP Q45878
C	1	PRO	-	expression tag	UNP Q45878

- Molecule 3 is beta-D-galactopyranose (CCD ID: GAL) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			12	6	6		
3	B	1	Total	C	O	0	0
			12	6	6		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	97	Total	O	0	0
			97	97		
4	C	64	Total	O	0	0
			64	64		
4	B	75	Total	O	0	0
			75	75		

- Molecule 1: HA-33



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	107.13Å 118.70Å 162.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.17 – 2.25 36.17 – 2.25	Depositor EDS
% Data completeness (in resolution range)	99.7 (36.17-2.25) 99.7 (36.17-2.25)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.81 (at 2.24Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.2_869)	Depositor
R, R_{free}	0.199 , 0.227 0.193 , 0.222	Depositor DCC
R_{free} test set	2481 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	41.4	Xtriage
Anisotropy	0.687	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 51.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6107	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/2393	0.66	0/3267
1	B	0.29	1/2385 (0.0%)	0.69	2/3256 (0.1%)
2	C	0.29	0/1217	0.68	0/1653
All	All	0.28	1/5995 (0.0%)	0.68	2/8176 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	81	ASN	C-N	5.70	1.41	1.33

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	113	ASN	CA-C-N	5.22	124.89	119.56
1	B	113	ASN	C-N-CA	5.22	124.89	119.56

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2334	0	2250	10	0
1	B	2326	0	2244	13	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	1187	0	1150	11	0
3	A	12	0	12	0	0
3	B	12	0	12	1	0
4	A	97	0	0	3	0
4	B	75	0	0	3	0
4	C	64	0	0	2	0
All	All	6107	0	5668	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:119:TYR:CD2	1:B:128:LYS:NZ	2.53	0.71
1:A:62:LYS:NZ	1:A:69:THR:O	2.24	0.68
1:B:139:LYS:NZ	4:B:407:HOH:O	2.24	0.68
1:A:65:ASP:O	4:A:451:HOH:O	2.17	0.57
2:C:100:LEU:HD23	2:C:111:ALA:HB2	1.87	0.55
2:C:94:VAL:HB	4:C:260:HOH:O	2.08	0.54
1:A:223:VAL:HG13	1:A:276:GLN:HA	1.91	0.53
1:A:122:THR:O	1:A:125:ARG:NE	2.37	0.53
1:A:186:ARG:HB2	1:A:207:ILE:HD13	1.92	0.52
1:B:196:GLU:CD	1:B:196:GLU:H	2.19	0.51
2:C:50:MET:HE2	2:C:53:ASN:ND2	2.28	0.49
2:C:6:THR:OG1	2:C:7:PHE:N	2.34	0.49
1:B:223:VAL:HG13	1:B:276:GLN:HA	1.94	0.47
1:A:196:GLU:H	1:A:196:GLU:CD	2.22	0.47
1:B:186:ARG:HB2	1:B:207:ILE:HD13	1.97	0.47
1:B:276:GLN:NE2	3:B:301:GAL:O6	2.38	0.46
2:C:93:ALA:HB3	2:C:96:THR:HB	1.97	0.46
4:A:445:HOH:O	2:C:106:ASN:HB3	2.16	0.45
1:A:245:ASP:O	4:A:464:HOH:O	2.21	0.44
2:C:122:LEU:HD23	2:C:122:LEU:HA	1.83	0.43
1:B:155:CYS:HA	1:B:293:PRO:HD3	2.00	0.43
1:B:63:SER:OG	4:B:427:HOH:O	2.21	0.43
1:A:9:ASN:O	1:A:9:ASN:OD1	2.37	0.43
1:A:207:ILE:HD12	1:A:207:ILE:HA	1.85	0.43
2:C:73:PHE:CE1	1:B:128:LYS:NZ	2.64	0.42
1:B:23:ASP:OD2	4:B:447:HOH:O	2.21	0.42
2:C:55:CYS:SG	2:C:102:LEU:HD13	2.61	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:60:LYS:NZ	1:A:87:ASP:OD2	2.52	0.41
2:C:75:PHE:CG	1:B:80:HIS:NE2	2.89	0.41
1:B:176:ASN:HA	1:B:224:ARG:HG2	2.03	0.41
2:C:50:MET:CE	2:C:53:ASN:HD21	2.33	0.40
4:C:216:HOH:O	1:B:80:HIS:HE1	2.05	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	285/296 (96%)	275 (96%)	10 (4%)	0	100	100
1	B	284/296 (96%)	272 (96%)	12 (4%)	0	100	100
2	C	142/147 (97%)	135 (95%)	7 (5%)	0	100	100
All	All	711/739 (96%)	682 (96%)	29 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	263/270 (97%)	258 (98%)	5 (2%)	50	61
1	B	262/270 (97%)	259 (99%)	3 (1%)	65	75

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	C	136/138 (99%)	134 (98%)	2 (2%)	57	68
All	All	661/678 (98%)	651 (98%)	10 (2%)	57	68

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	80	HIS
1	A	84	THR
1	A	175	LEU
1	A	223	VAL
1	A	292	ASN
2	C	26	ASN
2	C	122	LEU
1	B	186	ARG
1	B	223	VAL
1	B	292	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	67	HIS
1	A	152	ASN
1	A	236	GLN
1	A	276	GLN
1	A	289	ASN
2	C	138	ASN
1	B	115	ASN
1	B	134	ASN
1	B	152	ASN
1	B	202	GLN
1	B	236	GLN
1	B	276	GLN
1	B	289	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GAL	A	301	-	12,12,12	0.56	0	17,17,17	0.68	0
3	GAL	B	301	-	12,12,12	0.53	0	17,17,17	0.65	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GAL	A	301	-	-	1/2/22/22	0/1/1/1
3	GAL	B	301	-	-	2/2/22/22	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	301	GAL	O5-C5-C6-O6
3	B	301	GAL	C4-C5-C6-O6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	301	GAL	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	301	GAL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	286/296 (96%)	0.13	11 (3%) 44 44	26, 51, 83, 128	1 (0%)
1	B	285/296 (96%)	0.56	26 (9%) 15 14	29, 62, 130, 164	2 (0%)
2	C	144/147 (97%)	0.03	4 (2%) 55 55	28, 53, 76, 99	1 (0%)
All	All	715/739 (96%)	0.28	41 (5%) 29 27	26, 55, 118, 164	4 (0%)

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	146	ILE	5.8
1	B	230	ASP	4.7
1	B	294	PRO	4.7
1	B	231	GLN	4.2
1	A	9	ASN	4.0
1	A	231	GLN	3.5
2	C	3	VAL	3.4
1	B	227	SER	3.3
1	B	269	THR	3.2
1	B	128	LYS	3.1
1	B	261	ALA	3.1
1	B	10	SER	3.0
1	B	228	SER	3.0
1	A	208	LEU	3.0
1	A	269	THR	2.9
1	B	160	ILE	2.9
1	B	161	LEU	2.8
1	A	226	SER	2.7
1	B	232	ASN	2.7
1	B	66	ILE	2.6
1	B	184	TYR	2.6
1	B	216	ILE	2.6
1	A	294	PRO	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	217	PHE	2.4
1	B	218	SER	2.4
1	B	265	TYR	2.4
1	A	173	THR	2.4
2	C	94	VAL	2.4
1	A	245	ASP	2.3
1	A	184	TYR	2.3
1	B	67	HIS	2.3
1	B	244	SER	2.3
1	B	280	TYR	2.2
1	B	215	TRP	2.2
1	B	283	ASP	2.2
1	A	228	SER	2.2
1	B	245	ASP	2.1
1	B	275	ILE	2.1
2	C	55	CYS	2.1
1	B	164	ASN	2.1
1	A	217	PHE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GAL	B	301	12/12	0.63	0.14	89,96,101,107	0
3	GAL	A	301	12/12	0.80	0.11	60,69,75,76	0

6.5 Other polymers [i](#)

There are no such residues in this entry.