



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 12:41 PM UTC

PDB ID : 5LOU / pdb_00005lou
Title : human NUDT22
Authors : Carter, M.; Stenmark, P.
Deposited on : 2016-08-10
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	330	12% 75% 12% 14%
1	B	330	29% 82% 8% 9%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4825 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nucleoside diphosphate-linked moiety X motif 22.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	285	Total	C	N	O	S	0	0	0
			2167	1372	378	411	6			
1	B	301	Total	C	N	O	S	0	0	0
			2292	1445	405	434	8			

There are 54 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-26	MET	-	initiating methionine	UNP Q9BRQ3
A	-25	GLY	-	expression tag	UNP Q9BRQ3
A	-24	SER	-	expression tag	UNP Q9BRQ3
A	-23	SER	-	expression tag	UNP Q9BRQ3
A	-22	HIS	-	expression tag	UNP Q9BRQ3
A	-21	HIS	-	expression tag	UNP Q9BRQ3
A	-20	HIS	-	expression tag	UNP Q9BRQ3
A	-19	HIS	-	expression tag	UNP Q9BRQ3
A	-18	HIS	-	expression tag	UNP Q9BRQ3
A	-17	HIS	-	expression tag	UNP Q9BRQ3
A	-16	SER	-	expression tag	UNP Q9BRQ3
A	-15	SER	-	expression tag	UNP Q9BRQ3
A	-14	GLY	-	expression tag	UNP Q9BRQ3
A	-13	LEU	-	expression tag	UNP Q9BRQ3
A	-12	VAL	-	expression tag	UNP Q9BRQ3
A	-11	PRO	-	expression tag	UNP Q9BRQ3
A	-10	ARG	-	expression tag	UNP Q9BRQ3
A	-9	GLY	-	expression tag	UNP Q9BRQ3
A	-8	SER	-	expression tag	UNP Q9BRQ3
A	-7	HIS	-	expression tag	UNP Q9BRQ3
A	-6	MET	-	expression tag	UNP Q9BRQ3
A	-5	SER	-	expression tag	UNP Q9BRQ3
A	-4	CYS	-	expression tag	UNP Q9BRQ3
A	-3	PRO	-	expression tag	UNP Q9BRQ3
A	-2	VAL	-	expression tag	UNP Q9BRQ3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLN	-	expression tag	UNP Q9BRQ3
A	0	THR	-	expression tag	UNP Q9BRQ3
B	-26	MET	-	initiating methionine	UNP Q9BRQ3
B	-25	GLY	-	expression tag	UNP Q9BRQ3
B	-24	SER	-	expression tag	UNP Q9BRQ3
B	-23	SER	-	expression tag	UNP Q9BRQ3
B	-22	HIS	-	expression tag	UNP Q9BRQ3
B	-21	HIS	-	expression tag	UNP Q9BRQ3
B	-20	HIS	-	expression tag	UNP Q9BRQ3
B	-19	HIS	-	expression tag	UNP Q9BRQ3
B	-18	HIS	-	expression tag	UNP Q9BRQ3
B	-17	HIS	-	expression tag	UNP Q9BRQ3
B	-16	SER	-	expression tag	UNP Q9BRQ3
B	-15	SER	-	expression tag	UNP Q9BRQ3
B	-14	GLY	-	expression tag	UNP Q9BRQ3
B	-13	LEU	-	expression tag	UNP Q9BRQ3
B	-12	VAL	-	expression tag	UNP Q9BRQ3
B	-11	PRO	-	expression tag	UNP Q9BRQ3
B	-10	ARG	-	expression tag	UNP Q9BRQ3
B	-9	GLY	-	expression tag	UNP Q9BRQ3
B	-8	SER	-	expression tag	UNP Q9BRQ3
B	-7	HIS	-	expression tag	UNP Q9BRQ3
B	-6	MET	-	expression tag	UNP Q9BRQ3
B	-5	SER	-	expression tag	UNP Q9BRQ3
B	-4	CYS	-	expression tag	UNP Q9BRQ3
B	-3	PRO	-	expression tag	UNP Q9BRQ3
B	-2	VAL	-	expression tag	UNP Q9BRQ3
B	-1	GLN	-	expression tag	UNP Q9BRQ3
B	0	THR	-	expression tag	UNP Q9BRQ3

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	182	Total	O	0	0
			182	182		
3	B	174	Total	O	0	0
			174	174		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nucleoside diphosphate-linked moiety X motif 22



- Molecule 1: Nucleoside diphosphate-linked moiety X motif 22



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	52.21Å 46.87Å 119.61Å 90.00° 93.92° 90.00°	Depositor
Resolution (Å)	48.98 – 1.80 48.98 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.98-1.80) 99.9 (48.98-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.10 (at 1.79Å)	Xtriage
Refinement program	PHENIX (1.11.1 _2575: ???)	Depositor
R, R_{free}	0.199 , 0.242 0.235 , 0.269	Depositor DCC
R_{free} test set	2694 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	27.9	Xtriage
Anisotropy	0.388	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 43.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4825	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/2218	0.77	2/3026 (0.1%)
1	B	0.45	0/2347	0.76	1/3202 (0.0%)
All	All	0.46	0/4565	0.76	3/6228 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	164	PRO	N-CA-CB	7.19	110.80	103.25
1	A	242	GLY	CA-C-N	5.01	124.79	119.28
1	A	242	GLY	C-N-CA	5.01	124.79	119.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2167	0	2140	37	0
1	B	2292	0	2246	30	0
2	A	5	0	0	0	0
2	B	5	0	0	1	0
3	A	182	0	0	14	0
3	B	174	0	0	8	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	4825	0	4386	67	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1:MET:HE1	1:A:263:LEU:CD1	1.30	1.59
1:A:1:MET:CE	1:A:263:LEU:HD13	1.42	1.46
1:B:263:LEU:CD2	1:B:268:TRP:CE2	2.00	1.43
1:B:263:LEU:HD21	1:B:268:TRP:CZ2	1.50	1.42
1:B:263:LEU:HD23	1:B:268:TRP:CE2	1.62	1.28
1:B:263:LEU:HD23	1:B:268:TRP:CD2	1.69	1.25
1:A:1:MET:CE	1:A:263:LEU:CD1	2.04	1.25
1:B:263:LEU:CD2	1:B:268:TRP:CZ2	2.18	1.21
1:A:1:MET:HE3	1:A:263:LEU:HD13	1.20	1.15
1:A:1:MET:HE1	1:A:263:LEU:HD12	1.37	1.04
1:A:263:LEU:CD2	1:A:268:TRP:CE3	2.44	0.93
1:A:1:MET:HE1	1:A:263:LEU:HD11	1.48	0.91
1:A:263:LEU:HD23	1:A:268:TRP:CE3	1.56	0.89
1:B:263:LEU:HD21	1:B:268:TRP:CH2	2.14	0.82
1:B:263:LEU:HD22	1:B:268:TRP:CE2	2.17	0.77
1:A:46:GLU:OE1	3:A:501:HOH:O	2.05	0.74
1:B:111:GLY:O	3:B:501:HOH:O	2.05	0.72
1:B:263:LEU:HD23	1:B:268:TRP:CE3	2.27	0.69
1:B:-10:ARG:NH1	3:B:507:HOH:O	2.26	0.68
1:A:75:GLY:O	3:A:502:HOH:O	2.11	0.68
1:B:247:GLU:H	1:B:247:GLU:CD	2.00	0.68
1:B:263:LEU:CD2	1:B:268:TRP:CH2	2.78	0.65
1:A:159:PRO:O	3:A:503:HOH:O	2.14	0.65
1:A:88:ARG:NE	3:A:512:HOH:O	2.31	0.64
1:A:1:MET:CE	1:A:263:LEU:HD11	2.18	0.63
1:B:263:LEU:CD2	1:B:268:TRP:NE1	2.60	0.63
1:A:145:GLU:OE1	3:A:504:HOH:O	2.17	0.59
1:A:101:TRP:O	1:A:105:GLN:HG3	2.03	0.58
1:B:263:LEU:HD22	1:B:268:TRP:NE1	2.19	0.57
1:B:285:VAL:HG12	1:B:286:GLN:H	1.71	0.55
1:B:38:ASP:HA	3:B:515:HOH:O	2.06	0.54
1:B:168:PRO:O	3:B:502:HOH:O	2.19	0.53
1:A:298:ALA:O	3:A:505:HOH:O	2.19	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:88:ARG:NE	3:B:518:HOH:O	2.40	0.52
1:A:260:GLN:HG3	3:A:558:HOH:O	2.09	0.52
1:B:-7:HIS:NE2	1:B:119:ASP:OD2	2.44	0.51
1:A:257:GLN:H	1:A:257:GLN:CD	2.20	0.50
1:A:59:PRO:HB2	1:A:159:PRO:HG2	1.93	0.50
1:A:46:GLU:O	1:A:50:LYS:HD3	2.12	0.50
1:A:186:VAL:HG11	1:A:223:PHE:CD2	2.47	0.50
1:A:39:GLU:HA	1:A:39:GLU:OE1	2.12	0.49
1:A:260:GLN:CG	3:A:558:HOH:O	2.60	0.49
1:A:255:GLU:OE1	3:A:506:HOH:O	2.20	0.49
1:B:71:ILE:C	1:B:73:SER:H	2.23	0.47
1:A:105:GLN:NE2	3:A:519:HOH:O	2.47	0.47
1:A:176:GLN:OE1	3:A:509:HOH:O	2.21	0.46
1:A:255:GLU:HB3	1:A:257:GLN:OE1	2.15	0.46
1:B:164:PRO:HA	1:B:165:GLY:HA2	1.55	0.46
1:B:25:LEU:HD13	1:B:82:LEU:HB2	1.99	0.45
1:B:34:LEU:HD12	1:B:35:PRO:HD2	1.99	0.45
1:B:74:ARG:N	1:B:74:ARG:HD3	2.32	0.45
1:B:158:GLU:HG2	3:B:567:HOH:O	2.17	0.44
1:A:173:LEU:HA	3:A:635:HOH:O	2.19	0.43
1:A:28:ALA:O	1:A:32:ARG:NH2	2.51	0.42
1:A:202:SER:OG	3:A:508:HOH:O	2.21	0.42
1:B:218:ARG:NH2	2:B:401:PO4:O3	2.51	0.42
1:A:34:LEU:O	1:A:36:GLY:HA2	2.20	0.42
1:A:231:SER:OG	3:A:507:HOH:O	2.21	0.42
1:A:133:PHE:HB3	1:A:253:PHE:HB3	2.02	0.41
1:A:150:VAL:HB	1:A:271:LEU:HD23	2.01	0.41
1:A:258:ASN:O	1:A:261:ARG:HG2	2.19	0.41
1:B:103:ARG:HD2	1:B:113:THR:HB	2.01	0.41
1:B:-10:ARG:HD3	3:B:606:HOH:O	2.20	0.41
1:A:140:SER:O	1:A:143:VAL:HG22	2.21	0.41
1:A:104:GLN:O	1:A:108:THR:HG23	2.21	0.41
1:B:163:CYS:C	3:B:514:HOH:O	2.63	0.40
1:B:-12:VAL:HA	1:B:-11:PRO:HD2	1.94	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:504:HOH:O	3:B:507:HOH:O[1_565]	2.14	0.06

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	281/330 (85%)	273 (97%)	8 (3%)	0	100	100
1	B	299/330 (91%)	286 (96%)	11 (4%)	2 (1%)	18	8
All	All	580/660 (88%)	559 (96%)	19 (3%)	2 (0%)	36	25

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	164	PRO
1	B	285	VAL

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	76/267 (28%)	75 (99%)	1 (1%)	61	54
1	B	187/267 (70%)	183 (98%)	4 (2%)	47	36
All	All	263/534 (49%)	258 (98%)	5 (2%)	50	41

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	141	ARG
1	B	71	ILE
1	B	104	GLN
1	B	244	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	261	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	104	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	B	401	-	4,4,4	0.98	0	6,6,6	0.46	0
2	PO4	A	401	-	4,4,4	0.98	0	6,6,6	0.45	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	PO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	285/330 (86%)	0.89	40 (14%) 6 5	20, 35, 70, 89	0
1	B	301/330 (91%)	1.44	97 (32%) 1 1	23, 41, 81, 110	0
All	All	586/660 (88%)	1.17	137 (23%) 2 1	20, 38, 76, 110	0

All (137) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	-11	PRO	7.7
1	A	101	TRP	7.0
1	A	263	LEU	7.0
1	A	298	ALA	6.2
1	A	260	GLN	5.8
1	A	173	LEU	5.8
1	A	71	ILE	5.8
1	B	263	LEU	5.4
1	B	-12	VAL	4.6
1	B	71	ILE	4.6
1	B	229	LEU	4.6
1	B	35	PRO	4.6
1	B	230	THR	4.5
1	B	199	LEU	4.4
1	B	260	GLN	4.1
1	B	72	GLY	4.0
1	B	101	TRP	4.0
1	A	95	TRP	3.9
1	B	34	LEU	3.9
1	B	243	PRO	3.9
1	B	130	ALA	3.8
1	A	1	MET	3.7
1	A	100	ALA	3.7
1	B	36	GLY	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	1	MET	3.7
1	B	163	CYS	3.7
1	A	36	GLY	3.6
1	B	-10	ARG	3.6
1	B	-9	GLY	3.5
1	B	285	VAL	3.5
1	B	226	GLN	3.5
1	B	-4	CYS	3.4
1	A	74	ARG	3.4
1	B	69	ALA	3.4
1	B	240	SER	3.4
1	B	-3	PRO	3.4
1	B	162	LEU	3.4
1	B	13	GLY	3.3
1	A	264	GLU	3.3
1	B	247	GLU	3.2
1	A	70	PRO	3.2
1	B	197	PRO	3.2
1	B	68	LEU	3.2
1	B	-7	HIS	3.2
1	A	104	GLN	3.1
1	B	228	SER	3.1
1	A	35	PRO	3.1
1	A	69	ALA	3.1
1	B	195	ASN	3.1
1	B	198	LEU	3.0
1	B	224	TYR	3.0
1	B	70	PRO	3.0
1	A	56	PHE	3.0
1	B	14	GLY	2.9
1	B	164	PRO	2.9
1	B	245	ALA	2.9
1	A	105	GLN	2.9
1	A	257	GLN	2.9
1	A	297	PRO	2.9
1	B	193	GLU	2.8
1	B	110	TRP	2.8
1	B	21	ILE	2.8
1	B	205	LEU	2.8
1	B	15	GLY	2.8
1	B	165	GLY	2.8
1	B	200	THR	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	166	GLY	2.8
1	A	54	TRP	2.8
1	B	227	CYS	2.8
1	B	129	THR	2.7
1	A	261	ARG	2.7
1	B	128	ALA	2.7
1	A	141	ARG	2.7
1	B	202	SER	2.7
1	A	39	GLU	2.7
1	B	40	ALA	2.6
1	B	244	GLU	2.6
1	B	73	SER	2.6
1	B	167	SER	2.6
1	B	246	HIS	2.6
1	B	241	GLY	2.6
1	A	262	LEU	2.6
1	B	201	LEU	2.6
1	A	108	THR	2.6
1	B	131	ASP	2.5
1	A	174	ALA	2.5
1	B	74	ARG	2.5
1	B	286	GLN	2.5
1	B	234	VAL	2.5
1	B	33	PRO	2.5
1	B	-6	MET	2.5
1	A	72	GLY	2.4
1	B	237	HIS	2.4
1	A	265	THR	2.4
1	B	104	GLN	2.4
1	B	196	LEU	2.4
1	B	239	LEU	2.4
1	B	-2	VAL	2.4
1	B	232	GLU	2.4
1	B	75	GLY	2.4
1	A	55	LEU	2.4
1	B	256	THR	2.4
1	B	177	LEU	2.3
1	B	77	GLN	2.3
1	B	-8	SER	2.3
1	B	225	VAL	2.3
1	B	191	CYS	2.3
1	B	182	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	230	THR	2.3
1	B	257	GLN	2.3
1	A	51	ALA	2.3
1	B	132	ASP	2.3
1	B	18	GLN	2.3
1	B	16	LEU	2.3
1	B	12	PRO	2.2
1	B	19	GLU	2.2
1	B	37	GLY	2.2
1	A	249	THR	2.2
1	B	100	ALA	2.2
1	B	186	VAL	2.2
1	B	231	SER	2.2
1	B	242	GLY	2.2
1	B	133	PHE	2.2
1	A	143	VAL	2.2
1	A	140	SER	2.2
1	B	-14	GLY	2.2
1	B	204	PRO	2.2
1	A	266	GLU	2.1
1	B	233	GLN	2.1
1	B	127	LEU	2.1
1	B	190	ILE	2.1
1	B	-13	LEU	2.1
1	B	264	GLU	2.0
1	A	37	GLY	2.0
1	A	34	LEU	2.0
1	A	50	LYS	2.0
1	A	289	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PO4	A	401	5/5	0.92	0.10	44,46,49,51	0
2	PO4	B	401	5/5	0.94	0.09	44,47,51,55	0

6.5 Other polymers [i](#)

There are no such residues in this entry.