



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 07:57 AM UTC

PDB ID : 1LPZ / pdb_00001lpz
Title : CRYSTAL STRUCTURE OF FXA IN COMPLEX WITH 41.
Authors : Schreuder, H.A.; Brachvogel, V.; Liesum, A.
Deposited on : 2002-05-08
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	134	 28% 10% • 60%
2	B	254	 67% 21% • 8%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2571 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Blood coagulation factor Xa.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	53	Total	C	N	O	S	0	0	0
			395	237	70	81	7			


- Molecule 2 is a protein called Blood coagulation factor Xa.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	234	Total	C	N	O	S	0	0	0
			1852	1166	323	349	14			

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Ca	0	0
			1	1		

- Molecule 4 is 1-(3-carbamimidoylbenzyl)-N-(3,5-dichlorobenzyl)-4-methyl-1H-indole-2-carboxamide (CCD ID: CMB) (formula: C₂₅H₂₂Cl₂N₄O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	Cl	N	O	0	0
			32	25	2	4	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	60	Total	O	0	0
			60	60		
5	B	231	Total	O	0	0
			231	231		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Blood coagulation factor Xa

Chain A: 

GLU GLU MET LYS LYS GLY HIS LEU GLU ARG GLU MET GLU THR CYS TYR GLU ALA ARG GLU VAL PHE GLU ASP SER ASP LYS THR ASN GLU PHE TRP ASN LYS TYR LYS ASP GLY ASP GLN CYS GLU THR SER PRO CYS GLN ASN GLN GLY LYS CYS LYS ASP GLY LEU

GLY TYR THR CYS THR CYS LEU GLU PHE THR R1A K1B L1C C1 S2 L3 G6 H13 E14 E15 Q16 V19 T39 G40 P41 Y42 P43 C44 L49 E50 ARG

- Molecule 2: Blood coagulation factor Xa

Chain B: 

I16 E21 E26 L32 N35 E36 E37 N38 E39 G40 Y61 H57 C58 L59 Y60 Q61 A61A K62 R63 N72 T73 E76 E86 V87 H91 N92 R93 F94 T95 K109 R115 M116 C122 L123 S130 T131 L131A Q133 G136 H145 E147 K148

T153 M157 N166 S167 L170 S171 S172 I175 I176 T177 Q178 A183 D185A T185B K186 Q187 E188 D189 A190 G191 Q192 S195 G196 G197 P198 H199 V200 K204 D205 V209 C220 K223 C223A K224 Y228 S241 T244 ARG GLY LEU PRO LYS ALA LYS SER HIS

ALA PRO GLU VAL ILE THR SER SER LYS

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	56.30Å 72.00Å 78.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.40 8.00 – 2.41	Depositor EDS
% Data completeness (in resolution range)	91.1 (8.00-2.40) 93.9 (8.00-2.41)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.83 (at 2.42Å)	Xtriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.179 , 0.279 0.168 , 0.257	Depositor DCC
R_{free} test set	1234 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å ²)	30.5	Xtriage
Anisotropy	0.562	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 109.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2571	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CMB, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	0/401	1.00	2/541 (0.4%)
2	B	0.57	0/1890	0.95	12/2546 (0.5%)
All	All	0.57	0/2291	0.96	14/3087 (0.5%)

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	40	GLY	CA-C-N	7.54	127.07	118.85
1	A	40	GLY	C-N-CA	7.54	127.07	118.85
2	B	199	HIS	N-CA-C	-6.68	96.32	108.02
2	B	51	TYR	N-CA-C	6.45	119.46	109.07
2	B	136	GLY	N-CA-C	-6.25	101.80	111.12
2	B	190	ALA	N-CA-C	-5.87	103.19	110.41
2	B	123	LEU	N-CA-C	-5.76	102.52	109.72
2	B	172	SER	N-CA-C	5.51	118.91	110.20
2	B	175	ILE	N-CA-C	5.39	116.65	109.37
2	B	133	GLN	N-CA-C	-5.31	103.88	110.41
2	B	197	GLY	CA-C-N	5.28	126.09	120.13
2	B	197	GLY	C-N-CA	5.28	126.09	120.13
2	B	26	GLU	N-CA-C	5.03	117.88	111.69
2	B	153	THR	N-CA-C	-5.02	107.00	113.23

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	395	0	365	10	0
2	B	1852	0	1808	34	0
3	B	1	0	0	0	0
4	B	32	0	21	1	0
5	A	60	0	0	5	0
5	B	231	0	0	7	0
All	All	2571	0	2194	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:185(B):THR:HG22	2:B:223:LYS:HG2	1.65	0.77
2:B:157:MET:SD	5:B:511:HOH:O	2.54	0.67
2:B:166:ASN:HB2	5:B:394:HOH:O	1.96	0.65
2:B:86:GLU:HB2	2:B:109:LYS:HG3	1.80	0.62
2:B:220:CYS:SG	4:B:301:CMB:HC2	2.39	0.62
2:B:204:LYS:O	2:B:205:ASP:HB2	2.03	0.59
2:B:175:ILE:HD13	2:B:177:THR:HG22	1.90	0.53
2:B:185(B):THR:HG22	2:B:223:LYS:CG	2.37	0.52
2:B:177:THR:HB	5:B:466:HOH:O	2.09	0.52
1:A:1(C):LEU:HB3	1:A:19:VAL:CG2	2.41	0.50
1:A:44:CYS:C	2:B:122:CYS:SG	2.95	0.50
1:A:1(C):LEU:HB2	1:A:14:GLU:HG2	1.93	0.50
2:B:91:HIS:CE1	2:B:93:ARG:HB2	2.47	0.50
2:B:72:ASN:HB3	5:B:473:HOH:O	2.11	0.50
2:B:35:ASN:OD1	2:B:37:GLU:HG2	2.12	0.49
1:A:1(C):LEU:HB3	1:A:19:VAL:HG23	1.95	0.49
2:B:185(A):ASP:OD1	2:B:223(A):GLY:HA2	2.13	0.49
2:B:200:VAL:HG12	2:B:209:VAL:HA	1.95	0.48
2:B:185(B):THR:HA	2:B:223:LYS:HG3	1.96	0.48
1:A:1(B):LYS:O	1:A:3:LEU:HB3	2.14	0.47
2:B:32:LEU:HD11	2:B:40:GLY:HA3	1.97	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:192:GLN:HA	2:B:192:GLN:OE1	2.15	0.47
2:B:178:GLN:NE2	5:B:466:HOH:O	2.47	0.47
1:A:13:HIS:HB3	5:A:85:HOH:O	2.15	0.46
2:B:131(A):LEU:O	2:B:133:GLN:HG2	2.16	0.46
2:B:57:HIS:CE1	2:B:195:SER:HB3	2.51	0.45
2:B:183:ALA:HB3	2:B:228:TYR:CE1	2.52	0.45
2:B:72:ASN:C	2:B:72:ASN:HD22	2.23	0.44
2:B:35:ASN:OD1	2:B:39:GLU:HG3	2.17	0.44
2:B:166:ASN:O	2:B:170:LEU:HG	2.18	0.43
1:A:6:GLY:HA2	5:A:70:HOH:O	2.18	0.43
2:B:145:HIS:HB2	5:B:402:HOH:O	2.18	0.43
2:B:115:ARG:HD3	2:B:116:MET:O	2.19	0.42
1:A:43:PRO:HB3	5:A:79:HOH:O	2.19	0.42
1:A:2:SER:HB2	5:A:95:HOH:O	2.20	0.42
2:B:148:LYS:HG3	5:B:431:HOH:O	2.20	0.41
2:B:186:LYS:HG2	2:B:188:GLU:HG3	2.01	0.41
2:B:61:GLN:H	2:B:61:GLN:HG2	1.63	0.41
2:B:86:GLU:HG2	2:B:87:VAL:HG23	2.02	0.41
2:B:224:LYS:HA	2:B:224:LYS:HD3	1.57	0.41
2:B:58:CYS:C	2:B:60:TYR:H	2.28	0.41
2:B:72:ASN:HD22	2:B:73:THR:N	2.19	0.40
1:A:42:TYR:HB2	5:A:98:HOH:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	51/134 (38%)	45 (88%)	5 (10%)	1 (2%)	6	7
2	B	232/254 (91%)	221 (95%)	11 (5%)	0	100	100
All	All	283/388 (73%)	266 (94%)	16 (6%)	1 (0%)	30	43

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1(C)	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	45/118 (38%)	42 (93%)	3 (7%)	15	26
2	B	199/216 (92%)	181 (91%)	18 (9%)	9	15
All	All	244/334 (73%)	223 (91%)	21 (9%)	10	16

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	GLN
1	A	39	THR
1	A	49	LEU
2	B	21	GLU
2	B	37	GLU
2	B	39	GLU
2	B	61	GLN
2	B	62	LYS
2	B	63	ARG
2	B	76	GLU
2	B	95	THR
2	B	115	ARG
2	B	130	SER
2	B	148	LYS
2	B	167	SER
2	B	175	ILE
2	B	178	GLN
2	B	209	VAL
2	B	223	LYS
2	B	241	SER
2	B	244	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	16	GLN
2	B	38	ASN
2	B	72	ASN
2	B	178	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	CMB	B	301	-	35,35,35	1.61	5 (14%)	43,50,50	1.81	11 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	CMB	B	301	-	-	0/17/17/17	0/4/4/4

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	301	CMB	C9-C8	4.22	1.45	1.37
4	B	301	CMB	C4-C5	3.66	1.46	1.41
4	B	301	CMB	C8-C11	-3.55	1.42	1.49
4	B	301	CMB	C2-C3	2.13	1.42	1.39
4	B	301	CMB	C30-C29	2.03	1.41	1.38

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	301	CMB	C4-C9-C8	-6.00	102.77	107.50
4	B	301	CMB	C17-C18-C25	-3.29	114.08	120.27
4	B	301	CMB	C9-C8-N7	3.04	111.13	109.42
4	B	301	CMB	C3-N7-C8	-2.71	106.45	108.44
4	B	301	CMB	C29-C28-C27	2.67	120.63	117.43
4	B	301	CMB	C25-C24-C31	-2.61	115.71	120.25
4	B	301	CMB	C3-C4-C9	2.44	110.42	107.56
4	B	301	CMB	C28-C27-C26	-2.34	118.79	121.64
4	B	301	CMB	C17-C18-C21	2.23	124.85	120.75
4	B	301	CMB	C27-C26-C14	2.10	121.86	119.71
4	B	301	CMB	C9-C8-C11	-2.10	124.25	129.48

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	301	CMB	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	53/134 (39%)	-0.46	0 100 100	13, 27, 72, 81	0
2	B	234/254 (92%)	-0.82	0 100 100	5, 23, 54, 78	0
All	All	287/388 (73%)	-0.75	0 100 100	5, 24, 58, 81	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CA	B	1	1/1	0.91	0.04	30,30,30,30	0
4	CMB	B	301	32/32	0.94	0.06	17,44,48,49	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around CMB B 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.