



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 08:14 PM UTC

PDB ID : 3LP2 / pdb_00003lp2
Title : HIV-1 reverse transcriptase with inhibitor
Authors : Yan, Y.; Munshi, S.K.; Prasad, G.S.; Su, H.P.
Deposited on : 2010-02-04
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	563	4% 77% 18% ••
2	B	443	4% 74% 15% • 10%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7913 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Reverse transcriptase/ribonuclease H.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	554	Total	C	N	O	S	0	0	0
			4508	2913	754	833	8			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	MET	-	expression tag	UNP P04585
A	-1	ASN	-	expression tag	UNP P04585
A	0	SER	-	expression tag	UNP P04585


- Molecule 2 is a protein called p51 RT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	400	Total	C	N	O	S	0	0	0
			3311	2155	548	602	6			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	MET	-	expression tag	UNP P04585
B	-1	ASN	-	expression tag	UNP P04585
B	0	SER	-	expression tag	UNP P04585

- Molecule 3 is 3-[4-(diethylamino)phenoxy]-6-(ethoxycarbonyl)-5,8-dihydroxy-7-oxo-7,8-dihydro-1,8-naphthyridin-1-ium (CCD ID: LP9) (formula: C₂₁H₂₃N₃O₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			30	21	3	6		

- Molecule 4 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	2	Total Mn 2 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	29	Total O 29 29	0	0
5	B	33	Total O 33 33	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Reverse transcriptase/ribonuclease H



• Molecule 2: p51 RT



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	117.51Å 154.78Å 155.79Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.80 50.00 – 2.80	Depositor EDS
% Data completeness (in resolution range)	97.0 (50.00-2.80) 97.0 (50.00-2.80)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.04 (at 2.81Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.249 , 0.296 0.245 , 0.290	Depositor DCC
R_{free} test set	1728 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	68.3	Xtriage
Anisotropy	0.073	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 31.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7913	wwPDB-VP
Average B, all atoms (Å ²)	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MN, LP9

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/4624	0.80	4/6285 (0.1%)
2	B	0.45	0/3404	0.81	3/4627 (0.1%)
All	All	0.46	0/8028	0.81	7/10912 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	320	ASP	CA-C-N	6.08	125.96	119.28
1	A	320	ASP	C-N-CA	6.08	125.96	119.28
2	B	139	THR	CA-C-N	-5.25	115.17	120.21
2	B	139	THR	C-N-CA	-5.25	115.17	120.21
1	A	224	GLU	CA-C-N	5.06	123.30	119.66
1	A	224	GLU	C-N-CA	5.06	123.30	119.66
2	B	348	ASN	N-CA-C	5.01	116.91	109.25

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4508	0	4549	49	0
2	B	3311	0	3326	34	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	30	0	22	1	0
4	A	2	0	0	0	0
5	A	29	0	0	1	0
5	B	33	0	0	0	0
All	All	7913	0	7897	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:337:TRP:HE1	2:B:367:GLN:HE21	1.36	0.74
1:A:458:VAL:HG22	1:A:548:VAL:HG22	1.69	0.73
2:B:388:LYS:HE2	2:B:415:GLU:HG3	1.74	0.70
2:B:388:LYS:HD2	2:B:413:GLU:HB3	1.75	0.69
1:A:235:HIS:HB3	1:A:236:PRO:HD2	1.77	0.67
2:B:209:LEU:HB3	2:B:214:LEU:HB2	1.77	0.66
1:A:175:ASN:HB3	1:A:178:ILE:HD12	1.78	0.64
1:A:447:ASN:OD1	1:A:449:GLU:HG2	1.97	0.64
2:B:13:LYS:HE3	2:B:85:GLN:HB3	1.82	0.61
1:A:199:ARG:HE	1:A:219:LYS:HE3	1.68	0.58
2:B:365:VAL:O	2:B:369:THR:HG23	2.05	0.57
1:A:181:TYR:HB2	1:A:188:TYR:HB3	1.86	0.57
2:B:209:LEU:HD12	2:B:214:LEU:HD12	1.86	0.56
1:A:60:VAL:HG12	1:A:75:VAL:HG22	1.87	0.56
1:A:125:ARG:HD3	1:A:147:ASN:HD22	1.71	0.56
2:B:169:GLU:HB3	2:B:170:PRO:HD3	1.89	0.55
2:B:253:THR:HG22	2:B:256:ASP:CG	2.32	0.54
1:A:501:TYR:O	1:A:505:ILE:HD12	2.09	0.53
2:B:116:PHE:CD1	2:B:116:PHE:C	2.86	0.53
1:A:365:VAL:O	1:A:369:THR:HG23	2.08	0.53
2:B:324:ASP:O	2:B:343:GLN:HG2	2.08	0.53
1:A:406:TRP:CH2	2:B:418:ASN:HA	2.45	0.52
2:B:195:ILE:O	2:B:199:ARG:HG3	2.10	0.51
1:A:169:GLU:HB3	1:A:170:PRO:HD3	1.93	0.51
1:A:430:GLU:CD	1:A:530:LYS:HE3	2.36	0.51
2:B:396:GLU:O	2:B:400:THR:HG23	2.10	0.51
1:A:182:GLN:HB3	1:A:187:LEU:HD23	1.93	0.50
1:A:364:ASP:HB3	1:A:423:VAL:HG13	1.94	0.49
1:A:90:VAL:O	1:A:161:GLN:NE2	2.45	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:38:CYS:SG	1:A:132:ILE:HD11	2.52	0.49
2:B:373:GLN:NE2	2:B:407:GLN:H	2.11	0.48
2:B:13:LYS:CE	2:B:85:GLN:HB3	2.43	0.48
1:A:46:LYS:HE3	1:A:116:PHE:HB3	1.95	0.48
2:B:47:ILE:HD12	2:B:144:TYR:CD1	2.50	0.47
1:A:544:GLY:HA2	2:B:286:THR:HG22	1.97	0.47
1:A:3:SER:OG	1:A:5:ILE:HG23	2.14	0.46
2:B:168:LEU:HD22	2:B:180:ILE:HD13	1.98	0.46
1:A:547:GLN:O	1:A:550:LYS:HG2	2.16	0.46
2:B:42:GLU:C	2:B:44:GLU:H	2.24	0.45
1:A:111:VAL:HG21	1:A:164:MET:HE1	1.98	0.45
1:A:5:ILE:HG22	1:A:212:TRP:CE3	2.52	0.45
1:A:474:ASN:O	1:A:478:GLU:HG2	2.17	0.45
2:B:275:LYS:H	2:B:306:ASN:HD21	1.63	0.45
1:A:100:LEU:O	1:A:318:TYR:HB3	2.16	0.45
2:B:373:GLN:HE22	2:B:407:GLN:H	1.64	0.45
2:B:373:GLN:O	2:B:377:THR:CG2	2.65	0.45
1:A:227:PHE:HB3	3:A:601:LP9:H	2.00	0.44
1:A:519:ASN:HA	1:A:522:ILE:HD12	1.98	0.44
2:B:157:PRO:HG3	2:B:184:MET:HA	2.00	0.44
2:B:86:ASP:O	2:B:91:GLN:HB2	2.17	0.44
1:A:0:SER:HA	1:A:1:PRO:HD3	1.89	0.43
1:A:372:VAL:HG11	1:A:411:ILE:HG23	1.98	0.43
1:A:460:ASN:ND2	2:B:288:ALA:HB2	2.33	0.43
2:B:344:GLU:O	2:B:347:LYS:HB2	2.18	0.43
1:A:545:ASN:O	1:A:549:ASP:HB2	2.17	0.43
2:B:317:VAL:HG12	2:B:349:LEU:HD13	2.00	0.43
1:A:412:PRO:O	1:A:413:GLU:C	2.62	0.43
1:A:303:LEU:O	1:A:307:ARG:HG3	2.18	0.43
1:A:312:GLU:HA	1:A:313:PRO:HD3	1.90	0.43
2:B:293:ILE:O	2:B:293:ILE:HG13	2.19	0.43
2:B:337:TRP:NE1	2:B:367:GLN:HE21	2.09	0.43
1:A:21:VAL:HG13	1:A:59:PRO:HD3	2.00	0.42
2:B:388:LYS:CE	2:B:415:GLU:HG3	2.47	0.42
1:A:255:ASN:HB2	1:A:289:LEU:HD22	2.01	0.42
1:A:180:ILE:HG12	1:A:189:VAL:HG13	2.01	0.42
1:A:442:VAL:HB	1:A:481:ALA:HB1	2.00	0.42
2:B:58:THR:HG23	2:B:76:ASP:O	2.18	0.42
1:A:320:ASP:C	1:A:320:ASP:OD1	2.63	0.42
1:A:354:TYR:HD1	1:A:374:LYS:HD2	1.85	0.42
2:B:203:GLU:CD	2:B:206:ARG:HH12	2.27	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:297:GLU:O	1:A:301:LEU:HD12	2.20	0.41
1:A:373:GLN:HG3	5:A:562:HOH:O	2.21	0.41
1:A:344:GLU:O	1:A:345:PRO:C	2.63	0.41
1:A:460:ASN:HD22	2:B:288:ALA:HB2	1.86	0.41
1:A:335:GLY:O	1:A:355:ALA:HA	2.20	0.40
1:A:138:GLU:HB3	1:A:139:THR:H	1.56	0.40
1:A:520:GLN:O	1:A:523:GLU:HG2	2.21	0.40
1:A:17:ASP:O	1:A:83:ARG:HD3	2.22	0.40
1:A:362:THR:OG1	1:A:363:ASN:N	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	550/563 (98%)	517 (94%)	29 (5%)	4 (1%)	18	47
2	B	392/443 (88%)	369 (94%)	21 (5%)	2 (0%)	24	55
All	All	942/1006 (94%)	886 (94%)	50 (5%)	6 (1%)	21	51

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	346	PHE
1	A	288	ALA
1	A	543	GLY
1	A	345	PRO
2	B	90	VAL
1	A	276	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	493/503 (98%)	451 (92%)	42 (8%)	10	31
2	B	364/403 (90%)	334 (92%)	30 (8%)	10	33
All	All	857/906 (95%)	785 (92%)	72 (8%)	10	32

All (72) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	ILE
1	A	7	THR
1	A	21	VAL
1	A	26	LEU
1	A	63	ILE
1	A	70	LYS
1	A	72	ARG
1	A	90	VAL
1	A	101	LYS
1	A	107	THR
1	A	118	VAL
1	A	126	LYS
1	A	138	GLU
1	A	161	GLN
1	A	173	LYS
1	A	182	GLN
1	A	202	ILE
1	A	205	LEU
1	A	210	LEU
1	A	215	THR
1	A	260	LEU
1	A	276	VAL
1	A	279	LEU
1	A	284	ARG
1	A	287	LYS
1	A	289	LEU
1	A	303	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	305	GLU
1	A	312	GLU
1	A	344	GLU
1	A	368	LEU
1	A	373	GLN
1	A	402	TRP
1	A	422	LEU
1	A	428	GLN
1	A	449	GLU
1	A	479	LEU
1	A	493	VAL
1	A	496	VAL
1	A	503	LEU
1	A	547	GLN
1	A	548	VAL
2	B	6	GLU
2	B	8	VAL
2	B	11	LYS
2	B	12	LEU
2	B	26	LEU
2	B	58	THR
2	B	72	ARG
2	B	80	LEU
2	B	91	GLN
2	B	111	VAL
2	B	120	LEU
2	B	162	SER
2	B	179	VAL
2	B	205	LEU
2	B	209	LEU
2	B	253	THR
2	B	260	LEU
2	B	270	ILE
2	B	275	LYS
2	B	279	LEU
2	B	295	LEU
2	B	301	LEU
2	B	310	LEU
2	B	336	GLN
2	B	347	LYS
2	B	349	LEU
2	B	356	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	361	HIS
2	B	377	THR
2	B	400	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (24) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	57	ASN
1	A	147	ASN
1	A	175	ASN
1	A	182	GLN
1	A	242	GLN
1	A	258	GLN
1	A	330	GLN
1	A	332	GLN
1	A	418	ASN
1	A	428	GLN
1	A	475	GLN
1	A	480	GLN
1	A	509	GLN
1	A	520	GLN
2	B	96	HIS
2	B	147	ASN
2	B	175	ASN
2	B	182	GLN
2	B	258	GLN
2	B	306	ASN
2	B	336	GLN
2	B	367	GLN
2	B	373	GLN
2	B	394	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	LP9	A	601	-	32,32,32	1.52	5 (15%)	43,45,45	1.40	6 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	LP9	A	601	-	-	8/19/19/19	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601	LP9	O11-C6	5.21	1.33	1.23
3	A	601	LP9	C4-C8	-3.32	1.40	1.45
3	A	601	LP9	O13-C14	-2.60	1.34	1.39
3	A	601	LP9	O13-C1	-2.46	1.34	1.39
3	A	601	LP9	C3-N5	-2.30	1.35	1.38

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	LP9	N-C3-N5	3.78	121.61	117.35
3	A	601	LP9	C15-C16-C17	2.71	123.73	120.30
3	A	601	LP9	C2-N-C3	2.63	123.33	115.19
3	A	601	LP9	C1-C2-N	-2.62	119.43	122.67

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	LP9	C18-C17-C16	-2.27	114.86	119.18
3	A	601	LP9	C19-C18-C17	2.22	123.11	120.30

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	601	LP9	C7-C9-O25-C27
3	A	601	LP9	C16-C17-N20-C23
3	A	601	LP9	O10-C9-O25-C27
3	A	601	LP9	C18-C17-N20-C23
3	A	601	LP9	C6-C7-C9-O10
3	A	601	LP9	C6-C7-C9-O25
3	A	601	LP9	C16-C17-N20-C21
3	A	601	LP9	C18-C17-N20-C21

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	601	LP9	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	554/563 (98%)	0.53	25 (4%) 38 30	48, 71, 90, 99	0
2	B	400/443 (90%)	0.39	17 (4%) 40 31	42, 63, 102, 106	0
All	All	954/1006 (94%)	0.47	42 (4%) 39 31	42, 69, 95, 106	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	0	SER	3.9
2	B	362	THR	3.6
1	A	7	THR	3.5
1	A	195	ILE	3.3
2	B	252	TRP	3.3
1	A	215	THR	3.2
1	A	359	GLY	3.2
2	B	293	ILE	3.1
1	A	92	LEU	2.9
1	A	460	ASN	2.8
2	B	90	VAL	2.7
2	B	346	PHE	2.7
1	A	289	LEU	2.7
2	B	301	LEU	2.6
2	B	282	LEU	2.5
1	A	286	THR	2.4
2	B	286	THR	2.4
1	A	247	PRO	2.4
2	B	24	TRP	2.4
2	B	288	ALA	2.4
1	A	24	TRP	2.3
2	B	8	VAL	2.3
1	A	450	THR	2.3
2	B	292	VAL	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	549	ASP	2.3
1	A	198	HIS	2.3
2	B	253	THR	2.3
2	B	231	GLY	2.3
1	A	209	LEU	2.2
2	B	275	LYS	2.2
1	A	250	ASP	2.2
1	A	106	VAL	2.2
1	A	301	LEU	2.2
1	A	64	LYS	2.1
1	A	207	GLN	2.1
1	A	107	THR	2.1
2	B	289	LEU	2.1
1	A	290	THR	2.1
1	A	134	SER	2.0
1	A	373	GLN	2.0
2	B	294	PRO	2.0
1	A	295	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	MN	A	602	1/1	0.78	0.15	149,149,149,149	0
3	LP9	A	601	30/30	0.82	0.15	78,80,81,82	0
4	MN	A	603	1/1	0.94	0.06	92,92,92,92	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.