



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 01:31 AM UTC

PDB ID : 5LPH / pdb_00005lph
Title : Structure of the TOG domain of human Cep104
Authors : van Breugel, M.
Deposited on : 2016-08-13
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1898 (2.26-2.26)
Clashscore	190562	2005 (2.26-2.26)
Ramachandran outliers	187476	1965 (2.26-2.26)
RSRZ outliers	180081	1898 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	288	28% 87% 10%
1	B	288	8% 88% 9%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4250 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Centrosomal protein of 104 kDa.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	260	Total	C	N	O	S	Se	0	2	0
			2061	1317	355	378	2	9			
1	B	261	Total	C	N	O	S	Se	0	3	0
			2074	1324	359	379	2	10			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	389	GLY	-	expression tag	UNP O60308
A	390	PRO	-	expression tag	UNP O60308
A	391	HIS	-	expression tag	UNP O60308
B	389	GLY	-	expression tag	UNP O60308
B	390	PRO	-	expression tag	UNP O60308
B	391	HIS	-	expression tag	UNP O60308


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	30	Total	O	0	0
			30	30		
2	B	85	Total	O	0	0
			85	85		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Centrosomal protein of 104 kDa



- Molecule 1: Centrosomal protein of 104 kDa



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	52.29Å 53.45Å 155.36Å 90.00° 91.87° 90.00°	Depositor
Resolution (Å)	49.50 – 2.25 49.50 – 2.26	Depositor EDS
% Data completeness (in resolution range)	98.6 (49.50-2.25) 98.6 (49.50-2.26)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.98 (at 2.27Å)	Xtriage
Refinement program	REFMAC 5.8.0137	Depositor
R, R_{free}	0.234 , 0.253 (Not available) , 0.234	Depositor DCC
R_{free} test set	2031 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	35.4	Xtriage
Anisotropy	0.371	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 28.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.015 for -k,-h,-l 0.017 for k,h,-l 0.039 for h,-k,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4250	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.63	1/2091 (0.0%)	0.92	2/2809 (0.1%)
1	B	0.62	0/2107	0.92	1/2829 (0.0%)
All	All	0.63	1/4198 (0.0%)	0.92	3/5638 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	530	ILE	CA-CB	5.83	1.57	1.53

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	467	VAL	N-CA-C	6.63	118.18	110.62
1	B	530	ILE	N-CA-CB	6.29	114.46	110.50
1	A	530	ILE	N-CA-CB	5.46	113.94	110.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2061	0	2167	6	0
1	B	2074	0	2183	6	0
2	A	30	0	0	0	0
2	B	85	0	0	0	0
All	All	4250	0	4350	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:585[A]:MSE:HE1	1:B:624:GLU:HB2	1.66	0.76
1:B:588:MSE:HE2	1:B:628:THR:HB	1.85	0.59
1:A:530:ILE:CD1	1:A:552:ILE:HG12	2.40	0.51
1:A:577:ALA:O	1:A:621:ARG:NH1	2.44	0.50
1:A:515:HIS:HB2	1:A:517:LEU:HD13	1.96	0.47
1:B:588:MSE:HE1	1:B:632:ILE:HD12	1.96	0.47
1:A:532:VAL:O	1:A:535:THR:OG1	2.32	0.46
1:A:577:ALA:HB1	1:B:657:ARG:HD3	1.99	0.44
1:A:530:ILE:CG2	1:A:531:PRO:HD3	2.49	0.43
1:B:654:ASN:OD1	1:B:655:THR:N	2.52	0.43
1:B:585[A]:MSE:CE	1:B:624:GLU:HB2	2.44	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	260/288 (90%)	257 (99%)	3 (1%)	0	100	100
1	B	262/288 (91%)	259 (99%)	3 (1%)	0	100	100
All	All	522/576 (91%)	516 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	252/288 (87%)	1.56	81 (32%) 1 0	27, 56, 99, 113	1 (0%)
1	B	253/288 (87%)	0.53	22 (8%) 16 15	16, 38, 76, 117	1 (0%)
All	All	505/576 (87%)	1.04	103 (20%) 3 2	16, 46, 95, 117	2 (0%)

All (103) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	414	LEU	7.3
1	A	669	ALA	5.4
1	A	665	PHE	5.4
1	A	673	GLY	5.1
1	A	650	PRO	4.9
1	B	653	SER	4.5
1	A	659	ILE	4.4
1	A	645	LEU	4.4
1	B	675	ALA	4.2
1	B	654	ASN	4.1
1	A	642	ALA	4.1
1	B	650	PRO	4.0
1	A	467	VAL	3.9
1	A	671	ILE	3.8
1	A	601	GLY	3.8
1	A	618	LEU	3.8
1	A	619	GLU	3.8
1	A	660	LEU	3.8
1	B	651	ASP	3.7
1	A	644	ILE	3.7
1	B	655	THR	3.7
1	A	668	PHE	3.7
1	A	658	ASN	3.6
1	B	656	ARG	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	661	TYR	3.5
1	B	657	ARG	3.5
1	B	658	ASN	3.4
1	A	558	PHE	3.4
1	A	655	THR	3.4
1	B	415	GLY	3.3
1	A	625	VAL	3.3
1	A	648	LEU	3.2
1	A	615	VAL	3.2
1	A	468	GLY	3.2
1	A	600	THR	3.2
1	A	613	PHE	3.2
1	A	471	LYS	3.2
1	B	660	LEU	3.1
1	A	623	TYR	3.0
1	B	577	ALA	3.0
1	B	673	GLY	3.0
1	A	634	LEU	3.0
1	A	622	VAL	2.9
1	A	559	LYS	2.9
1	A	670	LYS	2.9
1	A	638	ARG	2.9
1	A	640	HIS	2.9
1	A	603	SER	2.9
1	A	635	ASP	2.9
1	A	643	SER	2.9
1	A	657	ARG	2.9
1	A	631	ARG	2.8
1	A	656	ARG	2.8
1	A	562	LYS	2.8
1	A	629	ALA	2.8
1	B	644	ILE	2.7
1	A	479[A]	ARG	2.7
1	A	581	VAL	2.7
1	A	666	GLU	2.7
1	B	659	ILE	2.7
1	A	557	LEU	2.7
1	B	652	ASP	2.6
1	A	664	ILE	2.6
1	A	639	GLN	2.6
1	A	415	GLY	2.6
1	A	530	ILE	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	627	GLU	2.6
1	A	632	ILE	2.5
1	A	637	TYR	2.5
1	A	564	LEU	2.5
1	A	466	PRO	2.5
1	A	649	PRO	2.5
1	A	624	GLU	2.5
1	A	609	ASN	2.5
1	A	654	ASN	2.5
1	A	573	GLN	2.5
1	A	599	GLY	2.5
1	A	630	VAL	2.4
1	B	641	GLN	2.4
1	A	560	GLU	2.4
1	A	617	ALA	2.3
1	B	639	GLN	2.3
1	A	651	ASP	2.3
1	A	516	LYS	2.3
1	A	596	LYS	2.2
1	A	602	SER	2.2
1	A	456	LEU	2.2
1	A	616	SER	2.2
1	A	597	ASP	2.2
1	A	502	LYS	2.2
1	B	464	GLU	2.2
1	A	612	LYS	2.1
1	A	641	GLN	2.1
1	A	627	GLU	2.1
1	A	470	PRO	2.1
1	A	459	SER	2.1
1	A	450	TYR	2.1
1	A	565	GLN	2.1
1	A	667	GLY	2.1
1	A	607	ILE	2.0
1	B	649	PRO	2.0
1	A	646	GLU	2.0
1	B	575	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.