



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 06:52 AM UTC

PDB ID : 3LQH / pdb_00003lqh
Title : Crystal structure of MLL1 PHD3-Bromo in the free form
Authors : Wang, Z.; Patel, D.J.
Deposited on : 2010-02-09
Resolution : 1.72 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1039 (1.72-1.72)
Clashscore	190562	1049 (1.72-1.72)
Ramachandran outliers	187476	1041 (1.72-1.72)
Sidechain outliers	187428	1041 (1.72-1.72)
RSRZ outliers	180081	1039 (1.72-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	183	5% 67% 25% • 5%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1571 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone-lysine N-methyltransferase MLL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	174	Total	C	N	O	S	0	0	0
			1420	893	244	269	14			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1565	SER	-	expression tag	UNP Q03164

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	149	Total	O	0	0
			149	149		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Histone-lysine N-methyltransferase MLL



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	63.22Å 71.88Å 93.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.72 50.00 – 1.72	Depositor EDS
% Data completeness (in resolution range)	93.6 (50.00-1.72) 93.6 (50.00-1.72)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.61 (at 1.60Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.206 , 0.234 0.202 , 0.223	Depositor DCC
R_{free} test set	2409 reflections (9.29%)	wwPDB-VP
Wilson B-factor (Å ²)	11.7	Xtriage
Anisotropy	0.627	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 45.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	1571	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	2.07	58/1448 (4.0%)	1.15	10/1948 (0.5%)

All (58) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1655	ARG	C-O	-15.87	1.05	1.24
1	A	1733	ILE	C-O	-15.11	1.06	1.24
1	A	1607	TYR	C-O	-13.78	1.08	1.24
1	A	1605	GLU	C-O	-12.99	1.09	1.24
1	A	1625	THR	C-O	-12.22	1.08	1.24
1	A	1745	PRO	C-O	-11.73	1.08	1.24
1	A	1596	HIS	C-O	-11.73	1.09	1.23
1	A	1707	ASP	C-O	-11.28	1.09	1.23
1	A	1600	GLU	C-O	-11.15	1.09	1.23
1	A	1705	PRO	C-O	-10.66	1.12	1.23
1	A	1627	ARG	C-O	-10.43	1.10	1.23
1	A	1630	ALA	CA-CB	-10.27	1.36	1.53
1	A	1645	LYS	C-O	-10.27	1.12	1.24
1	A	1744	GLN	C-O	-10.18	1.11	1.24
1	A	1626	GLU	C-O	-10.09	1.12	1.24
1	A	1606	MET	C-O	-9.59	1.12	1.24
1	A	1602	LEU	C-O	-9.35	1.13	1.23
1	A	1767	PRO	C-O	-9.32	1.12	1.24
1	A	1659	HIS	C-O	-9.32	1.13	1.24
1	A	1707	ASP	N-CA	-8.95	1.34	1.45
1	A	1706	LEU	C-O	-8.86	1.12	1.24
1	A	1633	ARG	C-O	-8.63	1.13	1.24
1	A	1775	ARG	CD-NE	-8.58	1.34	1.46
1	A	1573	ASP	CG-OD1	-7.97	1.10	1.25
1	A	1653	ASN	C-O	-7.85	1.13	1.24
1	A	1639	GLU	C-O	-7.83	1.15	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1708	LEU	C-O	-7.55	1.14	1.24
1	A	1708	LEU	N-CA	-7.21	1.37	1.46
1	A	1615	GLU	C-O	-7.07	1.14	1.24
1	A	1602	LEU	CG-CD2	-7.07	1.29	1.52
1	A	1601	ASN	C-O	-7.05	1.15	1.23
1	A	1626	GLU	CA-C	7.04	1.61	1.52
1	A	1709	GLU	C-O	-6.96	1.16	1.24
1	A	1600	GLU	CA-C	6.85	1.60	1.52
1	A	1775	ARG	C-O	-6.76	1.15	1.24
1	A	1571	LEU	CG-CD2	-6.51	1.31	1.52
1	A	1705	PRO	CG-CD	-6.50	1.28	1.50
1	A	1571	LEU	C-O	-6.47	1.15	1.24
1	A	1706	LEU	CG-CD1	-6.27	1.31	1.52
1	A	1615	GLU	CA-C	-6.26	1.44	1.52
1	A	1571	LEU	N-CA	-6.23	1.37	1.46
1	A	1653	ASN	CG-OD1	-6.18	1.11	1.23
1	A	1630	ALA	C-O	-6.11	1.16	1.23
1	A	1607	TYR	CA-C	6.10	1.60	1.52
1	A	1625	THR	C-N	-5.93	1.26	1.33
1	A	1709	GLU	CD-OE1	-5.85	1.14	1.25
1	A	1745	PRO	CB-CG	-5.81	1.20	1.49
1	A	1744	GLN	CG-CD	-5.69	1.37	1.52
1	A	1704	GLN	CG-CD	-5.53	1.38	1.52
1	A	1627	ARG	CZ-NH1	-5.35	1.25	1.32
1	A	1705	PRO	CA-CB	-5.34	1.46	1.53
1	A	1639	GLU	CD-OE2	-5.31	1.15	1.25
1	A	1573	ASP	C-O	-5.30	1.15	1.24
1	A	1626	GLU	C-N	-5.27	1.26	1.33
1	A	1605	GLU	CB-CG	-5.26	1.36	1.52
1	A	1704	GLN	C-O	-5.21	1.13	1.23
1	A	1606	MET	CB-CG	-5.08	1.37	1.52
1	A	1601	ASN	CG-OD1	-5.06	1.14	1.23

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1625	THR	CA-C-N	6.54	131.47	120.88
1	A	1625	THR	C-N-CA	6.54	131.47	120.88
1	A	1626	GLU	N-CA-CB	-6.40	100.67	110.07
1	A	1626	GLU	CG-CD-OE2	-6.00	104.61	118.40
1	A	1627	ARG	NE-CZ-NH2	5.92	124.52	119.20
1	A	1630	ALA	N-CA-C	5.91	118.80	110.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1626	GLU	CA-CB-CG	-5.73	102.64	114.10
1	A	1661	LEU	N-CA-C	5.70	121.66	114.31
1	A	1655	ARG	CG-CD-NE	5.24	123.54	112.00
1	A	1596	HIS	N-CA-CB	5.03	117.42	109.83

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1420	0	1385	12	0
2	A	2	0	0	0	0
3	A	149	0	0	2	0
All	All	1571	0	1385	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1584:LYS:HG2	3:A:80:HOH:O	1.93	0.66
1:A:1743:GLY:HA2	1:A:1748:LYS:HE2	1.81	0.62
1:A:1628:HIS:HA	1:A:1629:PRO:C	2.29	0.58
1:A:1709:GLU:HG3	1:A:1713:ARG:NH1	2.18	0.57
1:A:1577:ASP:HA	3:A:91:HOH:O	2.08	0.52
1:A:1610:LEU:HD23	1:A:1610:LEU:C	2.35	0.51
1:A:1617:VAL:HG21	1:A:1636:LEU:HD13	1.92	0.51
1:A:1662:ARG:HA	1:A:1708:LEU:HB2	1.95	0.48
1:A:1568:PHE:CE1	1:A:1575:CYS:HB2	2.50	0.46
1:A:1591:CYS:SG	1:A:1593:ARG:HG2	2.56	0.46
1:A:1768:TRP:CD1	1:A:1768:TRP:H	2.39	0.40
1:A:1757:PHE:O	1:A:1761:GLN:HG2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	170/183 (93%)	169 (99%)	1 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	161/170 (95%)	157 (98%)	4 (2%)	42	18

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1586	MET
1	A	1626	GLU
1	A	1706	LEU
1	A	1744	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1612	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1719	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	174/183 (95%)	0.36	10 (5%)	29 35	7, 18, 32, 37	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1626	GLU	3.5
1	A	1631	GLU	3.4
1	A	1584	LYS	3.3
1	A	1628	HIS	2.8
1	A	1706	LEU	2.7
1	A	1598	LYS	2.7
1	A	1655	ARG	2.4
1	A	1568	PHE	2.4
1	A	1634	LEU	2.3
1	A	1566	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ZN	A	1001	1/1	0.98	0.03	18,18,18,18	0
2	ZN	A	1002	1/1	0.98	0.02	13,13,13,13	0

6.5 Other polymers [i](#)

There are no such residues in this entry.