



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 11:36 PM UTC

PDB ID : 1LRQ / pdb_00001lrq
Title : Aquifex aeolicus KDO8P synthase H185G mutant in complex with PEP, A5P and Cadmium
Authors : Wang, J.; Duewel, H.S.; Stuckey, J.A.; Woodard, R.W.; Gatti, D.L.
Deposited on : 2002-05-15
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	267	4% 81% 17% .
1	B	267	9% 74% 20% ..

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4352 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called KDO-8-phosphate synthetase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	263	Total	C	N	O	S	0	0	0
			2059	1327	343	383	6			
1	B	255	Total	C	N	O	S	0	0	0
			2007	1297	334	370	6			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1185	GLY	HIS	engineered mutation	UNP O66496
B	2185	GLY	HIS	engineered mutation	UNP O66496

- Molecule 2 is CADMIUM ION (CCD ID: CD) (formula: Cd).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cd	0	0
			1	1		
2	B	1	Total	Cd	0	0
			1	1		

- Molecule 3 is PHOSPHOENOLPYRUVATE (CCD ID: PEP) (formula: C₃H₅O₆P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	P	0	0
			10	3	6	1		

- Molecule 4 is ARABINOSE-5-PHOSPHATE (CCD ID: A5P) (formula: $C_5H_{13}O_8P$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	P	0	0
			14	5	8	1		

- Molecule 5 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	O	P	0	0
			5	4	1		
5	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	142	Total	O	0	0
			142	142		
6	B	108	Total	O	0	0
			108	108		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: KDO-8-phosphate synthetase



• Molecule 1: KDO-8-phosphate synthetase



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	84.47Å 84.47Å 160.48Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.29 – 1.80 20.29 – 1.80	Depositor EDS
% Data completeness (in resolution range)	95.0 (20.29-1.80) 95.0 (20.29-1.80)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.42 (at 1.74Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.207 , 0.228 0.207 , 0.229	Depositor DCC
R_{free} test set	6013 reflections (9.35%)	wwPDB-VP
Wilson B-factor (Å ²)	29.8	Xtriage
Anisotropy	0.046	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 34.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.024 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4352	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEP, A5P, PO4, CD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/2099	0.85	2/2830 (0.1%)
1	B	0.35	0/2046	0.85	2/2759 (0.1%)
All	All	0.36	0/4145	0.85	4/5589 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	2085	SER	N-CA-C	5.88	118.44	111.33
1	A	1085	SER	N-CA-C	5.57	118.07	111.33
1	A	1179	VAL	N-CA-C	5.04	115.17	108.11
1	B	2130	ALA	N-CA-C	-5.03	103.35	110.29

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2059	0	2096	34	0
1	B	2007	0	2047	48	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	10	0	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	14	0	10	0	0
5	B	10	0	0	0	0
6	A	142	0	0	4	0
6	B	108	0	0	0	0
All	All	4352	0	4155	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1161:ASN:HD21	1:B:2132:TRP:HE1	1.16	0.94
1:A:1174:LYS:HG3	1:A:1264:ILE:HD13	1.51	0.90
1:A:1132:TRP:HE1	1:B:2161:ASN:HD21	1.22	0.86
1:A:1189:LEU:HD21	1:A:1200:MET:HE2	1.57	0.85
1:B:2185:GLY:O	1:B:2188:GLN:HG2	1.81	0.80
1:B:2208:ILE:HD11	1:B:2219:VAL:HG11	1.62	0.79
1:A:1185:GLY:O	1:A:1188:GLN:HG2	1.88	0.73
1:B:2189:LEU:HG	1:B:2199:GLY:HA2	1.69	0.73
1:B:2011:CYS:H	1:B:2237:GLN:NE2	1.89	0.71
1:B:2208:ILE:CD1	1:B:2219:VAL:HG11	2.24	0.68
1:B:2025:GLU:HG2	1:B:2240:LEU:HD22	1.75	0.67
1:B:2025:GLU:O	1:B:2029:LEU:HD22	1.96	0.66
1:B:2048:ASN:H	1:B:2048:ASN:HD22	1.44	0.65
1:B:2263:THR:HG22	1:B:2264:ILE:HG22	1.79	0.64
1:B:2201:ARG:HG3	1:B:2204:ILE:HD12	1.81	0.63
1:A:1048:ASN:H	1:A:1048:ASN:HD22	1.47	0.63
1:A:1014:GLU:HG2	1:A:1231:LEU:HD12	1.83	0.60
1:B:2011:CYS:H	1:B:2237:GLN:HE22	1.49	0.60
6:A:24:HOH:O	1:B:2083:HIS:HD2	1.87	0.57
1:B:2016:GLU:O	1:B:2020:LEU:HG	2.04	0.56
1:A:1201:ARG:HG3	1:A:1204:ILE:HD12	1.88	0.55
1:B:2055:PHE:CZ	1:B:2057:GLY:HA2	2.42	0.55
1:A:1228:GLU:HG3	1:A:1229:LYS:HG3	1.89	0.55
1:A:1041:LYS:HD3	1:A:1041:LYS:C	2.32	0.54
1:B:2019:LEU:HD12	1:B:2066:ALA:HB1	1.89	0.54
1:B:2129:LEU:HG	1:B:2133:ASP:HB2	1.91	0.53
1:B:2041:LYS:HD3	1:B:2041:LYS:C	2.35	0.52
1:B:2162:ASN:HD22	1:B:2163:LEU:H	1.56	0.52
1:A:1060:LEU:C	1:A:1060:LEU:HD13	2.34	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2205:PHE:O	1:B:2208:ILE:HG22	2.11	0.51
1:A:1048:ASN:HD22	1:A:1048:ASN:N	2.07	0.50
1:A:1099:GLN:HG2	6:A:233:HOH:O	2.11	0.50
1:A:1041:LYS:HD3	1:A:1042:SER:N	2.27	0.49
1:A:1026:ILE:HD12	1:A:1040:PHE:CD1	2.47	0.49
1:B:2048:ASN:HD22	1:B:2048:ASN:N	2.06	0.49
1:A:1212:VAL:HG11	1:A:1250:ILE:HB	1.95	0.49
1:A:1161:ASN:ND2	1:B:2132:TRP:HE1	1.98	0.48
1:B:2068:ARG:O	1:B:2072:GLU:HG3	2.14	0.48
1:B:2044:PHE:CZ	1:B:2060:LEU:HD13	2.48	0.48
1:A:1132:TRP:HE1	1:B:2161:ASN:ND2	2.02	0.47
1:B:2238:LEU:HD21	1:B:2246:ILE:HD12	1.96	0.47
1:B:2041:LYS:HD3	1:B:2042:SER:N	2.29	0.47
1:B:2205:PHE:HB3	1:B:2206:PRO:HD3	1.96	0.47
1:A:1132:TRP:NE1	1:B:2161:ASN:HD21	2.00	0.46
1:A:1182:ASP:HA	1:A:1220:PHE:HB3	1.97	0.46
1:A:1014:GLU:O	1:A:1058:HIS:HE1	1.99	0.45
1:B:2040:PHE:HB3	1:B:2078:ILE:HD13	1.98	0.45
1:A:1205:PHE:O	1:A:1208:ILE:HG22	2.16	0.45
1:A:1050:SER:O	1:B:2136:ASN:HB3	2.17	0.45
1:A:1176:TRP:HE3	6:A:249:HOH:O	1.99	0.45
1:A:1193:LEU:HD23	1:A:1198:GLY:CA	2.47	0.45
1:A:1233:ASP:O	1:A:1237:GLN:HG3	2.16	0.45
1:B:2200:MET:HB2	1:B:2203:PHE:CD2	2.53	0.44
1:A:1003:LYS:HE2	1:A:1036:VAL:C	2.42	0.44
1:A:1026:ILE:HD12	1:A:1040:PHE:HD1	1.83	0.44
1:B:2162:ASN:HD22	1:B:2163:LEU:N	2.15	0.44
6:A:95:HOH:O	1:B:2053:HIS:HE1	2.01	0.43
1:A:1161:ASN:HD21	1:B:2132:TRP:NE1	1.98	0.43
1:B:2200:MET:HB2	1:B:2203:PHE:HD2	1.83	0.43
1:A:1060:LEU:O	1:A:1064:VAL:HG23	2.19	0.42
1:B:2200:MET:HB3	1:B:2202:GLU:HG2	1.99	0.42
1:B:2263:THR:O	1:B:2264:ILE:C	2.62	0.42
1:B:2013:ILE:HG23	1:B:2019:LEU:HD11	2.01	0.42
1:B:2022:VAL:O	1:B:2026:ILE:HG12	2.19	0.42
1:B:2005:LEU:HD12	1:B:2037:GLU:O	2.20	0.42
1:B:2219:VAL:HG12	1:B:2220:PHE:N	2.35	0.42
1:B:2010:PRO:HD3	1:B:2022:VAL:HG11	2.01	0.41
1:A:1103:PHE:HE2	1:B:2106:ARG:NH1	2.18	0.41
1:A:1162:ASN:HD22	1:A:1163:LEU:H	1.69	0.41
1:A:1005:LEU:HD12	1:A:1037:GLU:HB3	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2068:ARG:NH1	1:B:2094:VAL:HG22	2.35	0.41
1:B:2068:ARG:HH11	1:B:2094:VAL:HG22	1.86	0.40
1:A:1140:LYS:HG2	1:B:2052:ILE:HB	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	261/267 (98%)	257 (98%)	4 (2%)	0	100	100
1	B	251/267 (94%)	246 (98%)	5 (2%)	0	100	100
All	All	512/534 (96%)	503 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	219/223 (98%)	216 (99%)	3 (1%)	59	52
1	B	214/223 (96%)	210 (98%)	4 (2%)	50	41
All	All	433/446 (97%)	426 (98%)	7 (2%)	55	47

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1048	ASN
1	A	1106	ARG
1	A	1112	LEU
1	B	2029	LEU
1	B	2048	ASN
1	B	2106	ARG
1	B	2240	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1048	ASN
1	A	1058	HIS
1	A	1099	GLN
1	A	1161	ASN
1	A	1162	ASN
1	A	1175	GLN
1	B	2048	ASN
1	B	2058	HIS
1	B	2083	HIS
1	B	2122	ASN
1	B	2161	ASN
1	B	2162	ASN
1	B	2188	GLN
1	B	2237	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	PO4	B	2269	-	4,4,4	1.72	0	6,6,6	0.46	0
5	PO4	B	2268	-	4,4,4	1.67	0	6,6,6	0.45	0
3	PEP	A	1268	2	9,9,9	2.68	3 (33%)	11,13,13	2.39	2 (18%)
4	A5P	A	1269	-	13,13,13	1.51	1 (7%)	17,18,18	4.85	9 (52%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEP	A	1268	2	-	0/9/9/9	-
4	A5P	A	1269	-	-	7/16/16/16	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1268	PEP	C2-C1	6.07	1.55	1.49
4	A	1269	A5P	O1-C1	-4.86	1.22	1.42
3	A	1268	PEP	O2'-C1	-4.01	1.19	1.30
3	A	1268	PEP	O1-C1	2.09	1.27	1.22

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1269	A5P	O2-C2-C3	9.03	130.37	109.25
4	A	1269	A5P	O4-C4-C3	8.62	129.42	109.25
4	A	1269	A5P	C1-C2-C3	-7.35	97.19	112.17
4	A	1269	A5P	O4-C4-C5	-7.15	94.22	109.99
4	A	1269	A5P	O3-C3-C2	6.29	123.22	108.93

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1268	PEP	O2'-C1-C2	5.89	123.96	113.91
4	A	1269	A5P	C5-C4-C3	-5.26	102.29	112.22
4	A	1269	A5P	O1-C1-C2	5.23	122.14	111.16
4	A	1269	A5P	C4-C3-C2	-4.48	106.11	113.57
3	A	1268	PEP	O1-C1-C2	-4.06	115.66	121.79
4	A	1269	A5P	O2-C2-C1	3.95	118.00	109.03

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1269	A5P	O1-C1-C2-O2
4	A	1269	A5P	O3-C3-C4-O4
4	A	1269	A5P	C2-C3-C4-O4
4	A	1269	A5P	O3-C3-C4-C5
4	A	1269	A5P	C2-C3-C4-C5
4	A	1269	A5P	C1-C2-C3-O3
4	A	1269	A5P	C1-C2-C3-C4

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	263/267 (98%)	0.19	11 (4%) 40 40	20, 28, 45, 58	0
1	B	255/267 (95%)	0.57	24 (9%) 14 12	21, 32, 50, 62	0
All	All	518/534 (97%)	0.38	35 (6%) 23 22	20, 30, 48, 62	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2264	ILE	6.2
1	B	2199	GLY	5.2
1	A	1264	ILE	4.9
1	B	2094	VAL	3.9
1	B	2200	MET	3.9
1	B	2035	GLU	3.7
1	A	1228	GLU	3.3
1	A	1195	ASP	3.3
1	A	1197	SER	3.3
1	A	1198	GLY	3.2
1	B	2068	ARG	3.2
1	B	2060	LEU	3.2
1	A	1193	LEU	3.2
1	B	2230	ALA	3.0
1	B	2061	GLU	2.6
1	B	2147	LYS	2.6
1	A	1002	GLU	2.6
1	A	1263	THR	2.4
1	B	2234	ALA	2.4
1	B	2263	THR	2.4
1	A	1194	GLY	2.4
1	B	2235	SER	2.4
1	B	2093	GLU	2.4
1	B	2017	GLU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	2188	GLN	2.3
1	A	1201	ARG	2.2
1	B	2034	LYS	2.2
1	B	2036	VAL	2.2
1	B	2070	VAL	2.2
1	B	2191	GLY	2.1
1	B	2092	ALA	2.1
1	B	2016	GLU	2.1
1	B	2090	PRO	2.1
1	A	1200	MET	2.0
1	B	2072	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	A5P	A	1269	14/14	0.90	0.15	44,60,64,64	0
3	PEP	A	1268	10/10	0.93	0.13	30,36,45,47	0
5	PO4	B	2269	5/5	0.96	0.07	47,48,49,49	0
5	PO4	B	2268	5/5	0.98	0.10	31,32,33,35	0
2	CD	B	2270	1/1	0.98	0.03	30,30,30,30	1
2	CD	A	1270	1/1	0.99	0.02	26,26,26,26	1

6.5 Other polymers [i](#)

There are no such residues in this entry.