



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 03:03 PM UTC

PDB ID : 5LRF / pdb_00005lrf
Title : Crystal structure of Glycogen Phosphorylase b in complex with KS389
Authors : Kantsadi, A.L.; Leonidas, D.D.
Deposited on : 2016-08-18
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	842	5% 89% 7% .

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 6990 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycogen phosphorylase, muscle form.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	810	Total	C	N	O	S	0	1	0
			6596	4206	1162	1199	29			

- Molecule 2 is (2 {R},3 {S},4 {R},5 {R},6 {S})-2-(hydroxymethyl)-6-(3-naphthalen-2-yl-1 {H})-1,2,4-triazol-5-yl)oxane-3,4,5-triol (CCD ID: KS3) (formula: C₁₈H₁₉N₃O₅).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			26	18	3	5		
2	A	1	Total	C	N	O	0	0
			26	18	3	5		

- Molecule 3 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	S	0	0
			4	2	1	1		
3	A	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 4 is PYRIDOXAL-5'-PHOSPHATE (CCD ID: PLP) (formula: $C_8H_{10}NO_6P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	319	Total 319	O 319	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glycogen phosphorylase, muscle form



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	128.56Å 128.56Å 116.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.14 – 1.75 37.14 – 1.75	Depositor EDS
% Data completeness (in resolution range)	11.4 (37.14-1.75) 92.2 (37.14-1.75)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.41 (at 1.75Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.165 , 0.185 (Not available) , 0.196	Depositor DCC
R_{free} test set	4438 reflections (4.52%)	wwPDB-VP
Wilson B-factor (Å ²)	26.7	Xtriage
Anisotropy	0.011	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 40.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6990	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DMS, PLP, KS3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	0/6747	0.85	4/9131 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	398	ARG	NE-CZ-NH2	6.29	124.86	119.20
1	A	751	SER	CA-C-N	5.28	124.73	119.24
1	A	751	SER	C-N-CA	5.28	124.73	119.24
1	A	507	ILE	N-CA-C	5.22	117.23	112.43

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6596	0	6536	24	0
2	A	52	0	0	5	2
3	A	8	0	12	2	0
4	A	15	0	6	0	0
5	A	319	0	0	4	1
All	All	6990	0	6554	27	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:192:ALA:H	2:A:902:KS3:C6'	1.68	1.06
1:A:549:LEU:HB3	1:A:555:VAL:HG23	1.52	0.91
1:A:65:GLY:O	1:A:69:ARG:HG2	1.76	0.85
1:A:759:LYS:HE3	1:A:763:ASN:OD1	1.92	0.70
1:A:184:ARG:NH1	5:A:1002:HOH:O	2.28	0.67
1:A:192:ALA:N	2:A:902:KS3:C6'	2.51	0.66
1:A:355:ASP:OD2	1:A:398:ARG:HD3	1.98	0.64
1:A:549:LEU:HB3	1:A:555:VAL:CG2	2.26	0.63
1:A:739:ARG:O	1:A:743:GLU:HG3	2.03	0.59
1:A:235:ASN:HA	1:A:833:ARG:HG3	1.86	0.57
1:A:329:PHE:HB3	1:A:330:PRO:HD3	1.89	0.55
1:A:325:ASN:ND2	1:A:327:ASP:OD1	2.36	0.53
1:A:358:ARG:NH1	5:A:1006:HOH:O	2.42	0.52
3:A:904:DMS:H21	5:A:1313:HOH:O	2.10	0.51
1:A:60:ARG:HD3	2:A:902:KS3:C10	2.42	0.49
3:A:903:DMS:C1	3:A:904:DMS:H22	2.44	0.48
1:A:60:ARG:CG	2:A:902:KS3:C15	2.92	0.47
1:A:550:GLU:OE2	1:A:556:HIS:ND1	2.49	0.45
2:A:902:KS3:O4'	5:A:1001:HOH:O	2.21	0.43
1:A:144:LEU:HD23	1:A:147:MET:CE	2.49	0.42
1:A:426:ARG:HG3	1:A:427:ARG:N	2.34	0.42
1:A:433:GLU:OE1	1:A:433:GLU:HA	2.18	0.42
1:A:708:PHE:HB3	1:A:710:ILE:CD1	2.50	0.41
1:A:730:GLU:O	1:A:734:ARG:CG	2.68	0.41
1:A:730:GLU:O	1:A:734:ARG:HG3	2.20	0.41
1:A:553:TYR:O	1:A:555:VAL:HG13	2.21	0.41
1:A:562:LEU:C	1:A:562:LEU:HD23	2.46	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:902:KS3:O2'	2:A:902:KS3:O2'[7_556]	1.53	0.67
2:A:902:KS3:O3'	2:A:902:KS3:O3'[7_556]	1.70	0.50
5:A:1054:HOH:O	5:A:1221:HOH:O[7_556]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	805/842 (96%)	782 (97%)	23 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	702/731 (96%)	678 (97%)	24 (3%)	32	13

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	SER
1	A	22	GLU
1	A	43	ARG
1	A	76	GLU
1	A	102	LEU
1	A	115	LEU
1	A	210	SER
1	A	211	GLN
1	A	234	ARG
1	A	253	ASN
1	A	271	LEU
1	A	339	ASP
1	A	398	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	552	GLU
1	A	554	LYS
1	A	565	VAL
1	A	568	LYS
1	A	586	LEU
1	A	596	LYS
1	A	706	GLU
1	A	721	LEU
1	A	734	ARG
1	A	754	GLN
1	A	765	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	96	GLN
1	A	211	GLN
1	A	284	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	DMS	A	904	-	3,3,3	0.40	0	3,3,3	0.98	0
3	DMS	A	903	-	3,3,3	0.46	0	3,3,3	0.55	0
4	PLP	A	905	1	15,15,16	2.45	3 (20%)	21,22,23	1.50	5 (23%)
2	KS3	A	902	-	29,29,29	1.81	8 (27%)	36,42,42	2.91	16 (44%)
2	KS3	A	901	-	29,29,29	1.05	2 (6%)	36,42,42	1.53	7 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	KS3	A	902	-	-	2/6/30/30	0/4/4/4
2	KS3	A	901	-	-	0/6/30/30	0/4/4/4
4	PLP	A	905	1	-	2/6/6/8	0/1/1/1

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	905	PLP	C5-C4	6.03	1.47	1.40
4	A	905	PLP	C3-C2	5.79	1.47	1.41
2	A	902	KS3	C1'-C1	-4.27	1.44	1.50
4	A	905	PLP	C3-C4	3.35	1.46	1.40
2	A	902	KS3	O5'-C1'	-3.17	1.37	1.43
2	A	902	KS3	C9-C8	2.90	1.48	1.42
2	A	902	KS3	O5'-C5'	-2.76	1.37	1.44
2	A	902	KS3	C4-N3	2.67	1.36	1.32
2	A	902	KS3	C2'-C1'	-2.57	1.49	1.53
2	A	902	KS3	C7-C8	-2.42	1.36	1.42
2	A	902	KS3	C6'-C5'	-2.22	1.44	1.51
2	A	901	KS3	C9-C8	2.19	1.47	1.42
2	A	901	KS3	C1-N5	2.00	1.35	1.32

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	902	KS3	C1'-C1-N2	7.62	135.80	122.28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	902	KS3	C5'-O5'-C1'	-6.06	101.33	112.86
2	A	902	KS3	O2'-C2'-C1'	-5.62	96.92	109.76
2	A	902	KS3	C3'-C4'-C5'	-5.22	100.76	110.23
2	A	901	KS3	C3'-C4'-C5'	4.52	118.43	110.23
2	A	902	KS3	C3'-C2'-C1'	-4.43	101.69	109.30
2	A	902	KS3	O4'-C4'-C3'	4.32	120.55	110.38
2	A	902	KS3	O5'-C1'-C2'	-4.17	101.25	109.48
2	A	902	KS3	O5'-C5'-C4'	-3.82	102.81	109.70
2	A	902	KS3	C6-C4-N3	3.58	128.97	122.57
2	A	902	KS3	O2'-C2'-C3'	3.28	118.11	110.38
2	A	901	KS3	O4'-C4'-C5'	2.97	116.65	109.32
2	A	902	KS3	C11-C6-C7	2.89	122.71	119.21
2	A	902	KS3	N3-C4-N5	-2.86	110.11	114.17
2	A	902	KS3	O4'-C4'-C5'	-2.66	102.77	109.32
4	A	905	PLP	O4P-P-O1P	-2.57	99.49	106.44
2	A	902	KS3	O5'-C5'-C6'	-2.56	100.11	106.44
2	A	901	KS3	N3-C4-N5	-2.54	110.55	114.17
4	A	905	PLP	C4A-C4-C5	-2.47	118.39	120.94
2	A	902	KS3	O6'-C6'-C5'	-2.41	103.13	111.33
2	A	901	KS3	C11-C6-C7	2.36	122.06	119.21
4	A	905	PLP	C4A-C4-C3	2.29	124.34	120.52
2	A	901	KS3	C6-C7-C8	-2.16	117.87	121.19
4	A	905	PLP	O3P-P-O2P	2.15	115.87	107.80
2	A	901	KS3	C6-C4-N5	2.12	126.63	122.92
2	A	901	KS3	O4'-C4'-C3'	2.07	115.26	110.38
4	A	905	PLP	O3-C3-C2	2.00	121.73	117.58
2	A	902	KS3	O3'-C3'-C4'	-2.00	105.66	110.38

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	902	KS3	O5'-C5'-C6'-O6'
2	A	902	KS3	C4'-C5'-C6'-O6'
4	A	905	PLP	C6-C5-C5A-O4P
4	A	905	PLP	C4-C5-C5A-O4P


There are no ring outliers.

3 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	904	DMS	2	0
3	A	903	DMS	1	0
2	A	902	KS3	5	2

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	810/842 (96%)	0.27	40 (4%) 35 40	14, 28, 52, 84	1 (0%)

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	324	THR	4.5
1	A	252	PHE	4.5
1	A	13	ILE	4.4
1	A	555	VAL	4.2
1	A	553	TYR	3.9
1	A	556	HIS	3.9
1	A	836	ALA	3.7
1	A	14	SER	3.2
1	A	435	ALA	3.0
1	A	209	THR	2.7
1	A	21	VAL	2.7
1	A	72	GLN	2.7
1	A	19	ALA	2.7
1	A	551	ARG	2.6
1	A	76	GLU	2.6
1	A	253	ASN	2.5
1	A	325	ASN	2.5
1	A	548	TYR	2.5
1	A	720	ARG	2.5
1	A	113	TYR	2.4
1	A	271	LEU	2.3
1	A	532	ARG	2.3
1	A	270	ASN	2.3
1	A	212	GLY	2.3
1	A	547	ALA	2.3
1	A	725	GLY	2.3
1	A	420	GLY	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	550	GLU	2.2
1	A	213	ALA	2.2
1	A	754	GLN	2.2
1	A	18	LEU	2.2
1	A	45	VAL	2.1
1	A	426	ARG	2.1
1	A	768	HIS	2.1
1	A	171	CYS	2.1
1	A	211	GLN	2.1
1	A	15	VAL	2.0
1	A	69	ARG	2.0
1	A	436	VAL	2.0
1	A	549	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	DMS	A	904	4/4	0.36	0.27	78,79,82,85	0
3	DMS	A	903	4/4	0.74	0.18	55,56,57,60	0
2	KS3	A	902	26/26	0.80	0.15	35,43,57,65	0
2	KS3	A	901	26/26	0.97	0.05	17,18,20,21	0
4	PLP	A	905	15/16	0.98	0.05	17,19,20,21	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around KS3 A 902:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around KS3 A 901:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.