



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2026 – 07:45 PM UTC

PDB ID : 5LRY / pdb_00005lry
Title : E coli [NiFe] Hydrogenase Hyd-1 mutant E28D
Authors : Carr, S.B.; Phillips, S.E.V.; Evans, R.M.; Brooke, E.J.; Armstrong, F.A.
Deposited on : 2016-08-22
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.40-1.40)
Clashscore	190562	2660 (1.40-1.40)
Ramachandran outliers	187476	2611 (1.40-1.40)
Sidechain outliers	187428	2610 (1.40-1.40)
RSRZ outliers	180081	2561 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	S	335	% 76% 21%
1	T	335	74% 21%
2	L	582	% 97% .
2	M	582	% 96% .

2 Entry composition

There are 13 unique types of molecules in this entry. The entry contains 15125 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hydrogenase-1 small chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	S	263	Total	C	N	O	S	0	10	0
			2090	1327	358	385	20			
1	T	263	Total	C	N	O	S	0	12	0
			2113	1340	366	386	21			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
S	328	ARG	-	expression tag	UNP P69740
S	329	SER	-	expression tag	UNP P69740
S	330	HIS	-	expression tag	UNP P69740
S	331	HIS	-	expression tag	UNP P69740
S	332	HIS	-	expression tag	UNP P69740
S	333	HIS	-	expression tag	UNP P69740
S	334	HIS	-	expression tag	UNP P69740
S	335	HIS	-	expression tag	UNP P69740
T	328	ARG	-	expression tag	UNP P69740
T	329	SER	-	expression tag	UNP P69740
T	330	HIS	-	expression tag	UNP P69740
T	331	HIS	-	expression tag	UNP P69740
T	332	HIS	-	expression tag	UNP P69740
T	333	HIS	-	expression tag	UNP P69740
T	334	HIS	-	expression tag	UNP P69740
T	335	HIS	-	expression tag	UNP P69740


- Molecule 2 is a protein called Hydrogenase-1 large chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	581	Total	C	N	O	S	0	25	0
			4684	2975	814	865	30			
2	M	581	Total	C	N	O	S	0	28	0
			4702	2992	814	866	30			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
L	28	ASP	GLU	conflict	UNP P0ACD8
M	28	ASP	GLU	conflict	UNP P0ACD8

- Molecule 3 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe_4S_4).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	S	1	Total	Fe	S	0	0
			8	4	4		
3	T	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 4 is FE3-S4 CLUSTER (CCD ID: F3S) (formula: Fe_3S_4).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	S	1	Total	Fe	S	0	0
			7	3	4		
4	T	1	Total	Fe	S	0	0
			7	3	4		

- Molecule 5 is FE4-S3 CLUSTER (CCD ID: SF3) (formula: Fe_4S_3).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	S	1	Total	Fe	S	0	0
			7	4	3		
5	T	1	Total	Fe	S	0	0
			7	4	3		

- Molecule 6 is DODECYL-BETA-D-MALTOSE (CCD ID: LMT) (formula: C₂₄H₄₆O₁₁).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	S	1	Total	C	O	0	0
			14	13	1		
6	T	1	Total	C	O	0	0
			14	13	1		

- Molecule 7 is CHLORIDE ION (CCD ID: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	S	1	Total	Cl	0	0
			1	1		
7	T	1	Total	Cl	0	0
			1	1		

- Molecule 8 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	S	1	Total	O	S	0	0
			5	4	1		
8	L	1	Total	O	S	0	0
			5	4	1		
8	M	1	Total	O	S	0	0
			5	4	1		

- Molecule 9 is CARBONMONOXIDE-(DICYANO) IRON (CCD ID: FCO) (formula: C₃FeN₂O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	L	1	Total	C	Fe	N	O	0	0
			7	3	1	2	1		
9	M	1	Total	C	Fe	N	O	0	0
			7	3	1	2	1		

- Molecule 10 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	L	1	Total	Ni	0	0
			1	1		
10	M	1	Total	Ni	0	0
			1	1		

- Molecule 11 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	L	1	Total	Mg	0	0
			1	1		
11	M	1	Total	Mg	0	0
			1	1		

- Molecule 12 is LITHIUM ION (CCD ID: LI) (formula: Li).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	M	1	Total	Li	0	0
			1	1		

- Molecule 13 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
13	S	226	Total	O	0	0
			226	226		
13	L	481	Total	O	0	0
			481	481		
13	T	200	Total	O	0	0
			200	200		
13	M	521	Total	O	0	0
			521	521		

- Molecule 1: Hydrogenase-1 small chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	94.76Å 98.75Å 185.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	92.79 – 1.40 92.62 – 1.40	Depositor EDS
% Data completeness (in resolution range)	99.2 (92.79-1.40) 99.2 (92.62-1.40)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.79 (at 1.40Å)	Xtriage
Refinement program	REFMAC 5.8.0155	Depositor
R, R_{free}	0.146 , 0.163 0.147 , 0.163	Depositor DCC
R_{free} test set	16837 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	10.2	Xtriage
Anisotropy	0.103	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 39.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.018 for k,h,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	15125	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SF3, FCO, F3S, LI, SO4, SF4, NI, CSD, LMT, MG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	S	1.11	1/2169 (0.0%)	0.95	3/2943 (0.1%)
1	T	1.11	2/2195 (0.1%)	0.93	2/2977 (0.1%)
2	L	1.08	0/4865	0.88	0/6615
2	M	1.08	2/4889 (0.0%)	0.88	1/6647 (0.0%)
All	All	1.09	5/14118 (0.0%)	0.90	6/19182 (0.0%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	S	260	ARG	CZ-NH1	6.01	1.41	1.32
2	M	244	ASP	N-CA	5.88	1.52	1.46
1	T	260	ARG	CZ-NH1	5.36	1.40	1.32
1	T	188	ASP	CG-OD2	-5.10	1.15	1.25
2	M	299	LYS	CA-C	5.01	1.56	1.52

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	T	193	ARG	CG-CD-NE	-6.13	98.52	112.00
1	S	193	ARG	CG-CD-NE	-5.77	99.30	112.00
2	M	530	VAL	N-CA-CB	-5.37	108.13	111.83
1	T	260	ARG	CB-CA-C	-5.35	102.98	111.17
1	S	38	SER	N-CA-C	5.11	120.90	114.31
1	S	78	ASN	N-CA-C	5.03	114.08	108.25

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	S	2090	0	2039	5	0
1	T	2113	0	2067	10	0
2	L	4684	0	4603	9	0
2	M	4702	0	4638	25	0
3	S	8	0	0	0	0
3	T	8	0	0	0	0
4	S	7	0	0	0	0
4	T	7	0	0	0	0
5	S	7	0	0	0	0
5	T	7	0	0	0	0
6	S	14	0	25	0	0
6	T	14	0	25	0	0
7	S	1	0	0	0	0
7	T	1	0	0	0	0
8	L	5	0	0	0	0
8	M	5	0	0	0	0
8	S	5	0	0	0	0
9	L	7	0	0	0	0
9	M	7	0	0	0	0
10	L	1	0	0	0	0
10	M	1	0	0	0	0
11	L	1	0	0	0	0
11	M	1	0	0	0	0
12	M	1	0	0	0	0
13	L	481	0	0	4	0
13	M	521	0	0	5	0
13	S	226	0	0	3	0
13	T	200	0	0	3	0
All	All	15125	0	13397	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:M:434[B]:MET:HE1	2:M:457[B]:HIS:CD2	1.33	1.62
2:M:434[B]:MET:HE3	2:M:457[B]:HIS:CE1	1.67	1.30
2:M:434[B]:MET:CE	2:M:457[B]:HIS:NE2	1.96	1.28
2:M:434[B]:MET:CE	2:M:457[B]:HIS:CD2	2.18	1.25
2:M:434[B]:MET:CE	2:M:457[B]:HIS:CE1	2.35	1.04
1:T:70:LYS:NZ	13:T:501:HOH:O	1.92	1.02
2:M:434[B]:MET:HE1	2:M:457[B]:HIS:CG	1.96	1.00
2:L:488[B]:GLU:OE2	13:L:701:HOH:O	1.99	0.78
2:M:434[B]:MET:CE	2:M:457[B]:HIS:CG	2.65	0.75
2:L:457[A]:HIS:ND1	13:L:702:HOH:O	2.20	0.74
2:M:556:LYS:NZ	13:M:701:HOH:O	2.27	0.67
1:T:61[B]:GLU:OE2	1:T:101:ARG:NH2	2.30	0.64
2:M:434[B]:MET:HE2	2:M:457[B]:HIS:NE2	2.09	0.63
2:L:52[B]:MET:HG2	13:L:1073:HOH:O	1.99	0.62
2:M:486[A]:SER:OG	2:M:488[A]:GLU:OE2	2.12	0.62
2:M:52[B]:MET:HG2	13:M:1080:HOH:O	2.01	0.61
2:L:486:SER:OG	2:L:488[B]:GLU:OE1	2.13	0.61
2:M:60[B]:LEU:HD11	2:M:72:VAL:CG1	2.34	0.58
2:L:169:ARG:O	2:L:172:LYS:HE3	2.04	0.57
1:S:234[A]:ARG:NH2	1:S:244:GLN:HE22	2.04	0.55
2:M:434[B]:MET:HE3	2:M:457[B]:HIS:ND1	2.17	0.54
1:S:61[A]:GLU:HG2	13:S:613:HOH:O	2.08	0.54
1:T:61[B]:GLU:CD	1:T:101:ARG:HH12	2.18	0.52
2:M:162:TYR:OH	2:M:208[B]:GLU:OE2	2.23	0.51
2:L:254[B]:MET:HE1	1:T:166:PHE:HE1	1.77	0.49
1:T:76[B]:GLU:HG3	13:T:532:HOH:O	2.12	0.49
2:L:135:LEU:HD22	2:L:187:TRP:CD1	2.48	0.48
1:T:89:ILE:HG22	13:M:704:HOH:O	2.13	0.48
2:L:530:VAL:CG1	2:L:531:PRO:HD2	2.45	0.46
2:M:30:HIS:CD2	2:M:52[B]:MET:SD	3.09	0.46
2:L:169:ARG:NH1	13:L:708:HOH:O	2.45	0.45
1:T:217:TYR:HE1	1:T:267:VAL:HG12	1.81	0.45
2:M:280:LEU:HG	13:M:705:HOH:O	2.17	0.45
2:M:434[B]:MET:CE	2:M:457[B]:HIS:ND1	2.73	0.45
2:M:254[B]:MET:HE2	2:M:254[B]:MET:HA	1.98	0.45
2:M:30:HIS:CG	2:M:52[B]:MET:SD	3.11	0.44
2:M:488[A]:GLU:H	2:M:488[A]:GLU:CD	2.25	0.44
1:T:186:ILE:HD11	1:T:228:ASN:HB3	2.00	0.44
1:S:76[B]:GLU:HG3	13:S:573:HOH:O	2.18	0.44
1:T:61[B]:GLU:HG2	13:T:578:HOH:O	2.17	0.43
1:S:234[A]:ARG:NH1	1:S:244:GLN:OE1	2.52	0.43
2:M:530:VAL:CG1	2:M:531:PRO:HD2	2.50	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:T:89:ILE:HB	2:M:51:THR:HB	2.01	0.42
2:M:457[B]:HIS:NE2	13:M:705:HOH:O	2.37	0.42
2:M:540:ASP:HB2	2:M:541:PRO:CD	2.50	0.41
1:S:101[B]:ARG:HD3	13:S:678:HOH:O	2.20	0.41
2:M:445:ILE:O	2:M:450:GLY:HA3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	S	271/335 (81%)	259 (96%)	12 (4%)	0	100	100
1	T	273/335 (82%)	260 (95%)	13 (5%)	0	100	100
2	L	603/582 (104%)	588 (98%)	15 (2%)	0	100	100
2	M	606/582 (104%)	590 (97%)	16 (3%)	0	100	100
All	All	1753/1834 (96%)	1697 (97%)	56 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	S	226/274 (82%)	226 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	T	228/274 (83%)	227 (100%)	1 (0%)	84	67
2	L	504/480 (105%)	501 (99%)	3 (1%)	78	56
2	M	507/480 (106%)	507 (100%)	0	100	100
All	All	1465/1508 (97%)	1461 (100%)	4 (0%)	86	70

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	L	172	LYS
2	L	433	ARG
2	L	524	ASP
1	T	242	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	S	68	ASN
2	L	150	GLN
2	L	343	GLN
2	L	360	GLN
1	T	55	GLN
2	M	4	GLN
2	M	150	GLN
2	M	213	GLN
2	M	261	GLN
2	M	392	GLN
2	M	467	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	CSD	M	576	2,10	4,7,8	2.81	1 (25%)	1,8,10	1.06	0
2	CSD	L	576	2,10	4,7,8	2.33	1 (25%)	1,8,10	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CSD	M	576	2,10	-	0/2/6/8	-
2	CSD	L	576	2,10	-	0/2/6/8	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	M	576	CSD	OD1-SG	5.51	1.52	1.47
2	L	576	CSD	OD1-SG	4.28	1.51	1.47

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 7 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SF3	S	403	1,13	0,8,8	-	-	-		
8	SO4	L	601	-	4,4,4	0.44	0	6,6,6	0.32	0
6	LMT	S	404	-	13,13,36	0.45	0	12,12,47	0.60	0
3	SF4	S	401	1	0,12,12	-	-	-		
9	FCO	L	602	2,13	0,6,6	-	-	-		
9	FCO	M	601	2,13	0,6,6	-	-	-		
4	F3S	S	402	1	0,9,9	-	-	-		
8	SO4	S	406	-	4,4,4	0.47	0	6,6,6	0.24	0
8	SO4	M	604	-	4,4,4	0.49	0	6,6,6	0.34	0
3	SF4	T	401	1	0,12,12	-	-	-		
6	LMT	T	404	-	13,13,36	0.36	0	12,12,47	0.37	0
4	F3S	T	402	1	0,9,9	-	-	-		
5	SF3	T	403	1,13	0,8,8	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	SF3	S	403	1,13	-	-	0/2/2/2
6	LMT	S	404	-	-	3/11/11/61	-
3	SF4	S	401	1	-	-	0/6/5/5
4	F3S	S	402	1	-	-	0/3/3/3
3	SF4	T	401	1	-	-	0/6/5/5
6	LMT	T	404	-	-	0/11/11/61	-
4	F3S	T	402	1	-	-	0/3/3/3
5	SF3	T	403	1,13	-	-	0/2/2/2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	S	404	LMT	O1'-C1-C2-C3
6	S	404	LMT	C5-C6-C7-C8
6	S	404	LMT	C2-C3-C4-C5

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	S	263/335 (78%)	-0.41	2 (0%) 82 85	5, 10, 17, 27	10 (3%)
1	T	263/335 (78%)	-0.31	1 (0%) 88 91	6, 11, 18, 30	12 (4%)
2	L	580/582 (99%)	-0.37	4 (0%) 84 87	5, 11, 22, 38	25 (4%)
2	M	580/582 (99%)	-0.44	4 (0%) 84 87	5, 11, 19, 35	28 (4%)
All	All	1686/1834 (91%)	-0.39	11 (0%) 84 87	5, 11, 20, 38	75 (4%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	T	267	VAL	4.5
1	S	267	VAL	3.9
2	L	285	PHE	3.3
2	M	143	ARG	2.5
2	L	172	LYS	2.5
2	M	285	PHE	2.3
2	L	425	ALA	2.2
2	L	437	ALA	2.2
2	M	2	SER	2.2
2	M	174	VAL	2.2
1	S	260	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CSD	L	576	8/9	0.98	0.05	8,9,10,11	2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CSD	M	576	8/9	0.99	0.05	8,9,10,11	2

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	LMT	S	404	14/35	0.76	0.19	25,27,32,32	0
8	SO4	S	406	5/5	0.77	0.12	60,63,65,67	0
8	SO4	M	604	5/5	0.77	0.17	33,34,35,35	5
6	LMT	T	404	14/35	0.78	0.17	27,29,39,40	0
12	LI	M	605	1/1	0.94	0.06	13,13,13,13	0
8	SO4	L	601	5/5	0.96	0.07	14,15,17,17	5
7	CL	T	405	1/1	0.98	0.06	14,14,14,14	0
7	CL	S	405	1/1	0.99	0.05	14,14,14,14	0
5	SF3	S	403	7/7	0.99	0.05	9,11,12,16	0
5	SF3	T	403	7/7	0.99	0.04	10,11,13,17	0
4	F3S	S	402	7/7	1.00	0.02	8,8,9,9	0
4	F3S	T	402	7/7	1.00	0.02	8,9,9,10	0
3	SF4	S	401	8/8	1.00	0.01	7,7,7,8	0
9	FCO	L	602	7/7	1.00	0.02	7,7,7,8	0
9	FCO	M	601	7/7	1.00	0.03	7,7,8,8	0
10	NI	L	603	1/1	1.00	0.06	11,11,11,11	0
10	NI	M	602	1/1	1.00	0.05	11,11,11,11	0
11	MG	L	604	1/1	1.00	0.04	3,3,3,3	0
11	MG	M	603	1/1	1.00	0.04	3,3,3,3	0
3	SF4	T	401	8/8	1.00	0.01	7,7,7,8	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.