



wwPDB EM Validation Summary Report ⓘ

Mar 8, 2026 – 05:16 PM UTC

PDB ID : 6LRS / pdb_00006lrs
EMDB ID : EMD-0960
Title : Cryo-EM structure of RbcL8-RbcS4 from Anabaena sp. PCC 7120
Authors : Xia, L.Y.; Jiang, Y.L.; Kong, W.W.; Sun, H.; Li, W.F.; Chen, Y.; Zhou, C.Z.
Deposited on : 2020-01-16
Resolution : 3.37 Å(reported)
Based on initial model : 1RBL

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDb archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.









Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	14287 (2.87 - 3.87)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	Q	109	70% 77% 20% .
1	R	109	69% 77% 20% .
1	S	109	69% 77% 20% .
1	T	109	69% 77% 20% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	A	476	
2	B	476	
2	C	476	
2	D	476	
2	E	476	
2	F	476	
2	G	476	
2	H	476	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 30156 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribulose biphosphate carboxylase small chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	T	106	Total	C	N	O	S	0	0
			877	571	144	159	3		
1	S	106	Total	C	N	O	S	0	0
			877	571	144	159	3		
1	R	106	Total	C	N	O	S	0	0
			877	571	144	159	3		
1	Q	106	Total	C	N	O	S	0	0
			877	571	144	159	3		


- Molecule 2 is a protein called Ribulose biphosphate carboxylase large chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	G	413	Total	C	N	O	S	0	0
			3229	2056	566	595	12		
2	H	439	Total	C	N	O	S	0	0
			3433	2180	608	632	13		
2	E	413	Total	C	N	O	S	0	0
			3229	2056	566	595	12		
2	F	439	Total	C	N	O	S	0	0
			3433	2180	608	632	13		
2	C	413	Total	C	N	O	S	0	0
			3229	2056	566	595	12		
2	D	439	Total	C	N	O	S	0	0
			3433	2180	608	632	13		
2	A	413	Total	C	N	O	S	0	0
			3229	2056	566	595	12		
2	B	439	Total	C	N	O	S	0	0
			3433	2180	608	632	13		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ribulose biphosphate carboxylase small chain




- Molecule 1: Ribulose biphosphate carboxylase small chain



- Molecule 1: Ribulose biphosphate carboxylase small chain




- Molecule 1: Ribulose bisphosphate carboxylase small chain






• Molecule 2: Ribulose biphosphate carboxylase large chain


Chain G: 30% 64% 23% 13%




• Molecule 2: Ribulose bisphosphate carboxylase large chain

Chain H: 21% 65% 27% 8%







• Molecule 2: Ribulose biphosphate carboxylase large chain




• Molecule 2: Ribulose biphosphate carboxylase large chain







• Molecule 2: Ribulose biphosphate carboxylase large chain




• Molecule 2: Ribulose bisphosphate carboxylase large chain





• Molecule 2: Ribulose biphosphate carboxylase large chain



• Molecule 2: Ribulose biphosphate carboxylase large chain



R422	L425	V429	N433	E434	G435	R436	N437	L438	A439	R440	E441	G442	N443	D444	R447	E448	A449	A450	K451	W452	S453	P454	E455	L456	A457	C460	GLU	LEU	TRP	LYS	GLU	ILE	LYS	PHE	GLU	PHE	GLU	ALA	MET	ASP	THR	VAL													
L336	E337	G338	E339	R340	T343	F346	L349	L350	E356	S360	R361	Q367	A370	S371	L372	P373	G374	V375	V378	I383	H384	V385	W386	H387	A390	L391	V392	E393	D397	V400	G404	G405	G406	T407	L408	G409	H410	P411	A415	P416	G417	A418	T419	A420	N421										
R254	I265	I266	M267	H268	A273	T276	T279	T280	R283	R286	L291	L292	H293	I294	H295	R296	A297	M298	H299	A300	V301	I302	D303	R304	H308	G309	I310	H311	F312	R313	V314	L315	A316	K317	A318	L319	R320	L321	S322	G323	G324	D325	H326	I327	H328	T331	V332	V333	G334	K335					
P152	G155	I156	Q157	V158	E159	R160	D161	K162	L163	Y166	G167	R168	P169	L170	L171	G172	C173	T174	I175	K176	P177	K178	L179	G180	Y186	G187	R188	A189	V190	Y191	E192	R195	F200	T201	K202	D203	D204	E205	N206	I207	N208	S209	W215	V222	L241	N242	V243	T244	E249						
L74	L75	T76	D77	L78	D79	R80	D87	I88	E89	P90	V91	G93	E94	D95	N96	Q97	Y101	P105	L108	F109	E110	S113	I114	T115	N116	V117	S120	I121	N124	V125	F126	G127	F128	K129	A130	A133	L134	R135	L136	R140	F141	P142	V143	A144	Y145	I146	Q150	G151							
MET	SER	TYR	ALA	GLN	THR	LYS	THR	GLN	LYS	THR	LYS	SER	GLY	TYR	LYS	ALA	GLY	VAL	GLN	ASP	TYR	R22	L23	T27	P28	D29	Y30	T31	P32	D36	I37	L38	R42	Q46	P47	G48	V49	P50	F51	E52	E53	A54	E61	S62	S63	T64	G65	T66	W67	T68	T69	V70	W71	T72	D73

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C4	Depositor
Number of particles used	340899	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.187	Depositor
Minimum map value	-0.117	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.009	Depositor
Recommended contour level	0.04	Depositor
Map size ($\text{\AA}$)	249.59999, 249.59999, 249.59999	wwPDB
Map dimensions	240, 240, 240	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	1.04, 1.04, 1.04	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	Q	0.28	0/901	0.69	2/1226 (0.2%)
1	R	0.28	0/901	0.69	2/1226 (0.2%)
1	S	0.28	0/901	0.69	2/1226 (0.2%)
1	T	0.28	0/901	0.69	2/1226 (0.2%)
2	A	0.27	0/3309	0.61	0/4497
2	B	0.29	0/3516	0.59	0/4777
2	C	0.27	0/3309	0.61	0/4497
2	D	0.29	0/3516	0.59	0/4777
2	E	0.27	0/3309	0.61	0/4497
2	F	0.29	0/3516	0.59	0/4777
2	G	0.27	0/3309	0.61	0/4497
2	H	0.29	0/3516	0.59	0/4777
All	All	0.28	0/30904	0.61	8/42000 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	Q	0	1
1	R	0	1
1	S	0	1
1	T	0	1
2	A	0	1
2	B	0	1
2	C	0	1
2	D	0	1
2	E	0	1
2	F	0	1
2	G	0	1
2	H	0	1
All	All	0	12

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	T	36	TYR	CA-C-N	7.48	131.48	123.25
1	T	36	TYR	C-N-CA	7.48	131.48	123.25
1	Q	36	TYR	CA-C-N	7.48	131.47	123.25
1	Q	36	TYR	C-N-CA	7.48	131.47	123.25
1	S	36	TYR	CA-C-N	7.43	131.42	123.25

There are no chirality outliers.

5 of 12 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	E	177	PRO	Peptide
2	G	177	PRO	Peptide
2	H	75	LEU	Peptide
1	S	89	GLY	Peptide
1	T	89	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	Q	877	0	882	12	0
1	R	877	0	882	12	0
1	S	877	0	882	12	0
1	T	877	0	882	12	0
2	A	3229	0	3165	81	0
2	B	3433	0	3368	82	0
2	C	3229	0	3165	85	0
2	D	3433	0	3368	85	0
2	E	3229	0	3165	80	0
2	F	3433	0	3368	83	0
2	G	3229	0	3165	77	0
2	H	3433	0	3368	84	0
All	All	30156	0	29660	644	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 644 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:209:SER:HB3	2:A:215:TRP:HD1	1.56	0.70
2:H:327:ILE:HD12	2:H:350:LEU:HD11	1.72	0.70
2:F:327:ILE:HD12	2:F:350:LEU:HD11	1.72	0.70
2:D:327:ILE:HD12	2:D:350:LEU:HD11	1.72	0.69
2:B:327:ILE:HD12	2:B:350:LEU:HD11	1.72	0.69

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	Q	104/109 (95%)	94 (90%)	10 (10%)	0	100	100
1	R	104/109 (95%)	94 (90%)	10 (10%)	0	100	100
1	S	104/109 (95%)	94 (90%)	10 (10%)	0	100	100
1	T	104/109 (95%)	94 (90%)	10 (10%)	0	100	100
2	A	411/476 (86%)	376 (92%)	35 (8%)	0	100	100
2	B	437/476 (92%)	418 (96%)	19 (4%)	0	100	100
2	C	411/476 (86%)	376 (92%)	35 (8%)	0	100	100
2	D	437/476 (92%)	419 (96%)	18 (4%)	0	100	100
2	E	411/476 (86%)	377 (92%)	34 (8%)	0	100	100
2	F	437/476 (92%)	418 (96%)	19 (4%)	0	100	100
2	G	411/476 (86%)	376 (92%)	35 (8%)	0	100	100
2	H	437/476 (92%)	417 (95%)	20 (5%)	0	100	100
All	All	3808/4244 (90%)	3553 (93%)	255 (7%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	Q	99/102 (97%)	98 (99%)	1 (1%)	68	75
1	R	99/102 (97%)	98 (99%)	1 (1%)	68	75
1	S	99/102 (97%)	98 (99%)	1 (1%)	68	75
1	T	99/102 (97%)	98 (99%)	1 (1%)	68	75
2	A	333/386 (86%)	332 (100%)	1 (0%)	86	84
2	B	353/386 (92%)	352 (100%)	1 (0%)	86	84
2	C	333/386 (86%)	332 (100%)	1 (0%)	86	84
2	D	353/386 (92%)	352 (100%)	1 (0%)	86	84
2	E	333/386 (86%)	332 (100%)	1 (0%)	86	84
2	F	353/386 (92%)	352 (100%)	1 (0%)	86	84
2	G	333/386 (86%)	332 (100%)	1 (0%)	86	84
2	H	353/386 (92%)	352 (100%)	1 (0%)	86	84
All	All	3140/3496 (90%)	3128 (100%)	12 (0%)	81	82

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	378	VAL
2	D	174	THR
2	B	174	THR
1	Q	43	ASN
1	S	43	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 95 such sidechains are listed below:

Mol	Chain	Res	Type
2	C	308	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	Q	2	GLN
2	C	353	ASN
2	D	230	GLN
2	A	97	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-0960. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 120



Y Index: 120




Z Index: 120


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 105



Y Index: 105




Z Index: 111


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.04. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 95 nm³; this corresponds to an approximate mass of 86 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.297 Å⁻¹

8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-0960 and PDB model 6LRS. Per-residue inclusion information can be found in section 3 on page 5.

9.1 Map-model overlay [i](#)


This section was not generated.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.04).

9.4 Atom inclusion [i](#)



At the recommended contour level, 61% of all backbone atoms, 51% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.04) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.5070	0.4400
A	0.5050	0.4360
B	0.5650	0.4610
C	0.5050	0.4370
D	0.5640	0.4620
E	0.5040	0.4360
F	0.5640	0.4610
G	0.5040	0.4360
H	0.5630	0.4600
Q	0.2910	0.3730
R	0.2970	0.3740
S	0.2960	0.3750
T	0.2940	0.3720

1.0

0.0

<0.0