



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:30 PM UTC

PDB ID : 3LSS / pdb_00003lss
Title : Trypanosoma brucei seryl-tRNA synthetase in complex with ATP
Authors : Larson, E.T.; Merritt, E.A.; Medical Structural Genomics of Pathogenic Protozoa (MSGPP)
Deposited on : 2010-02-12
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	484	17% 85% 6% 10%
1	B	484	15% 90% 6% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MLI	A	602	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7683 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Seryl-tRNA synthetase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	438	Total	C	N	O	S	0	9	0
			3509	2203	613	662	31			
1	B	468	Total	C	N	O	S	0	15	0
			3756	2355	653	718	30			

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	expression tag	UNP Q384V4
A	-3	PRO	-	expression tag	UNP Q384V4
A	-2	GLY	-	expression tag	UNP Q384V4
A	-1	SER	-	expression tag	UNP Q384V4
A	0	MET	-	expression tag	UNP Q384V4
B	-4	GLY	-	expression tag	UNP Q384V4
B	-3	PRO	-	expression tag	UNP Q384V4
B	-2	GLY	-	expression tag	UNP Q384V4
B	-1	SER	-	expression tag	UNP Q384V4
B	0	MET	-	expression tag	UNP Q384V4

- Molecule 2 is ADENOSINE-5'-TRIPHOSPHATE (CCD ID: ATP) (formula: C₁₀H₁₆N₅O₁₃P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 31	C 10	N 5	O 13	P 3	0	0
2	B	1	Total 31	C 10	N 5	O 13	P 3	0	0

- Molecule 3 is MALONATE ION (CCD ID: MLI) (formula: $\text{C}_3\text{H}_2\text{O}_4$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 7 3 4	0	0
3	A	1	Total C O 7 3 4	0	0

- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total 1	Na 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	167	Total 167	O 167	0	1
5	B	174	Total 174	O 174	0	1


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Seryl-tRNA synthetase



• Molecule 1: Seryl-tRNA synthetase



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	175.65Å 175.65Å 248.52Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	34.57 – 1.95 34.57 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.9 (34.57-1.95) 99.9 (34.57-1.95)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.26 (at 1.95Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.168 , 0.195 0.179 , 0.202	Depositor DCC
R_{free} test set	5392 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	27.7	Xtriage
Anisotropy	0.380	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 56.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7683	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, MLI, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	0/3601	0.83	1/4844 (0.0%)
1	B	0.75	0/3879	0.82	2/5223 (0.0%)
All	All	0.74	0/7480	0.83	3/10067 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	150	GLU	N-CA-C	5.84	117.65	111.28
1	B	34	ILE	CB-CA-C	-5.45	104.70	112.22
1	B	230	PHE	CB-CA-C	-5.17	100.08	109.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3509	0	3498	16	0
1	B	3756	0	3733	22	0
2	A	31	0	12	1	0
2	B	31	0	12	1	0
3	A	14	0	4	3	0
4	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	167	0	0	2	0
5	B	174	0	0	0	0
All	All	7683	0	7259	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:214:LEU:HD13	1:A:290[B]:MET:SD	2.08	0.94
1:B:2:VAL:CG2	1:B:130[B]:ASP:OD1	2.27	0.82
1:B:2:VAL:HG21	1:B:130[B]:ASP:OD1	1.85	0.76
1:B:2:VAL:HB	1:B:133[B]:MET:SD	2.28	0.73
1:B:134:LEU:HD11	1:B:242:LEU:HD12	1.71	0.72
1:B:252[B]:VAL:HG23	1:B:259:LYS:HB2	1.77	0.67
1:B:2:VAL:HG22	1:B:130[B]:ASP:OD1	1.95	0.65
1:B:2:VAL:CG2	1:B:133[B]:MET:SD	2.84	0.64
1:A:149:ASP:OD2	1:A:152:THR:HG22	2.00	0.62
1:A:219:LYS:NZ	5:A:911:HOH:O	2.35	0.60
1:A:50:ALA:O	1:A:54:LEU:HD12	2.06	0.55
1:B:31:VAL:HA	1:B:34:ILE:HD12	1.89	0.55
1:B:2:VAL:CB	1:B:133[B]:MET:SD	2.95	0.54
1:B:2:VAL:HG23	1:B:133[B]:MET:SD	2.48	0.54
1:A:276:ARG:HH11	3:A:602:MLI:H11	1.72	0.53
1:B:372:LEU:HB3	1:B:386:VAL:HB	1.91	0.52
1:A:290[B]:MET:HG2	1:A:317:ILE:HD12	1.92	0.52
1:A:252:VAL:HG22	1:B:252[B]:VAL:HG12	1.93	0.51
1:A:276:ARG:HD3	3:A:602:MLI:H11	1.93	0.50
1:A:231:LEU:CD2	1:B:190:MET:HG2	2.42	0.50
1:A:8:PHE:CE2	1:A:133[A]:MET:HE1	2.47	0.49
1:B:84:GLU:O	1:B:88[A]:ASN:ND2	2.45	0.49
1:A:172:VAL:HG21	1:A:306:LEU:HB3	1.94	0.49
1:A:282:LYS:O	5:A:960:HOH:O	2.20	0.48
1:A:157:VAL:HG12	1:A:158:ARG:HG3	1.96	0.47
1:B:2:VAL:HG21	1:B:130[B]:ASP:CG	2.37	0.47
1:B:247:GLU:HA	1:B:298:ALA:HB3	1.96	0.47
1:A:372:LEU:HB3	1:A:386:VAL:HB	1.97	0.47
2:B:501:ATP:O2G	2:B:501:ATP:O2A	2.33	0.46
3:A:601:MLI:O7	3:A:601:MLI:O9	2.33	0.45
1:B:3:LEU:O	1:B:133[B]:MET:SD	2.76	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:187:VAL:O	1:B:190:MET:HB3	2.19	0.43
1:A:384:GLU:OE2	2:A:501:ATP:O3G	2.39	0.41
1:A:391:CYS:O	1:A:392:THR:C	2.63	0.41
1:B:279:THR:HA	1:B:404:GLY:O	2.20	0.41
1:B:294:PHE:HA	1:B:312:HIS:O	2.22	0.40
1:B:271:ALA:O	1:B:274:ARG:HG2	2.22	0.40
1:B:141:HIS:HB2	1:B:331:TRP:CH2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	441/484 (91%)	429 (97%)	12 (3%)	0	100	100
1	B	481/484 (99%)	467 (97%)	13 (3%)	1 (0%)	43	36
All	All	922/968 (95%)	896 (97%)	25 (3%)	1 (0%)	48	41

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	28	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	387/412 (94%)	384 (99%)	3 (1%)	73	74
1	B	416/412 (101%)	412 (99%)	4 (1%)	68	67
All	All	803/824 (98%)	796 (99%)	7 (1%)	73	70

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	108	SER
1	A	305	THR
1	A	426	LEU
1	B	53[A]	LYS
1	B	53[B]	LYS
1	B	152	THR
1	B	264	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	81	GLN
1	B	95	GLN
1	B	345	ASN
1	B	356	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MLI	A	601	-	6,6,6	2.07	2 (33%)	7,7,7	1.14	0
3	MLI	A	602	-	6,6,6	1.70	2 (33%)	7,7,7	1.14	0
2	ATP	A	501	-	32,33,33	1.67	7 (21%)	48,52,52	1.62	11 (22%)
2	ATP	B	501	-	32,33,33	1.63	7 (21%)	48,52,52	1.74	11 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MLI	A	601	-	-	0/4/4/4	-
3	MLI	A	602	-	-	2/4/4/4	-
2	ATP	A	501	-	-	5/22/38/38	0/3/3/3
2	ATP	B	501	-	-	5/22/38/38	0/3/3/3

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	ATP	C5-C4	4.87	1.47	1.39
2	B	501	ATP	C5-C4	4.77	1.47	1.39
2	A	501	ATP	C8-N7	3.70	1.38	1.31
2	B	501	ATP	PB-O3B	3.66	1.63	1.59
2	A	501	ATP	PB-O3B	3.28	1.63	1.59
2	B	501	ATP	PB-O3A	3.27	1.63	1.59
3	A	601	MLI	C1-C2	3.22	1.56	1.51
2	A	501	ATP	PB-O3A	3.16	1.62	1.59
3	A	601	MLI	C1-C3	2.91	1.55	1.51
2	B	501	ATP	C8-N7	2.61	1.36	1.31
2	A	501	ATP	PA-O3A	2.50	1.62	1.59
3	A	602	MLI	C1-C3	2.36	1.54	1.51
2	B	501	ATP	PA-O3A	2.24	1.61	1.59
2	A	501	ATP	C4-N9	-2.18	1.33	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	ATP	C5-N7	-2.16	1.35	1.39
3	A	602	MLI	C1-C2	2.09	1.54	1.51
2	B	501	ATP	C4-N9	-2.04	1.33	1.37
2	B	501	ATP	C5-C6	2.01	1.46	1.41

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	ATP	C5-C4-N3	-4.37	120.70	126.72
2	A	501	ATP	C5-C4-N3	-4.35	120.73	126.72
2	B	501	ATP	N3-C4-N9	4.26	134.42	127.17
2	A	501	ATP	N3-C4-N9	3.88	133.77	127.17
2	B	501	ATP	C4-N9-C8	3.75	109.68	105.74
2	A	501	ATP	N3-C2-N1	-3.72	122.94	128.58
2	B	501	ATP	N3-C2-N1	-3.49	123.30	128.58
2	A	501	ATP	C4-N9-C8	3.26	109.16	105.74
2	A	501	ATP	N6-C6-N1	3.22	125.55	118.38
2	B	501	ATP	O2A-PA-O3A	2.98	115.32	107.27
2	A	501	ATP	C2-N3-C4	2.85	118.80	111.83
2	B	501	ATP	N6-C6-N1	2.69	124.37	118.38
2	A	501	ATP	C2-N1-C6	2.67	123.12	118.73
2	B	501	ATP	C2-N3-C4	2.63	118.27	111.83
2	A	501	ATP	C5-C6-N6	-2.48	117.16	123.29
2	B	501	ATP	C2-N1-C6	2.46	122.78	118.73
2	B	501	ATP	N9-C8-N7	-2.31	110.66	113.94
2	A	501	ATP	N9-C8-N7	-2.29	110.68	113.94
2	A	501	ATP	O3B-PG-O1G	-2.11	99.94	111.04
2	B	501	ATP	C5-C6-N6	-2.11	118.07	123.29
2	B	501	ATP	O2B-PB-O3B	2.08	112.91	107.27
2	A	501	ATP	C4-N9-C1'	-2.06	121.81	126.63

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	501	ATP	C5'-O5'-PA-O1A
2	A	501	ATP	PB-O3B-PG-O2G
2	A	501	ATP	PB-O3B-PG-O3G
2	A	501	ATP	C5'-O5'-PA-O1A
3	A	602	MLI	C2-C1-C3-O9
2	B	501	ATP	PA-O3A-PB-O3B
3	A	602	MLI	C2-C1-C3-O8

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	501	ATP	PG-O3B-PB-O1B
2	A	501	ATP	PG-O3B-PB-O2B
2	B	501	ATP	PB-O3B-PG-O1G
2	B	501	ATP	PG-O3B-PB-O2B
2	B	501	ATP	PG-O3B-PB-O1B


There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	601	MLI	1	0
3	A	602	MLI	2	0
2	A	501	ATP	1	0
2	B	501	ATP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	438/484 (90%)	0.88	84 (19%) 3 3	21, 41, 103, 140	9 (2%)
1	B	468/484 (96%)	0.76	71 (15%) 5 6	25, 43, 86, 130	16 (3%)
All	All	906/968 (93%)	0.82	155 (17%) 4 4	21, 42, 94, 140	25 (2%)

All (155) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	132	LEU	9.3
1	A	305	THR	9.1
1	A	71	ALA	8.4
1	A	306	LEU	7.8
1	B	305	THR	6.9
1	A	7[A]	LEU	6.7
1	A	304	ASP	6.5
1	A	300	ALA	6.5
1	B	300	ALA	6.2
1	B	306	LEU	6.1
1	B	412	ALA	6.1
1	B	-1	SER	6.0
1	A	57	ILE	5.9
1	A	301	HIS	5.9
1	A	255	ASP	5.8
1	B	2	VAL	5.7
1	A	102	LEU	5.7
1	B	-2	GLY	5.6
1	B	134	LEU	5.3
1	B	301	HIS	5.3
1	A	466	ASN	5.2
1	A	303	ARG	5.2
1	B	33	ALA	5.2
1	A	467	ASN	5.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	254	GLY	5.2
1	B	130[A]	ASP	5.2
1	A	407	LEU	5.2
1	A	405	PRO	5.1
1	B	411	ALA	5.1
1	B	3	LEU	5.1
1	B	136	VAL	5.1
1	B	0	MET	5.1
1	B	30	ILE	5.0
1	A	256	GLY	5.0
1	B	254	GLY	5.0
1	A	406	ASN	5.0
1	B	304	ASP	4.9
1	B	125	LEU	4.8
1	B	34	ILE	4.8
1	B	131	LYS	4.7
1	A	-2	GLY	4.6
1	B	151	GLU	4.6
1	B	7	LEU	4.4
1	B	122	ALA	4.4
1	B	302	GLY	4.4
1	A	63	GLY	4.3
1	A	302	GLY	4.3
1	B	133[A]	MET	4.3
1	A	68	ALA	4.2
1	A	298	ALA	4.2
1	A	69	LYS	4.1
1	A	120	LYS	4.1
1	A	67	LYS	4.1
1	A	0	MET	4.0
1	B	4	ASP	4.0
1	B	29	ASP	4.0
1	A	103	GLN	3.9
1	B	123	GLN	3.8
1	B	126	GLU	3.8
1	B	255	ASP	3.8
1	B	124	GLN	3.7
1	B	410	THR	3.7
1	A	101	VAL	3.7
1	B	32	ASP	3.7
1	A	299	GLY	3.7
1	A	408	ARG	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	256	GLY	3.6
1	A	258	LYS	3.6
1	B	129	ARG	3.6
1	A	-1	SER	3.5
1	B	303	ARG	3.5
1	B	152	THR	3.5
1	B	31	VAL	3.5
1	B	299	GLY	3.4
1	A	61	ALA	3.4
1	B	242	LEU	3.4
1	A	151	GLU	3.3
1	B	36	GLU	3.3
1	B	241	GLN	3.3
1	A	332[A]	ARG	3.3
1	A	58	CYS	3.3
1	B	415	VAL	3.3
1	A	413	GLN	3.3
1	A	241	GLN	3.2
1	B	150	GLU	3.2
1	B	413	GLN	3.1
1	A	50	ALA	3.1
1	A	56	ASN	3.1
1	B	414	ASN	3.1
1	A	54	LEU	3.1
1	A	105	LYS	3.1
1	A	415	VAL	3.0
1	B	257	ASP	3.0
1	A	242	LEU	3.0
1	B	1	MET	3.0
1	A	104	LEU	3.0
1	A	148	GLN	3.0
1	B	258	LYS	2.9
1	A	59	SER	2.9
1	B	135	ASN	2.9
1	A	66	LYS	2.9
1	A	46	PHE	2.9
1	B	332[A]	ARG	2.9
1	A	380	GLY	2.8
1	B	5	ILE	2.8
1	B	35	ILE	2.8
1	A	53	LYS	2.8
1	B	121	GLU	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	257	ASP	2.8
1	A	280	GLU	2.7
1	A	107	LEU	2.7
1	A	64	ALA	2.7
1	A	246	ASP	2.6
1	A	152	THR	2.6
1	B	443	GLU	2.6
1	B	149	ASP	2.6
1	B	119	ALA	2.6
1	A	3	LEU	2.6
1	A	127	GLU	2.5
1	B	127	GLU	2.5
1	A	125	LEU	2.5
1	A	100	CYS	2.5
1	A	149	ASP	2.5
1	B	8	PHE	2.5
1	B	11	GLU	2.5
1	A	62	VAL	2.5
1	A	134	LEU	2.4
1	A	4	ASP	2.4
1	B	298	ALA	2.4
1	A	70	GLU	2.4
1	A	2	VAL	2.4
1	A	49	GLU	2.3
1	B	280	GLU	2.3
1	A	65	LYS	2.3
1	B	128	GLU	2.3
1	A	123	GLN	2.3
1	A	1	MET	2.3
1	A	135	ASN	2.3
1	A	414	ASN	2.3
1	A	131	LYS	2.3
1	B	248	GLU	2.2
1	B	6	GLN	2.2
1	A	157	VAL	2.2
1	B	37	ALA	2.2
1	A	55	ILE	2.2
1	A	128	GLU	2.2
1	A	150	GLU	2.2
1	B	148	GLN	2.2
1	A	60	LYS	2.2
1	B	240	ALA	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	133[A]	MET	2.0
1	B	355	VAL	2.0
1	A	132	LEU	2.0
1	A	109	LYS	2.0
1	A	30	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MLI	A	601	7/7	0.70	0.24	57,62,68,70	0
3	MLI	A	602	7/7	0.83	0.14	48,64,70,73	0
2	ATP	A	501	31/31	0.92	0.11	35,44,62,67	8
2	ATP	B	501	31/31	0.93	0.10	33,44,75,79	5
4	NA	A	600	1/1	0.97	0.07	37,37,37,37	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ATP A 501:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ATP B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.