



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 06:46 AM UTC

PDB ID : 4LSX / pdb_00004lsx
Title : Plant steroid receptor ectodomain bound to brassinolide and SERK1 co-receptor ectodomain
Authors : Santiago, J.; Henzler, C.; Hothorn, M.
Deposited on : 2013-07-23
Resolution : 3.30 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1169 (3.32-3.28)
Clashscore	190562	1209 (3.32-3.28)
Ramachandran outliers	187476	1188 (3.32-3.28)
Sidechain outliers	187428	1187 (3.32-3.28)
RSRZ outliers	180081	1169 (3.32-3.28)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	774	71% (green), 24% (yellow), 5% (orange), 0% (red), 0% (grey)
1	B	774	68% (green), 23% (yellow), 8% (orange), 0% (red), 1% (grey)
2	C	203	65% (green), 27% (yellow), 9% (orange), 0% (red), 0% (grey)
2	D	203	67% (green), 24% (yellow), 9% (orange), 0% (red), 0% (grey)
3	E	4	50% (green), 25% (yellow), 25% (orange), 0% (red), 0% (grey)

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	H	4	 25% 50% 25%
3	J	4	 25% 50% 25%
4	F	2	 100%
4	K	2	 100%
5	G	5	 40% 40% 20%
6	I	3	 100%

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 13920 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein BRASSINOSTEROID INSENSITIVE 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	740	Total	C	N	O	S	0	0	0
			5454	3421	914	1088	31			
1	B	710	Total	C	N	O	S	0	0	0
			5217	3283	868	1037	29			

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	24	GLY	-	expression tag	UNP O22476
A	25	SER	-	expression tag	UNP O22476
A	26	SER	-	expression tag	UNP O22476
A	27	MET	-	expression tag	UNP O22476
A	28	GLY	-	expression tag	UNP O22476
A	643	GLU	GLY	engineered mutation	UNP O22476
A	789	LEU	-	expression tag	UNP O22476
A	790	GLU	-	expression tag	UNP O22476
A	791	ASN	-	expression tag	UNP O22476
A	792	LEU	-	expression tag	UNP O22476
A	793	TYR	-	expression tag	UNP O22476
A	794	PHE	-	expression tag	UNP O22476
A	795	GLN	-	expression tag	UNP O22476
A	796	GLY	-	expression tag	UNP O22476
A	797	ALA	-	expression tag	UNP O22476
B	24	GLY	-	expression tag	UNP O22476
B	25	SER	-	expression tag	UNP O22476
B	26	SER	-	expression tag	UNP O22476
B	27	MET	-	expression tag	UNP O22476
B	28	GLY	-	expression tag	UNP O22476
B	643	GLU	GLY	engineered mutation	UNP O22476
B	789	LEU	-	expression tag	UNP O22476
B	790	GLU	-	expression tag	UNP O22476
B	791	ASN	-	expression tag	UNP O22476
B	792	LEU	-	expression tag	UNP O22476

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	793	TYR	-	expression tag	UNP O22476
B	794	PHE	-	expression tag	UNP O22476
B	795	GLN	-	expression tag	UNP O22476
B	796	GLY	-	expression tag	UNP O22476
B	797	ALA	-	expression tag	UNP O22476

- Molecule 2 is a protein called Somatic embryogenesis receptor kinase 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
2	C	185	Total	C	N	O	S	0	0	0
			1397	881	238	273	5			
2	D	185	Total	C	N	O	S	0	0	0
			1380	871	234	271	4			

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	20	GLY	-	expression tag	UNP Q94AG2
C	21	SER	-	expression tag	UNP Q94AG2
C	22	SER	-	expression tag	UNP Q94AG2
C	23	MET	-	expression tag	UNP Q94AG2
C	214	LEU	-	expression tag	UNP Q94AG2
C	215	GLU	-	expression tag	UNP Q94AG2
C	216	ASN	-	expression tag	UNP Q94AG2
C	217	LEU	-	expression tag	UNP Q94AG2
C	218	TYR	-	expression tag	UNP Q94AG2
C	219	PHE	-	expression tag	UNP Q94AG2
C	220	GLN	-	expression tag	UNP Q94AG2
C	221	GLY	-	expression tag	UNP Q94AG2
C	222	ALA	-	expression tag	UNP Q94AG2
D	20	GLY	-	expression tag	UNP Q94AG2
D	21	SER	-	expression tag	UNP Q94AG2
D	22	SER	-	expression tag	UNP Q94AG2
D	23	MET	-	expression tag	UNP Q94AG2
D	214	LEU	-	expression tag	UNP Q94AG2
D	215	GLU	-	expression tag	UNP Q94AG2
D	216	ASN	-	expression tag	UNP Q94AG2
D	217	LEU	-	expression tag	UNP Q94AG2
D	218	TYR	-	expression tag	UNP Q94AG2
D	219	PHE	-	expression tag	UNP Q94AG2
D	220	GLN	-	expression tag	UNP Q94AG2
D	221	GLY	-	expression tag	UNP Q94AG2

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
D	222	ALA	-	expression tag	UNP Q94AG2

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	E	4	50	28	2	20	0	0	0
3	H	4	50	28	2	20	0	0	0
3	J	4	50	28	2	20	0	0	0

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	F	2	28	16	2	10	0	0	0
4	K	2	28	16	2	10	0	0	0

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-4)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
5	G	5	61	34	2	25	0	0	0

- Molecule 6 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
6	I	3	39	22	2	15	0	0	0

- Molecule 7 is Brassinolide (CCD ID: BLD) (formula: $C_{28}H_{48}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
7	A	1	34	28	6	0	0
7	B	1	34	28	6	0	0

- Molecule 8 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
8	A	1	Total 14	C 8	N 1	O 5	0	0
8	A	1	Total 14	C 8	N 1	O 5	0	0
8	B	1	Total 14	C 8	N 1	O 5	0	0
8	C	1	Total 14	C 8	N 1	O 5	0	0
8	C	1	Total 14	C 8	N 1	O 5	0	0
8	D	1	Total 14	C 8	N 1	O 5	0	0
8	D	1	Total 14	C 8	N 1	O 5	0	0


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Protein BRASSINOSTEROID INSENSITIVE 1




- Molecule 1: Protein BRASSINOSTEROID INSENSITIVE 1






- Molecule 2: Somatic embryogenesis receptor kinase 1

Chain C: 65% 27% 9%



- Molecule 2: Somatic embryogenesis receptor kinase 1

Chain D: 67% 24% 9%



- Molecule 3: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E: 50% 25% 25%



- Molecule 3: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H: 25% 50% 25%



- Molecule 3: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J: 25% 50% 25%



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  100%



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K:  100%




- Molecule 5: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-4)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  40% 40% 20%



- Molecule 6: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  100%



4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	69.90Å 69.90Å 873.55Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.53 – 3.30 48.53 – 3.30	Depositor EDS
% Data completeness (in resolution range)	98.2 (48.53-3.30) 96.4 (48.53-3.30)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.87 (at 3.33Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: dev_1334)	Depositor
R, R_{free}	0.246 , 0.285 0.241 , 0.281	Depositor DCC
R_{free} test set	2010 reflections (5.70%)	wwPDB-VP
Wilson B-factor (Å ²)	86.0	Xtrriage
Anisotropy	0.180	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 137.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.25$	Xtrriage
Estimated twinning fraction	0.458 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	13920	wwPDB-VP
Average B, all atoms (Å ²)	122.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, BLD, MAN, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/5553	0.86	10/7543 (0.1%)
1	B	0.29	0/5311	0.84	5/7218 (0.1%)
2	C	0.33	0/1427	0.86	1/1959 (0.1%)
2	D	0.32	0/1410	0.83	0/1940
All	All	0.30	0/13701	0.85	16/18660 (0.1%)

There are no bond length outliers.

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	332	VAL	CA-C-N	-6.29	113.91	120.38
1	A	332	VAL	C-N-CA	-6.29	113.91	120.38
1	B	759	ASN	CA-C-N	6.24	125.73	119.24
1	B	759	ASN	C-N-CA	6.24	125.73	119.24
1	A	332	VAL	N-CA-C	6.10	113.88	108.63

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5454	0	5251	115	0
1	B	5217	0	5032	110	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	1397	0	1366	33	0
2	D	1380	0	1332	32	0
3	E	50	0	43	1	0
3	H	50	0	43	2	0
3	J	50	0	43	2	0
4	F	28	0	25	2	0
4	K	28	0	25	0	0
5	G	61	0	52	1	0
6	I	39	0	34	0	0
7	A	34	0	47	5	0
7	B	34	0	47	5	0
8	A	28	0	26	3	0
8	B	14	0	13	1	0
8	C	28	0	26	0	0
8	D	28	0	26	0	0
All	All	13920	0	13431	292	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 292 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:551:ASN:HD21	8:A:814:NAG:H82	1.39	0.86
1:A:290:LEU:HB3	1:A:314:ALA:HB2	1.71	0.73
2:C:64:THR:HB	2:C:73:ARG:HB2	1.72	0.72
2:D:140:LEU:HD12	2:D:143:LEU:HD22	1.72	0.71
1:B:198:GLY:HA2	1:B:220:ARG:HH12	1.56	0.71

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	738/774 (95%)	710 (96%)	27 (4%)	1 (0%)	48	75
1	B	706/774 (91%)	680 (96%)	25 (4%)	1 (0%)	48	75
2	C	183/203 (90%)	176 (96%)	7 (4%)	0	100	100
2	D	183/203 (90%)	177 (97%)	6 (3%)	0	100	100
All	All	1810/1954 (93%)	1743 (96%)	65 (4%)	2 (0%)	48	75

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	637	ILE
1	B	637	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	611/680 (90%)	598 (98%)	13 (2%)	47	67
1	B	580/680 (85%)	569 (98%)	11 (2%)	50	68
2	C	165/182 (91%)	165 (100%)	0	100	100
2	D	161/182 (88%)	160 (99%)	1 (1%)	78	81
All	All	1517/1724 (88%)	1492 (98%)	25 (2%)	55	72

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	199	CYS
1	B	435	THR
2	D	78	ASN
1	B	410	LEU
1	B	439	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	127	ASN
2	D	118	ASN
2	C	78	ASN
2	D	94	ASN
1	B	705	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

24 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	E	1	3,1	14,14,15	0.25	0	17,19,21	0.48	0
3	NAG	E	2	3	14,14,15	0.47	0	17,19,21	1.34	2 (11%)
3	BMA	E	3	3	11,11,12	0.67	0	15,15,17	0.94	0
3	MAN	E	4	3	11,11,12	0.72	0	15,15,17	0.91	2 (13%)
4	NAG	F	1	4,1	14,14,15	0.35	0	17,19,21	0.52	0
4	NAG	F	2	4	14,14,15	0.41	0	17,19,21	0.64	1 (5%)
5	NAG	G	1	5,1	14,14,15	0.30	0	17,19,21	0.42	0
5	NAG	G	2	5	14,14,15	0.21	0	17,19,21	0.38	0
5	BMA	G	3	5	11,11,12	0.94	0	15,15,17	0.91	0
5	MAN	G	4	5	11,11,12	0.71	0	15,15,17	0.93	2 (13%)
5	MAN	G	5	5	11,11,12	0.75	0	15,15,17	1.52	3 (20%)
3	NAG	H	1	3,1	14,14,15	0.30	0	17,19,21	0.44	0
3	NAG	H	2	3	14,14,15	0.48	0	17,19,21	1.33	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	BMA	H	3	3	11,11,12	0.61	0	15,15,17	0.88	0
3	MAN	H	4	3	11,11,12	0.76	0	15,15,17	0.94	2 (13%)
6	NAG	I	1	6,1	14,14,15	0.61	1 (7%)	17,19,21	0.59	0
6	NAG	I	2	6	14,14,15	0.63	1 (7%)	17,19,21	0.61	0
6	BMA	I	3	6	11,11,12	0.64	0	15,15,17	0.96	1 (6%)
3	NAG	J	1	3,1	14,14,15	0.22	0	17,19,21	0.48	0
3	NAG	J	2	3	14,14,15	0.48	0	17,19,21	0.47	0
3	BMA	J	3	3	11,11,12	1.32	3 (27%)	15,15,17	1.44	2 (13%)
3	MAN	J	4	3	11,11,12	0.80	1 (9%)	15,15,17	0.99	2 (13%)
4	NAG	K	1	4,1	14,14,15	0.40	0	17,19,21	0.41	0
4	NAG	K	2	4	14,14,15	0.25	0	17,19,21	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	E	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	E	2	3	-	5/6/23/26	0/1/1/1
3	BMA	E	3	3	-	2/2/19/22	0/1/1/1
3	MAN	E	4	3	-	0/2/19/22	0/1/1/1
4	NAG	F	1	4,1	-	2/6/23/26	0/1/1/1
4	NAG	F	2	4	-	2/6/23/26	0/1/1/1
5	NAG	G	1	5,1	-	2/6/23/26	0/1/1/1
5	NAG	G	2	5	-	0/6/23/26	0/1/1/1
5	BMA	G	3	5	-	0/2/19/22	0/1/1/1
5	MAN	G	4	5	-	2/2/19/22	0/1/1/1
5	MAN	G	5	5	-	0/2/19/22	0/1/1/1
3	NAG	H	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	H	2	3	-	6/6/23/26	0/1/1/1
3	BMA	H	3	3	-	0/2/19/22	0/1/1/1
3	MAN	H	4	3	-	0/2/19/22	0/1/1/1
6	NAG	I	1	6,1	-	0/6/23/26	0/1/1/1
6	NAG	I	2	6	-	2/6/23/26	0/1/1/1
6	BMA	I	3	6	-	2/2/19/22	0/1/1/1
3	NAG	J	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	J	2	3	-	0/6/23/26	0/1/1/1
3	BMA	J	3	3	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MAN	J	4	3	-	0/2/19/22	0/1/1/1
4	NAG	K	1	4,1	-	1/6/23/26	0/1/1/1
4	NAG	K	2	4	-	1/6/23/26	0/1/1/1

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	J	3	BMA	O5-C1	-2.89	1.38	1.43
6	I	2	NAG	O5-C1	-2.22	1.40	1.43
3	J	3	BMA	C4-C3	2.19	1.58	1.52
6	I	1	NAG	O5-C1	-2.17	1.40	1.43
3	J	4	MAN	O5-C1	-2.15	1.40	1.43

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	2	NAG	C2-N2-C7	4.65	129.13	122.90
3	H	2	NAG	C2-N2-C7	4.64	129.12	122.90
5	G	5	MAN	C1-O5-C5	3.99	117.54	112.19
3	J	3	BMA	C2-C3-C4	3.88	117.69	110.86
3	J	3	BMA	C3-C4-C5	2.78	115.28	110.23

There are no chirality outliers.

5 of 27 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	F	1	NAG	C4-C5-C6-O6
4	F	1	NAG	O5-C5-C6-O6
4	F	2	NAG	O5-C5-C6-O6
3	H	2	NAG	O5-C5-C6-O6
5	G	4	MAN	O5-C5-C6-O6

There are no ring outliers.

8 monomers are involved in 8 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	J	2	NAG	1	0
5	G	5	MAN	1	0
5	G	3	BMA	1	0
3	E	2	NAG	1	0
4	F	1	NAG	2	0


Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	H	2	NAG	1	0
3	J	4	MAN	1	0
3	H	1	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.














5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
8	NAG	B	802	1	14,14,15	0.25	0	17,19,21	0.45	0
8	NAG	D	1001	2	14,14,15	0.39	0	17,19,21	0.45	0
8	NAG	A	814	1	14,14,15	0.59	0	17,19,21	1.21	1 (5%)
8	NAG	D	1002	2	14,14,15	1.81	1 (7%)	17,19,21	1.63	3 (17%)
8	NAG	C	1000	2	14,14,15	1.08	1 (7%)	17,19,21	0.63	0
7	BLD	A	801	-	36,37,37	3.09	12 (33%)	49,59,59	1.99	15 (30%)
8	NAG	C	1001	2	14,14,15	0.21	0	17,19,21	0.46	0
8	NAG	A	802	1	14,14,15	0.35	0	17,19,21	0.61	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	BLD	B	801	-	36,37,37	3.08	12 (33%)	49,59,59	1.89	15 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	NAG	B	802	1	-	2/6/23/26	0/1/1/1
8	NAG	D	1001	2	-	0/6/23/26	0/1/1/1
8	NAG	A	814	1	-	0/6/23/26	0/1/1/1
8	NAG	D	1002	2	-	2/6/23/26	0/1/1/1
8	NAG	C	1000	2	-	2/6/23/26	0/1/1/1
7	BLD	A	801	-	-	4/20/85/85	0/4/4/4
8	NAG	C	1001	2	-	2/6/23/26	0/1/1/1
8	NAG	A	802	1	-	2/6/23/26	0/1/1/1
7	BLD	B	801	-	-	4/20/85/85	0/4/4/4

The worst 5 of 26 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	801	BLD	O07-C07	-10.51	1.30	1.45
7	B	801	BLD	O07-C07	-10.35	1.30	1.45
7	B	801	BLD	O06-C06	9.91	1.46	1.21
7	A	801	BLD	O06-C06	9.82	1.46	1.21
8	D	1002	NAG	C1-C2	6.35	1.61	1.52

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	801	BLD	C15-C14-C08	-5.47	110.37	119.10
7	A	801	BLD	O07-C06-O06	-5.46	108.67	116.97
7	B	801	BLD	C15-C14-C08	-5.24	110.74	119.10
7	B	801	BLD	O07-C06-O06	-5.20	109.07	116.97
7	B	801	BLD	C01-C10-C05	5.18	115.22	107.11

There are no chirality outliers.

5 of 18 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
7	A	801	BLD	C20-C22-C23-O23
7	A	801	BLD	C20-C22-C23-C24
7	A	801	BLD	O22-C22-C23-C24
7	B	801	BLD	C20-C22-C23-O23
7	B	801	BLD	O22-C22-C23-C24

There are no ring outliers.

5 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	B	802	NAG	1	0
8	A	814	NAG	2	0
7	A	801	BLD	5	0
8	A	802	NAG	1	0
7	B	801	BLD	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	740/774 (95%)	-1.00	0	100 100	47, 114, 169, 224	0
1	B	710/774 (91%)	-0.94	0	100 100	57, 115, 166, 222	0
2	C	185/203 (91%)	-0.66	0	100 100	88, 145, 192, 210	0
2	D	185/203 (91%)	-0.65	1 (0%)	87 76	105, 146, 225, 258	0
All	All	1820/1954 (93%)	-0.91	1 (0%)	92 90	47, 121, 183, 258	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	204	PRO	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	BMA	H	3	11/12	0.90	0.08	125,136,149,156	0
3	MAN	H	4	11/12	0.90	0.07	150,157,165,169	0
3	MAN	E	4	11/12	0.95	0.06	83,104,121,128	0
4	NAG	F	2	14/15	0.95	0.07	107,112,122,127	0
4	NAG	K	2	14/15	0.96	0.04	72,85,107,107	0
3	BMA	J	3	11/12	0.97	0.06	98,131,145,150	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	BMA	E	3	11/12	0.97	0.06	77,110,116,116	0
3	NAG	H	2	14/15	0.97	0.07	109,116,136,136	0
5	BMA	G	3	11/12	0.97	0.04	101,118,146,149	0
5	MAN	G	4	11/12	0.97	0.05	128,135,139,143	0
6	NAG	I	2	14/15	0.97	0.05	93,114,124,132	0
6	BMA	I	3	11/12	0.97	0.05	136,138,139,143	0
3	NAG	J	2	14/15	0.98	0.06	98,120,128,130	0
3	NAG	E	2	14/15	0.98	0.05	108,114,118,120	0
3	MAN	J	4	11/12	0.98	0.03	84,92,96,99	0
5	MAN	G	5	11/12	0.98	0.03	76,87,117,130	0
3	NAG	J	1	14/15	0.98	0.04	86,99,107,109	0
4	NAG	K	1	14/15	0.98	0.07	93,107,121,122	0
3	NAG	H	1	14/15	0.99	0.06	110,124,137,137	0
3	NAG	E	1	14/15	0.99	0.04	48,61,73,95	0
4	NAG	F	1	14/15	0.99	0.04	92,102,111,111	0
6	NAG	I	1	14/15	0.99	0.06	72,85,96,100	0
5	NAG	G	1	14/15	0.99	0.05	111,121,148,149	0
5	NAG	G	2	14/15	0.99	0.05	88,102,106,111	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain H:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain J:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain F:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain K:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain G:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain I:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
8	NAG	A	814	14/15	0.97	0.05	88,94,101,103	0
8	NAG	D	1001	14/15	0.97	0.06	102,115,132,134	0
8	NAG	B	802	14/15	0.98	0.04	96,105,113,114	0
8	NAG	C	1000	14/15	0.98	0.03	96,99,105,106	0
8	NAG	C	1001	14/15	0.98	0.05	77,91,108,109	0
8	NAG	A	802	14/15	0.98	0.07	138,158,174,175	0
8	NAG	D	1002	14/15	0.98	0.03	81,98,108,109	0
7	BLD	B	801	34/34	0.99	0.08	85,127,138,149	0
7	BLD	A	801	34/34	0.99	0.07	61,109,134,140	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.