



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:13 PM UTC

PDB ID : 1LT4 / pdb_00001lt4
Title : HEAT-LABILE ENTEROTOXIN MUTANT S63K
Authors : Van Den Akker, F.; Hol, W.G.J.
Deposited on : 1997-04-14
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	D	103	87%12%.
1	E	103	88%10%.
1	F	103	%83%16%.
1	G	103	83%16%.
1	H	103	80%18%..

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	A	247	% 66%23%9%
3	B	2	100%
3	C	2	100%
3	I	2	100%
3	J	2	100%
3	K	2	100%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6452 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HEAT-LABILE ENTEROTOXIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	D	103	Total	C	N	O	S	0	0	0
			824	516	139	163	6			
1	E	103	Total	C	N	O	S	0	0	0
			824	516	139	163	6			
1	F	103	Total	C	N	O	S	0	0	0
			824	516	139	163	6			
1	G	103	Total	C	N	O	S	0	0	0
			824	516	139	163	6			
1	H	103	Total	C	N	O	S	0	0	0
			824	516	139	163	6			

- Molecule 2 is a protein called HEAT-LABILE ENTEROTOXIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	226	Total	C	N	O	S	0	0	0
			1861	1170	336	350	5			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	63	LYS	SER	engineered mutation	UNP P06717
A	188D	ASN	-	insertion	UNP P06717
A	188E	SER	-	insertion	UNP P06717
A	188F	SER	-	insertion	UNP P06717
A	188G	ARG	-	insertion	UNP P06717
A	188H	THR	-	insertion	UNP P06717
A	188I	ILE	-	insertion	UNP P06717
A	188J	THR	-	insertion	UNP P06717

- Molecule 3 is an oligosaccharide called beta-D-galactopyranose-(1-4)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	B	2	Total	C	O	0	0	0
			23	12	11			
3	C	2	Total	C	O	0	0	0
			23	12	11			
3	I	2	Total	C	O	0	0	0
			23	12	11			
3	J	2	Total	C	O	0	0	0
			23	12	11			
3	K	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	58	Total	O	0	0
			58	58		
4	E	54	Total	O	0	0
			54	54		
4	F	62	Total	O	0	0
			62	62		
4	G	61	Total	O	0	0
			61	61		
4	H	53	Total	O	0	0
			53	53		
4	A	68	Total	O	0	0
			68	68		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: HEAT-LABILE ENTEROTOXIN

Chain D:  87% 12% .



- Molecule 1: HEAT-LABILE ENTEROTOXIN

Chain E:  88% 10% .



- Molecule 1: HEAT-LABILE ENTEROTOXIN

Chain F:  83% 16% .



- Molecule 1: HEAT-LABILE ENTEROTOXIN

Chain G:  83% 16% .




- Molecule 1: HEAT-LABILE ENTEROTOXIN

Chain H:  80% 18% ..



- Molecule 2: HEAT-LABILE ENTEROTOXIN

Chain A:  66% 23% 9% .



- Molecule 3: beta-D-galactopyranose-(1-4)-beta-D-glucopyranose

Chain B: 100%

BGC1
GAL2

- Molecule 3: beta-D-galactopyranose-(1-4)-beta-D-glucopyranose

Chain C: 100%

BGC1
GAL2

- Molecule 3: beta-D-galactopyranose-(1-4)-beta-D-glucopyranose

Chain I: 100%

BGC1
GAL2

- Molecule 3: beta-D-galactopyranose-(1-4)-beta-D-glucopyranose

Chain J: 100%

BGC1
GAL2

- Molecule 3: beta-D-galactopyranose-(1-4)-beta-D-glucopyranose

Chain K: 100%

BGC1
GAL2

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	120.00Å 101.90Å 64.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.00 10.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	92.9 (10.00-2.00) 92.5 (10.00-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.19 (at 2.00Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.192 , 0.255 0.202 , 0.263	Depositor DCC
R_{free} test set	2553 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	22.4	Xtriage
Anisotropy	0.438	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 71.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6452	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, GAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	D	0.67	0/835	0.97	3/1124 (0.3%)
1	E	0.68	0/835	0.97	3/1124 (0.3%)
1	F	0.68	0/835	1.00	4/1124 (0.4%)
1	G	0.63	0/835	0.97	5/1124 (0.4%)
1	H	0.69	0/835	1.01	5/1124 (0.4%)
2	A	0.60	0/1913	1.03	13/2595 (0.5%)
All	All	0.65	0/6088	1.00	33/8215 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	1

There are no bond length outliers.

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	50	VAL	N-CA-C	-10.07	93.19	107.80
1	F	50	VAL	N-CA-C	-9.78	93.62	107.80
1	G	50	VAL	N-CA-C	-8.87	94.94	107.80
1	D	50	VAL	N-CA-C	-8.62	95.30	107.80
1	H	50	VAL	N-CA-C	-8.28	95.79	107.80
1	H	52	VAL	N-CA-C	-8.23	101.37	109.02
2	A	107	HIS	CA-C-N	7.66	127.55	119.05
2	A	107	HIS	C-N-CA	7.66	127.55	119.05
2	A	12	PRO	CA-C-N	7.51	129.23	119.84
2	A	12	PRO	C-N-CA	7.51	129.23	119.84

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	51	GLU	N-CA-C	7.46	120.50	110.35
1	G	52	VAL	N-CA-C	-7.38	100.91	108.96
2	A	222	ILE	N-CA-C	6.85	119.65	111.09
1	E	52	VAL	N-CA-C	-6.66	102.43	108.95
1	D	52	VAL	N-CA-C	-6.43	103.04	109.02
2	A	11	ARG	CA-C-N	6.12	126.68	120.38
2	A	11	ARG	C-N-CA	6.12	126.68	120.38
2	A	53	VAL	N-CA-C	-5.95	101.11	109.21
1	F	51	GLU	N-CA-C	5.87	118.48	110.55
1	E	51	GLU	N-CA-C	5.84	118.08	110.43
1	H	102	LYS	N-CA-C	5.64	118.65	109.06
1	F	96	ILE	N-CA-C	5.61	116.24	108.84
2	A	88	ILE	N-CA-C	5.48	116.38	108.48
1	H	56	GLN	N-CA-C	-5.45	106.61	113.20
2	A	176	GLU	N-CA-C	5.45	117.47	109.24
1	H	51	GLU	N-CA-C	5.38	117.48	110.43
1	F	86	CYS	N-CA-C	-5.37	99.78	108.52
2	A	76	ILE	N-CA-C	5.31	117.21	111.58
2	A	93	ASN	N-CA-C	-5.29	106.87	113.38
2	A	183	ALA	N-CA-C	5.26	117.06	109.48
1	G	102	LYS	N-CA-C	5.16	117.23	109.23
1	G	86	CYS	N-CA-C	-5.13	100.15	108.52
1	D	86	CYS	N-CA-C	-5.13	100.37	108.73

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	76	TYR	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	824	0	841	6	0
1	E	824	0	841	8	0
1	F	824	0	841	10	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	824	0	841	7	0
1	H	824	0	841	10	0
2	A	1861	0	1742	35	0
3	B	23	0	21	0	0
3	C	23	0	21	0	0
3	I	23	0	21	0	0
3	J	23	0	21	0	0
3	K	23	0	21	0	0
4	A	68	0	0	1	0
4	D	58	0	0	0	0
4	E	54	0	0	3	0
4	F	62	0	0	0	0
4	G	61	0	0	1	0
4	H	53	0	0	0	0
All	All	6452	0	6052	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:93:PRO:HG3	1:H:3:GLN:HG2	1.57	0.86
1:D:3:GLN:HG2	1:E:93:PRO:HG3	1.63	0.81
1:F:82:ILE:HD12	1:F:99:ILE:HD11	1.65	0.79
1:E:66:GLU:HG3	2:A:236:ILE:HD11	1.65	0.78
1:H:59:ASP:HA	1:H:62:LYS:HD3	1.67	0.77
1:H:63:LYS:HA	1:H:63:LYS:HE2	1.69	0.73
2:A:37:MET:HE1	2:A:117:GLY:HA2	1.71	0.73
2:A:84:TYR:CE2	2:A:129:ARG:HG2	2.24	0.72
1:H:75:THR:HG23	1:H:80:THR:HB	1.70	0.71
2:A:5:LEU:HD12	2:A:94:MET:HE3	1.71	0.70
2:A:168:PRO:HG2	2:A:171:HIS:HB2	1.75	0.69
2:A:186:GLY:O	2:A:199:CYS:HB2	1.94	0.67
2:A:106:PRO:O	2:A:108:PRO:HD3	1.96	0.65
2:A:94:MET:SD	2:A:115:ALA:HB2	2.38	0.63
1:E:37:MET:HE2	1:E:52:VAL:HG23	1.80	0.63
1:F:71:THR:O	1:F:75:THR:HG23	2.03	0.58
2:A:147:ASP:O	2:A:151:ARG:HB2	2.02	0.58
1:E:66:GLU:CG	2:A:236:ILE:HD11	2.33	0.57
2:A:122:SER:HB2	2:A:146:ARG:HG2	1.86	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:174:TRP:HB2	2:A:175:ARG:HH21	1.70	0.56
1:G:37:MET:HE2	1:G:39:ILE:HD11	1.88	0.56
2:A:208:THR:O	2:A:212:ARG:HG2	2.05	0.56
2:A:84:TYR:HE2	2:A:129:ARG:HG2	1.70	0.54
1:F:103:ASN:HB2	2:A:151:ARG:HH12	1.73	0.54
2:A:174:TRP:HB3	2:A:188:GLY:HA3	1.91	0.53
1:H:20:ILE:HG21	1:H:42:PHE:CE2	2.43	0.53
2:A:33:ARG:HD2	2:A:212:ARG:CZ	2.39	0.53
1:F:59:ASP:HA	1:F:62:LYS:HD3	1.90	0.52
2:A:106:PRO:C	2:A:108:PRO:HD3	2.34	0.52
1:F:61:GLN:O	1:F:65:ILE:HG13	2.10	0.51
2:A:46:ARG:HG3	2:A:106:PRO:HG2	1.91	0.51
1:F:58:ILE:O	1:F:61:GLN:HB2	2.11	0.50
2:A:174:TRP:HB2	2:A:175:ARG:NH2	2.26	0.50
1:H:54:GLY:HA3	1:H:56:GLN:NE2	2.26	0.50
1:G:33:GLY:O	1:G:34:LYS:HB2	2.12	0.49
2:A:30:TYR:CZ	2:A:120:PRO:HG3	2.48	0.49
1:F:103:ASN:ND2	1:G:76:TYR:CE1	2.80	0.49
1:E:81:LYS:HG3	4:E:136:HOH:O	2.13	0.48
1:D:44:SER:OG	1:D:46:GLU:HG2	2.14	0.48
1:E:35:ARG:NH1	4:E:130:HOH:O	2.48	0.46
1:H:51:GLU:HG3	1:H:95:SER:OG	2.16	0.46
2:A:184:PRO:HB2	2:A:187:CYS:SG	2.56	0.46
2:A:198:THR:HA	2:A:201:GLU:HG2	1.98	0.46
2:A:44:HIS:HB2	2:A:57:ASP:HB2	1.97	0.45
2:A:74:GLN:O	2:A:78:SER:HB3	2.17	0.44
2:A:82:THR:HA	2:A:130:VAL:O	2.17	0.44
2:A:149:TYR:HB2	4:A:293:HOH:O	2.17	0.44
1:D:8:LEU:C	1:D:8:LEU:HD23	2.43	0.44
2:A:172:GLN:OE1	2:A:172:GLN:N	2.51	0.44
1:D:46:GLU:HG2	1:D:46:GLU:H	1.55	0.43
1:H:13:ARG:O	1:H:14:ASN:HB2	2.18	0.43
1:E:3:GLN:NE2	1:F:92:THR:HG22	2.34	0.42
1:G:20:ILE:HG21	1:G:42:PHE:CE1	2.54	0.42
1:G:77:LEU:HD12	4:G:146:HOH:O	2.18	0.42
1:F:74:ILE:HD11	2:A:224:SER:HB3	2.01	0.42
2:A:174:TRP:HA	2:A:174:TRP:CE3	2.55	0.42
1:D:20:ILE:HG21	1:D:42:PHE:CE1	2.55	0.42
2:A:21:GLY:HA3	2:A:119:ILE:O	2.19	0.42
2:A:12:PRO:HA	2:A:13:PRO:HD3	1.86	0.41
2:A:42:TYR:HD1	2:A:101:LEU:HD13	1.85	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:177:GLU:HA	2:A:178:PRO:HA	1.81	0.41
2:A:33:ARG:HH11	2:A:33:ARG:HG3	1.85	0.41
1:H:24:ILE:HD11	1:H:82:ILE:HD11	2.02	0.41
1:G:102:LYS:O	1:G:103:ASN:HB2	2.21	0.41
1:H:58:ILE:HG22	1:H:61:GLN:OE1	2.20	0.41
1:G:61:GLN:O	1:G:65:ILE:HG13	2.21	0.41
4:E:135:HOH:O	1:F:34:LYS:HA	2.21	0.41
1:E:2:PRO:HG3	1:E:11:GLU:OE2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	101/103 (98%)	100 (99%)	1 (1%)	0	100	100
1	E	101/103 (98%)	100 (99%)	1 (1%)	0	100	100
1	F	101/103 (98%)	100 (99%)	1 (1%)	0	100	100
1	G	101/103 (98%)	99 (98%)	2 (2%)	0	100	100
1	H	101/103 (98%)	99 (98%)	2 (2%)	0	100	100
2	A	222/247 (90%)	211 (95%)	9 (4%)	2 (1%)	14	9
All	All	727/762 (95%)	709 (98%)	16 (2%)	2 (0%)	36	35

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	137	GLU
2	A	172	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	95/95 (100%)	91 (96%)	4 (4%)	26	25
1	E	95/95 (100%)	94 (99%)	1 (1%)	65	73
1	F	95/95 (100%)	92 (97%)	3 (3%)	34	35
1	G	95/95 (100%)	94 (99%)	1 (1%)	65	73
1	H	95/95 (100%)	94 (99%)	1 (1%)	65	73
2	A	195/215 (91%)	187 (96%)	8 (4%)	27	26
All	All	670/690 (97%)	652 (97%)	18 (3%)	39	42

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	13	ARG
1	D	46	GLU
1	D	91	LYS
1	D	103	ASN
1	E	3	GLN
1	F	34	LYS
1	F	102	LYS
1	F	103	ASN
1	G	59	ASP
1	H	56	GLN
2	A	37	MET
2	A	55	TYR
2	A	137	GLU
2	A	143	ARG
2	A	154	ASN
2	A	175	ARG
2	A	201	GLU
2	A	235	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	16	GLN
1	D	21	ASN
1	D	94	ASN
1	D	103	ASN
1	E	94	ASN
1	F	94	ASN
1	F	103	ASN
1	G	21	ASN
1	G	49	GLN
1	G	103	ASN
1	H	56	GLN
1	H	94	ASN
1	H	103	ASN
2	A	107	HIS
2	A	152	ASN
2	A	200	ASN
2	A	227	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

10 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	BGC	B	1	3	12,12,12	0.53	0	17,17,17	0.91	1 (5%)
3	GAL	B	2	3	11,11,12	1.24	1 (9%)	15,15,17	1.30	2 (13%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	BGC	C	1	3	12,12,12	1.26	1 (8%)	17,17,17	1.25	2 (11%)
3	GAL	C	2	3	11,11,12	1.20	1 (9%)	15,15,17	1.27	2 (13%)
3	BGC	I	1	3	12,12,12	2.21	3 (25%)	17,17,17	1.70	6 (35%)
3	GAL	I	2	3	11,11,12	0.92	1 (9%)	15,15,17	1.45	4 (26%)
3	BGC	J	1	3	12,12,12	1.38	1 (8%)	17,17,17	0.88	0
3	GAL	J	2	3	11,11,12	1.64	3 (27%)	15,15,17	1.40	2 (13%)
3	BGC	K	1	3	12,12,12	1.32	1 (8%)	17,17,17	1.52	5 (29%)
3	GAL	K	2	3	11,11,12	1.66	3 (27%)	15,15,17	1.95	7 (46%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BGC	B	1	3	-	0/2/22/22	0/1/1/1
3	GAL	B	2	3	-	0/2/19/22	0/1/1/1
3	BGC	C	1	3	-	0/2/22/22	0/1/1/1
3	GAL	C	2	3	-	0/2/19/22	0/1/1/1
3	BGC	I	1	3	-	1/2/22/22	0/1/1/1
3	GAL	I	2	3	-	0/2/19/22	0/1/1/1
3	BGC	J	1	3	-	0/2/22/22	0/1/1/1
3	GAL	J	2	3	-	0/2/19/22	0/1/1/1
3	BGC	K	1	3	-	2/2/22/22	0/1/1/1
3	GAL	K	2	3	-	0/2/19/22	0/1/1/1

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	I	1	BGC	C3-C2	5.09	1.65	1.52
3	I	1	BGC	C1-C2	3.72	1.60	1.52
3	C	1	BGC	C4-C5	3.64	1.60	1.53
3	K	1	BGC	C4-C3	3.60	1.61	1.52
3	K	2	GAL	C2-C3	3.42	1.57	1.52
3	J	1	BGC	C4-C5	3.22	1.59	1.53
3	K	2	GAL	C4-C3	3.01	1.60	1.52
3	I	1	BGC	C4-C5	2.98	1.59	1.53
3	B	2	GAL	C4-C5	2.85	1.59	1.53
3	J	2	GAL	C4-C5	2.73	1.58	1.53
3	J	2	GAL	C2-C3	2.51	1.56	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	2	GAL	C2-C3	-2.49	1.48	1.52
3	K	2	GAL	C4-C5	2.48	1.58	1.53
3	I	2	GAL	O2-C2	-2.35	1.38	1.43
3	J	2	GAL	O2-C2	-2.07	1.39	1.43

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	J	2	GAL	C1-O5-C5	3.98	117.52	112.19
3	C	1	BGC	O4-C4-C3	-3.81	101.39	110.38
3	K	2	GAL	C1-O5-C5	3.49	116.86	112.19
3	K	2	GAL	C6-C5-C4	3.29	121.09	113.02
3	I	1	BGC	C6-C5-C4	3.17	120.81	113.02
3	I	1	BGC	O3-C3-C2	3.15	117.81	110.38
3	I	2	GAL	C1-O5-C5	3.02	116.23	112.19
3	B	2	GAL	C6-C5-C4	2.95	120.25	113.02
3	K	2	GAL	O2-C2-C1	-2.78	102.85	109.22
3	C	2	GAL	O2-C2-C1	2.57	115.12	109.22
3	K	1	BGC	O3-C3-C2	-2.55	104.37	110.38
3	I	1	BGC	C3-C4-C5	-2.52	105.66	110.23
3	K	2	GAL	C2-C3-C4	-2.49	106.48	110.86
3	K	1	BGC	C6-C5-C4	2.49	119.13	113.02
3	J	2	GAL	C2-C3-C4	-2.49	106.48	110.86
3	I	1	BGC	O4-C4-C3	2.48	116.22	110.38
3	I	2	GAL	O5-C1-C2	-2.36	105.15	110.79
3	K	1	BGC	O5-C5-C4	-2.34	105.48	109.70
3	I	2	GAL	O3-C3-C2	-2.34	105.28	110.05
3	K	2	GAL	O3-C3-C2	2.28	114.70	110.05
3	K	1	BGC	O3-C3-C4	2.24	115.67	110.38
3	I	1	BGC	O5-C1-C2	2.16	114.10	110.30
3	K	1	BGC	O4-C4-C5	-2.16	104.01	109.32
3	B	1	BGC	C3-C4-C5	2.14	114.11	110.23
3	C	1	BGC	C6-C5-C4	2.12	118.23	113.02
3	K	2	GAL	O2-C2-C3	2.10	114.51	110.15
3	K	2	GAL	O5-C5-C4	-2.09	105.74	110.83
3	I	2	GAL	O4-C4-C3	-2.07	105.50	110.38
3	I	1	BGC	C4-C3-C2	-2.05	107.23	110.83
3	C	2	GAL	C1-O5-C5	2.03	114.91	112.19
3	B	2	GAL	C1-O5-C5	2.01	114.88	112.19

There are no chirality outliers.


All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	K	1	BGC	O5-C5-C6-O6
3	K	1	BGC	C4-C5-C6-O6
3	I	1	BGC	C4-C5-C6-O6


There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	D	103/103 (100%)	-0.90	0 100 100	7, 16, 34, 44	0
1	E	103/103 (100%)	-0.85	0 100 100	8, 17, 35, 49	0
1	F	103/103 (100%)	-0.79	1 (0%) 79 79	8, 17, 33, 49	0
1	G	103/103 (100%)	-0.85	0 100 100	8, 17, 34, 55	0
1	H	103/103 (100%)	-0.77	0 100 100	8, 19, 38, 57	0
2	A	226/247 (91%)	-0.03	3 (1%) 75 74	12, 33, 59, 73	0
All	All	741/762 (97%)	-0.59	4 (0%) 87 87	7, 21, 51, 73	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	47	GLY	3.8
2	A	197	ASP	2.7
2	A	236	ILE	2.2
1	F	103	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	BGC	B	1	12/12	-	-	37,54,59,63	0
3	GAL	B	2	11/12	-	-	21,23,30,30	0
3	BGC	C	1	12/12	-	-	32,53,58,61	0
3	GAL	C	2	11/12	-	-	17,20,24,27	0
3	BGC	I	1	12/12	0.67	0.14	47,58,62,65	0
3	BGC	K	1	12/12	0.77	0.09	48,59,63,63	0
3	BGC	J	1	12/12	0.87	0.07	26,37,40,43	0
3	GAL	I	2	11/12	0.88	0.08	26,34,40,42	0
3	GAL	K	2	11/12	0.88	0.08	26,32,41,42	0
3	GAL	J	2	11/12	0.95	0.05	14,19,25,29	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain I:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain J:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.