



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 11:44 PM UTC

PDB ID : 5LTN / pdb_00005ltn
Title : Crystal structure of Lymphocytic choriomeningitis mammarenavirus endonuclease complexed with DPBA
Authors : Saez-Ayala, M.; Yekwa, E.L.; Canard, B.; Alvarez, K.; Ferron, F.
Deposited on : 2016-09-07
Resolution : 1.88 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1220 (1.88-1.88)
Clashscore	190562	1234 (1.88-1.88)
Ramachandran outliers	187476	1222 (1.88-1.88)
Sidechain outliers	187428	1222 (1.88-1.88)
RSRZ outliers	180081	1220 (1.88-1.88)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	204	10% 79% 10% 10%
1	B	204	11% 82% 12% 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	B	204	-	-	X	-

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 3353 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA-directed RNA polymerase L.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	183	Total	C	N	O	S	0	3	0
			1492	946	246	290	10			
1	B	194	Total	C	N	O	S	6	1	0
			1572	998	258	306	10			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	initiating methionine	UNP A0A059U381
A	-6	LYS	-	expression tag	UNP A0A059U381
A	-5	HIS	-	expression tag	UNP A0A059U381
A	-4	HIS	-	expression tag	UNP A0A059U381
A	-3	HIS	-	expression tag	UNP A0A059U381
A	-2	HIS	-	expression tag	UNP A0A059U381
A	-1	HIS	-	expression tag	UNP A0A059U381
A	0	HIS	-	expression tag	UNP A0A059U381
A	173	ASP	ASN	conflict	UNP A0A059U381
B	-7	MET	-	initiating methionine	UNP A0A059U381
B	-6	LYS	-	expression tag	UNP A0A059U381
B	-5	HIS	-	expression tag	UNP A0A059U381
B	-4	HIS	-	expression tag	UNP A0A059U381
B	-3	HIS	-	expression tag	UNP A0A059U381
B	-2	HIS	-	expression tag	UNP A0A059U381
B	-1	HIS	-	expression tag	UNP A0A059U381
B	0	HIS	-	expression tag	UNP A0A059U381
B	173	ASP	ASN	conflict	UNP A0A059U381

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mg	0	0
			2	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Mg	0	0
			2	2		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is 2-4-DIOXO-4-PHENYLBUTANOIC ACID (CCD ID: XI7) (formula: $C_{10}H_8O_4$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	1	0
			14	10	4		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Cl	0	0
			1	1		
5	B	1	Total	Cl	0	0
			1	1		

- Molecule 6 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			7	4	3		
6	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 7 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	2	Total	Na	0	0
			2	2		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	118	Total	O	0	0
			118	118		
8	B	111	Total	O	0	0
			111	111		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: RNA-directed RNA polymerase L



• Molecule 1: RNA-directed RNA polymerase L



4 Data and refinement statistics

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, α , β , γ	107.19Å 107.19Å 53.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	53.59 – 1.88 53.59 – 1.88	Depositor EDS
% Data completeness (in resolution range)	99.7 (53.59-1.88) 99.7 (53.59-1.88)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.60 (at 1.88Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.190 , 0.216 0.193 , 0.219	Depositor DCC
R_{free} test set	2457 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	37.5	Xtriage
Anisotropy	0.198	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.029 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3353	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, XI7, PEG, NA, MG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/1526	0.62	0/2059
1	B	0.44	0/1601	0.64	0/2160
All	All	0.43	0/3127	0.63	0/4219

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	180	SER	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1492	0	1489	19	0
1	B	1572	0	1571	21	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	12	0	16	1	0
3	B	12	0	16	5	4
4	A	14	0	7	3	0
5	A	1	0	0	1	0
5	B	1	0	0	0	0
6	A	14	0	20	2	0
7	A	2	0	0	0	0
8	A	118	0	0	12	6
8	B	111	0	0	4	2
All	All	3353	0	3119	44	6

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:153:GLU:OE1	3:B:204:GOL:C3	2.22	0.87
1:A:102:CYS:O	8:A:301:HOH:O	1.95	0.83
1:A:88:ASP:OD1	8:A:301:HOH:O	1.97	0.83
1:A:181:GLU:OE1	8:A:302:HOH:O	1.97	0.81
1:B:153:GLU:OE1	3:B:204:GOL:H31	1.82	0.79
1:A:126[A]:ARG:NH2	5:A:206:CL:CL	2.53	0.79
1:B:153:GLU:OE1	3:B:204:GOL:H32	1.83	0.78
1:A:0:HIS:N	8:A:306:HOH:O	2.21	0.73
4:A:205:XI7:O11	8:A:301:HOH:O	2.06	0.72
4:A:205:XI7:O8	8:A:303:HOH:O	2.06	0.72
1:B:56:LYS:NZ	8:B:305:HOH:O	2.25	0.68
1:A:63:HIS:CD2	3:A:204:GOL:H11	2.33	0.64
1:B:101:GLU:OE2	8:B:301:HOH:O	2.17	0.57
1:A:43:LYS:NZ	8:A:313:HOH:O	2.38	0.55
1:A:178:GLU:HA	1:A:181:GLU:CD	2.36	0.51
1:B:188:SER:O	1:B:190:LYS:N	2.44	0.51
1:A:178:GLU:HA	1:A:181:GLU:OE2	2.13	0.49
6:A:207:PEG:H22	8:A:402:HOH:O	2.13	0.48
1:A:147:ASP:HB2	8:A:403:HOH:O	2.14	0.47
1:B:151:MET:HE2	1:B:151:MET:HB3	1.76	0.47
1:B:22:SER:OG	8:B:302:HOH:O	2.20	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:188:SER:C	1:B:190:LYS:N	2.72	0.47
1:B:67:SER:HA	1:B:86:ILE:HD13	1.96	0.46
1:B:153:GLU:OE2	3:B:204:GOL:H11	2.15	0.46
1:B:171:ASN:O	1:B:175:LYS:HG2	2.15	0.46
1:B:187:GLU:O	1:B:190:LYS:HE3	2.16	0.46
1:A:7:LEU:HB3	1:A:28:PHE:HZ	1.80	0.46
1:A:127:GLU:HG3	8:A:371:HOH:O	2.13	0.46
1:B:69:GLU:O	1:B:73:VAL:HG13	2.16	0.45
1:B:83:PRO:HD2	1:B:125:ILE:HG22	1.98	0.44
1:A:179:GLU:N	1:A:179:GLU:OE1	2.51	0.43
1:B:79:CYS:HB3	1:B:82:LEU:HD21	2.00	0.43
1:A:50:GLU:OE2	4:A:205:X17:O10	2.35	0.43
1:B:153:GLU:OE2	3:B:204:GOL:C1	2.66	0.43
1:B:191:THR:O	8:B:303:HOH:O	2.21	0.42
1:A:10:LEU:HD11	1:A:158:LYS:HD3	2.00	0.42
1:A:181:GLU:O	1:A:182:TYR:HB2	2.20	0.41
1:B:47:ARG:HE	1:B:47:ARG:HB2	1.60	0.41
1:A:174:GLU:O	1:A:178:GLU:N	2.48	0.41
1:B:43:LYS:HD2	1:B:43:LYS:HA	1.72	0.41
1:A:95:ASN:HA	6:A:208:PEG:H22	2.03	0.41
1:B:112:GLU:O	1:B:116:ILE:HG12	2.21	0.41
1:A:18:ASP:OD2	8:A:307:HOH:O	2.22	0.40

All (6) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:204:GOL:C2	8:A:373:HOH:O[2_455]	0.86	1.34
3:B:204:GOL:O2	8:A:373:HOH:O[2_455]	1.07	1.13
3:B:204:GOL:C3	8:A:373:HOH:O[2_455]	1.38	0.82
3:B:204:GOL:O3	8:A:373:HOH:O[2_455]	1.79	0.41
8:A:356:HOH:O	8:B:319:HOH:O[2_454]	1.99	0.21
8:A:381:HOH:O	8:B:323:HOH:O[2_455]	2.09	0.11

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	184/204 (90%)	179 (97%)	5 (3%)	0	100	100
1	B	193/204 (95%)	184 (95%)	4 (2%)	5 (3%)	4	0
All	All	377/408 (92%)	363 (96%)	9 (2%)	5 (1%)	9	2

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	82	LEU
1	B	81	GLY
1	B	189	LEU
1	B	190	LYS
1	B	80	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	172/190 (90%)	172 (100%)	0	100	100
1	B	181/190 (95%)	180 (99%)	1 (1%)	78	72
All	All	353/380 (93%)	352 (100%)	1 (0%)	86	82

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	82	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	31	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 8 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	GOL	A	203	-	5,5,5	0.34	0	5,5,5	0.85	0
6	PEG	A	208	7	6,6,6	0.38	0	5,5,5	0.30	0
3	GOL	B	204	-	5,5,5	0.31	0	5,5,5	0.59	0
6	PEG	A	207	-	6,6,6	0.49	0	5,5,5	0.48	0
3	GOL	B	203	-	5,5,5	0.34	0	5,5,5	0.35	0
4	XI7	A	205	2	14,14,14	1.63	2 (14%)	16,18,18	3.20	8 (50%)
3	GOL	A	204	-	5,5,5	0.60	0	5,5,5	1.50	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	203	-	-	4/4/4/4	-
6	PEG	A	208	7	-	1/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	204	-	-	0/4/4/4	-
6	PEG	A	207	-	-	2/4/4/4	-
3	GOL	B	203	-	-	2/4/4/4	-
4	XI7	A	205	2	-	0/12/12/12	0/1/1/1
3	GOL	A	204	-	-	4/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	205	XI7	O11-C10	-3.98	1.19	1.30
4	A	205	XI7	C9-C10	-3.70	1.47	1.53

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	205	XI7	O10-C10-C9	8.24	132.06	121.81
4	A	205	XI7	C2-C1-C6	4.74	124.59	118.57
4	A	205	XI7	O14-C7-C8	4.08	125.45	120.58
4	A	205	XI7	O14-C7-C1	-3.73	115.73	120.73
4	A	205	XI7	C5-C6-C1	-3.72	116.71	120.36
4	A	205	XI7	C3-C4-C5	2.61	123.45	119.87
4	A	205	XI7	O11-C10-O10	-2.54	117.85	123.90
3	A	204	GOL	O2-C2-C3	2.54	119.69	109.18
4	A	205	XI7	C3-C2-C1	-2.09	118.31	120.36

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	203	GOL	O1-C1-C2-C3
3	A	204	GOL	O1-C1-C2-C3
3	A	204	GOL	O2-C2-C3-O3
3	A	203	GOL	C1-C2-C3-O3
3	A	204	GOL	C1-C2-C3-O3
3	B	203	GOL	C1-C2-C3-O3
3	A	203	GOL	O1-C1-C2-O2
3	B	203	GOL	O2-C2-C3-O3
3	A	203	GOL	O2-C2-C3-O3
6	A	207	PEG	C1-C2-O2-C3
6	A	208	PEG	C1-C2-O2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	204	GOL	O1-C1-C2-O2
6	A	207	PEG	C4-C3-O2-C2

There are no ring outliers.

5 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	208	PEG	1	0
3	B	204	GOL	5	4
6	A	207	PEG	1	0
4	A	205	XI7	3	0
3	A	204	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	183/204 (89%)	0.64	21 (11%)	9 11	25, 43, 85, 115	7 (3%)
1	B	194/204 (95%)	0.63	22 (11%)	10 11	28, 44, 83, 116	1 (0%)
All	All	377/408 (92%)	0.63	43 (11%)	10 11	25, 44, 85, 116	8 (2%)

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	193	ILE	8.7
1	A	182	TYR	7.7
1	A	177	PHE	5.8
1	B	78	VAL	5.3
1	A	180	SER	4.7
1	A	169	TYR	4.7
1	B	80	PRO	4.5
1	B	192	THR	4.2
1	A	175	LYS	3.8
1	B	190	LYS	3.8
1	A	78	VAL	3.7
1	B	81	GLY	3.5
1	A	176	VAL	3.5
1	A	181	GLU	3.3
1	B	131	VAL	3.3
1	B	191	THR	3.2
1	A	80	PRO	3.2
1	B	83	PRO	3.2
1	A	172	GLN	3.1
1	B	169	TYR	3.0
1	A	179	GLU	3.0
1	A	173	ASP	2.9
1	A	170	SER	2.7
1	B	124	CYS	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	175	LYS	2.7
1	A	178	GLU	2.6
1	B	127	GLU	2.6
1	B	79	CYS	2.6
1	A	0	HIS	2.5
1	B	17	GLN	2.5
1	B	182	TYR	2.5
1	B	43	LYS	2.5
1	A	1	ASP	2.5
1	B	82	LEU	2.4
1	A	174	GLU	2.4
1	B	189	LEU	2.4
1	A	171	ASN	2.4
1	A	167	LEU	2.3
1	A	39	ILE	2.3
1	B	32	ARG	2.3
1	A	35	ARG	2.1
1	B	84	LEU	2.1
1	B	73	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	B	204	6/6	0.69	0.23	67,67,67,67	6
7	NA	A	209	1/1	0.69	0.17	89,89,89,89	0
3	GOL	B	203	6/6	0.78	0.16	69,82,88,89	0
7	NA	A	210	1/1	0.82	0.23	81,81,81,81	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	PEG	A	207	7/7	0.84	0.20	70,74,83,84	0
6	PEG	A	208	7/7	0.86	0.17	77,77,90,95	0
3	GOL	A	204	6/6	0.87	0.17	43,50,62,62	0
5	CL	A	206	1/1	0.90	0.19	112,112,112,112	0
3	GOL	A	203	6/6	0.91	0.13	41,50,53,54	6
5	CL	B	205	1/1	0.91	0.15	78,78,78,78	0
2	MG	B	201	1/1	0.93	0.10	72,72,72,72	0
2	MG	B	202	1/1	0.93	0.14	60,60,60,60	0
4	XI7	A	205	14/14	0.95	0.14	24,54,58,58	6
2	MG	A	202	1/1	0.97	0.05	42,42,42,42	0
2	MG	A	201	1/1	1.00	0.02	30,30,30,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.