



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 20, 2026 – 07:52 AM UTC

PDB ID : 6LT7 / pdb\_00006lt7  
Title : Crystal structure of human RPP20-RPP25 proteins in complex with the P3 domain of lncRNA RMRP  
Authors : Huang, J.; Yin, C.  
Deposited on : 2020-01-21  
Resolution : 2.70 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)
RNA backbone	3983	1044 (2.90-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	141	 64% 21% 12%
1	D	141	 65% 17% 6% 12%
2	B	198	 46% 12% 41%
2	E	198	 42% 16% 41%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	C	50	 50% 36% 10% ••
3	F	50	 42% 34% 14% 8% •

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6025 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribonuclease P protein subunit p20.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	S				Se
1	A	124	969	604	183	179	2	1	0	0	0
1	D	124	959	598	179	179	2	1	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP O75817
D	0	SER	-	expression tag	UNP O75817

- Molecule 2 is a protein called Ribonuclease P protein subunit p25.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	S				Se
2	B	116	871	546	168	152	2	3	0	0	0
2	E	116	871	545	167	154	2	3	0	0	0

- Molecule 3 is a RNA chain called 50-mer RNA.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
			Total	C	N	O				P
3	C	49	1030	460	172	349	49	0	0	0
3	F	49	1030	460	172	349	49	0	0	0

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	49	Total O 49 49	0	0
5	B	44	Total O 44 44	0	0
5	C	46	Total O 46 46	0	0
5	D	44	Total O 44 44	0	0
5	E	43	Total O 43 43	0	0
5	F	45	Total O 45 45	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ribonuclease P protein subunit p20

Chain A: 




- Molecule 1: Ribonuclease P protein subunit p20

Chain D: 



- Molecule 2: Ribonuclease P protein subunit p25

Chain B: 



- Molecule 2: Ribonuclease P protein subunit p25

Chain E: 



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	144.17Å 144.17Å 74.22Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	35.60 – 2.70 35.60 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.8 (35.60-2.70) 99.9 (35.60-2.70)	Depositor EDS
$R_{merge}$	0.19	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.58 (at 2.68Å)	Xtrriage
Refinement program	REFMAC 5.8.0232	Depositor
R, $R_{free}$	0.192 , 0.238 0.196 , 0.241	Depositor DCC
$R_{free}$ test set	2250 reflections (4.76%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	57.7	Xtrriage
Anisotropy	0.051	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 47.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.021 for -h,-k,l 0.468 for h,-h-k,-l 0.021 for -k,-h,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6025	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	67.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.94% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.13	1/982 (0.1%)	1.42	2/1327 (0.2%)
1	D	1.14	1/972 (0.1%)	1.37	0/1316
2	B	1.08	0/884	1.44	2/1190 (0.2%)
2	E	1.09	0/884	1.47	6/1191 (0.5%)
3	C	0.63	0/1145	1.14	9/1777 (0.5%)
3	F	0.62	0/1145	1.21	13/1777 (0.7%)
All	All	0.96	2/6012 (0.0%)	1.33	32/8578 (0.4%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	47	ALA	C-O	6.11	1.31	1.24
1	D	107	LEU	C-O	5.71	1.30	1.23

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	68	G	C3'-C2'-O2'	-13.40	94.50	114.60
3	C	68	G	C3'-C2'-O2'	-12.19	96.32	114.60
3	C	56	U	C4'-C3'-O3'	9.73	123.99	109.40
3	F	58	U	C2'-C3'-O3'	8.69	126.73	113.70
3	F	56	U	C2'-C3'-O3'	8.27	121.91	109.50

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	969	0	995	31	0
1	D	959	0	973	30	0
2	B	871	0	896	16	0
2	E	871	0	892	16	0
3	C	1030	0	528	12	0
3	F	1030	0	528	23	0
4	A	8	0	12	1	0
4	B	4	0	6	0	0
4	C	8	0	12	0	0
4	E	4	0	6	0	0
5	A	49	0	0	4	0
5	B	44	0	0	0	0
5	C	46	0	0	1	0
5	D	44	0	0	3	0
5	E	43	0	0	4	0
5	F	45	0	0	5	0
All	All	6025	0	4848	108	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 108 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:56:U:H4'	5:F:125:HOH:O	1.65	0.96
1:D:45:PHE:CD1	5:F:125:HOH:O	2.22	0.92
3:C:59:C:O2'	3:C:60:C:OP2	1.89	0.91
1:D:45:PHE:CG	5:F:125:HOH:O	2.23	0.90
3:F:66:G:N7	5:F:101:HOH:O	2.02	0.90

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	122/141 (86%)	112 (92%)	9 (7%)	1 (1%)	16	37
1	D	122/141 (86%)	113 (93%)	7 (6%)	2 (2%)	7	20
2	B	110/198 (56%)	104 (94%)	6 (6%)	0	100	100
2	E	110/198 (56%)	107 (97%)	2 (2%)	1 (1%)	14	35
All	All	464/678 (68%)	436 (94%)	24 (5%)	4 (1%)	14	35

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	E	12	ALA
1	A	27	PRO
1	D	27	PRO
1	D	62	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	105/117 (90%)	102 (97%)	3 (3%)	37	67
1	D	103/117 (88%)	93 (90%)	10 (10%)	8	20
2	B	90/146 (62%)	86 (96%)	4 (4%)	25	53
2	E	90/146 (62%)	84 (93%)	6 (7%)	15	36
All	All	388/526 (74%)	365 (94%)	23 (6%)	18	42

5 of 23 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	113	PRO
2	E	54	THR
1	D	125	ARG
2	E	84	LEU
2	B	129	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	101	ASN
1	D	64	ASN
1	D	101	ASN
1	D	127	ASN

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
3	C	48/50 (96%)	10 (20%)	5 (10%)
3	F	48/50 (96%)	14 (29%)	7 (14%)
All	All	96/100 (96%)	24 (25%)	12 (12%)

5 of 24 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
3	C	22	C
3	C	33	C
3	C	36	U
3	C	37	G
3	C	40	G

5 of 12 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
3	F	57	U
3	F	58	U
3	F	64	U
3	F	59	C
3	C	59	C

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	EDO	A	201	-	3,3,3	0.14	0	2,2,2	0.24	0
4	EDO	A	202	-	3,3,3	0.28	0	2,2,2	0.35	0
4	EDO	E	201	-	3,3,3	0.09	0	2,2,2	0.12	0
4	EDO	C	201	-	3,3,3	0.27	0	2,2,2	0.48	0
4	EDO	B	201	-	3,3,3	0.21	0	2,2,2	0.22	0
4	EDO	C	202	-	3,3,3	0.15	0	2,2,2	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	201	-	-	0/1/1/1	-
4	EDO	A	202	-	-	1/1/1/1	-
4	EDO	E	201	-	-	1/1/1/1	-
4	EDO	C	201	-	-	0/1/1/1	-
4	EDO	B	201	-	-	1/1/1/1	-
4	EDO	C	202	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	201	EDO	O1-C1-C2-O2
4	C	202	EDO	O1-C1-C2-O2
4	A	202	EDO	O1-C1-C2-O2
4	E	201	EDO	O1-C1-C2-O2


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	201	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	123/141 (87%)	-1.08	0 <a href="#">100</a> <a href="#">100</a>	35, 50, 121, 159	0
1	D	123/141 (87%)	-1.07	0 <a href="#">100</a> <a href="#">100</a>	34, 51, 122, 153	0
2	B	113/198 (57%)	-1.01	1 (0%) <a href="#">81</a> <a href="#">80</a>	35, 52, 114, 128	0
2	E	113/198 (57%)	-1.09	0 <a href="#">100</a> <a href="#">100</a>	35, 52, 116, 140	0
3	C	49/50 (98%)	-1.45	0 <a href="#">100</a> <a href="#">100</a>	38, 70, 141, 174	0
3	F	49/50 (98%)	-1.40	0 <a href="#">100</a> <a href="#">100</a>	40, 70, 143, 167	0
All	All	570/778 (73%)	-1.12	1 (0%) <a href="#">91</a> <a href="#">90</a>	34, 55, 125, 174	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	158	PRO	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	EDO	A	201	4/4	0.98	0.13	71,72,77,81	0
4	EDO	A	202	4/4	0.98	0.11	87,98,98,102	0
4	EDO	B	201	4/4	0.98	0.16	90,98,104,105	0
4	EDO	C	202	4/4	0.98	0.09	100,100,101,103	0
4	EDO	E	201	4/4	0.98	0.08	86,89,90,92	0
4	EDO	C	201	4/4	0.99	0.08	76,87,92,92	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around EDO A 202:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around EDO B 201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.