



wwPDB NMR Structure Validation Summary Report ⓘ

Mar 8, 2026 – 12:43 PM UTC

PDB ID : 1LUX / pdb_00001lux
Title : NMR SOLUTION STRUCTURE OF THE ANTICODON OF YEAST
TRNA-PHE WITH 3 MODIFICATIONS (OMC32 OMG34 M5C40)
Authors : Stuart, J.W.; Koshlap, K.M.; Guenther, R.H.; Agris, P.F.
Deposited on : 2002-05-23

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR


The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
RNA backbone	8273	777

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	17	 35% 35% 24% 6%

2 Ensemble composition and analysis ⓘ

This entry contains 10 models. This entry does not contain polypeptide chains, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.

3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 558 atoms, of which 193 are hydrogens and 0 are deuteriums.

- Molecule 1 is a RNA chain called 5'-R(*CP*CP*AP*GP*AP*(OMC)P*UP*(OMG)P*AP*AP*GP*AP*UP*(5MC)P*UP*GP*G)-3'.


Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	P	
1	A	17	558	166	193	68	115	16	0

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: 5'-R(*CP*CP*AP*GP*AP*(OMC)P*UP*(OMG)P*AP*AP*GP*AP*UP*(5MC)P*UP*GP*G)-3'



4.2 Residue scores for the first model from the NMR ensemble

No representative models were identified. Colouring as in section 4.1 above.

- Molecule 1: 5'-R(*CP*CP*AP*GP*AP*(OMC)P*UP*(OMG)P*AP*AP*GP*AP*UP*(5MC)P*UP*GP*G)-3'



5 Refinement protocol and experimental data overview ⓘ

The models were refined using the following method: *global fold by distance geometry. refinement by simulated annealing using the amber forcefield.*

Of the 50 calculated structures, 10 were deposited, based on the following criterion: *structures with the least restraint violations, structures with the lowest energy.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Discover	refinement	98

No chemical shift data was provided.

6 Model quality i

6.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: OMC, 5MC, OMG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.84±0.02	0±0/335 (0.0± 0.0%)	1.27±0.07	3±3/521 (0.5± 0.5%)
All	All	0.84	0/3350 (0.0%)	1.27	26/5210 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	4.7±1.2
All	All	0	47

There are no bond-length outliers.

5 of 13 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	35	A	O4'-C1'-N9	8.15	120.73	108.50	6	4
1	A	35	A	N9-C1'-C2'	7.96	123.94	112.00	2	4
1	A	34	OMG	O3'-P-O5'	-7.86	92.20	104.00	1	2
1	A	34	OMG	P-O3'-C3'	7.18	130.97	120.20	1	2
1	A	33	U	C3'-C2'-C1'	6.50	107.80	101.30	1	1

There are no chirality outliers.

5 of 13 unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	31	A	Sidechain	9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group	Models (Total)
1	A	33	U	Sidechain	8
1	A	35	A	Sidechain	6
1	A	37	G	Sidechain	5
1	A	41	U	Sidechain	3

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	365	193	192	1±1
All	All	3650	1930	1920	10

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

5 of 8 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:34:OMG:C8	1:A:35:A:C8	0.63	2.86	6	2
1:A:34:OMG:C8	1:A:34:OMG:H5''	0.55	2.37	6	2
1:A:34:OMG:C5'	1:A:34:OMG:C8	0.49	2.95	10	1
1:A:34:OMG:C8	1:A:35:A:C4	0.46	3.03	3	1
1:A:34:OMG:C8	1:A:34:OMG:C5'	0.43	3.01	6	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

6.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

6.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
1	A	16/17 (94%)	8±0 (50±0%)	2±1 (13±7%)	0.23±0.00
All	All	161/170 (95%)	80 (50%)	21 (13%)	0.23

The overall RNA backbone suiteness is 0.25.

5 of 8 unique RNA backbone outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
1	A	33	U	10
1	A	34	OMG	10
1	A	35	A	10
1	A	36	A	10
1	A	37	G	10

5 of 9 unique RNA pucker outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
1	A	34	OMG	6
1	A	35	A	3
1	A	39	U	3
1	A	33	U	2
1	A	37	G	2

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
1	OMC	A	32	1	19,22,23	0.49±0.02	0±0 (0±0%)
1	5MC	A	40	1	19,22,23	0.52±0.02	0±0 (0±0%)
1	OMG	A	34	1	23,26,27	0.71±0.04	0±0 (1±1%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types,

if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Counts	Bond angles	
						RMSZ	#Z>2
1	OMC	A	32	1	25,31,34	0.90±0.06	1±1 (5±3%)
1	5MC	A	40	1	26,32,35	0.88±0.03	1±1 (2±2%)
1	OMG	A	34	1	32,38,41	1.43±0.42	4±2 (11±6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	5MC	A	40	1	-	0±0,7,25,26	0±0,2,2,2
1	OMC	A	32	1	-	0±0,9,27,28	0±0,2,2,2
1	OMG	A	34	1	-	1±0,9,27,28	0±0,3,3,3

All unique bond outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	34	OMG	O4'-C4'	2.21	1.40	1.45	3	3

5 of 18 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	34	OMG	O4'-C4'-C3'	7.14	90.99	105.15	4	5
1	A	34	OMG	O2'-C2'-C1'	5.40	119.25	108.99	1	5
1	A	34	OMG	O3'-C3'-C4'	5.26	95.97	111.08	1	2
1	A	34	OMG	O4'-C4'-C5'	5.22	126.06	109.33	10	7
1	A	34	OMG	C2'-C3'-C4'	4.07	110.74	101.99	4	2

There are no chirality outliers.

All unique torsion outliers are listed below.

Mol	Chain	Res	Type	Atoms	Models (Total)
1	A	34	OMG	C1'-C2'-O2'-CM2	1

There are no ring outliers.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided