



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 12:06 PM UTC

PDB ID : 3LU8 / pdb_00003lu8
Title : Human serum albumin in complex with compound 3
Authors : Buttar, D.; Colclough, N.; Gerhardt, S.; MacFaul, P.A.; Phillips, S.D.; Plowright, A.; Whittamore, P.; Tam, K.; Maskos, K.; Steinbacher, S.; Steuber, H.
Deposited on : 2010-02-17
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	585	2% 82% 14% ..
1	B	585	% 84% 13% ..

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 9282 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serum albumin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	578	Total	C	N	O	S	0	2	0
			4609	2909	777	882	41			
1	B	573	Total	C	N	O	S	0	3	0
			4576	2889	776	870	41			

- Molecule 2 is N-[5-(5-[(2,4-dimethyl-1,3-thiazol-5-yl)sulfonyl]amino}-6-fluoropyridin-3-yl)-4-methyl-1,3-thiazol-2-yl]acetamide (CCD ID: IQX) (formula: C₁₆H₁₆FN₅O₃S₃).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	S	0	0
			28	16	1	5	3	3		
2	B	1	Total	C	F	N	O	S	0	0
			28	16	1	5	3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	24	Total 24	O 24	0	0
3	B	17	Total 17	O 17	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Serum albumin



• Molecule 1: Serum albumin



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	58.78Å 59.03Å 96.18Å 75.42° 88.06° 76.91°	Depositor
Resolution (Å)	21.74 – 2.60 21.74 – 2.60	Depositor EDS
% Data completeness (in resolution range)	90.1 (21.74-2.60) 90.1 (21.74-2.60)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.56 (at 2.60Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.245 , 0.266 (Not available) , 0.262	Depositor DCC
R_{free} test set	1342 reflections (3.58%)	wwPDB-VP
Wilson B-factor (Å ²)	44.3	Xtriage
Anisotropy	0.279	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 33.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	9282	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: IQX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.61	0/4704	0.84	1/6346 (0.0%)
1	B	0.61	0/4672	0.88	5/6301 (0.1%)
All	All	0.61	0/9376	0.86	6/12647 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	1	0
All	All	1	1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	170	GLN	N-CA-C	8.64	120.32	111.07
1	B	170	GLN	CB-CA-C	6.92	121.74	110.88
1	B	60	GLU	N-CA-C	5.85	118.41	111.33
1	B	170	GLN	N-CA-CB	5.44	117.90	110.01
1	B	271	ILE	CB-CA-C	-5.10	107.38	111.71
1	A	168	CYS	N-CA-C	5.09	116.91	111.36

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	B	170	GLN	CA

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	114	ARG	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4609	0	4528	45	0
1	B	4576	0	4511	41	0
2	A	28	0	16	0	0
2	B	28	0	16	0	0
3	A	24	0	0	0	0
3	B	17	0	0	1	0
All	All	9282	0	9071	86	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (86) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:503:ASN:HD21	1:A:505:GLU:HB3	1.56	0.69
1:B:558:CYS:SG	1:B:567:CYS:O	2.56	0.63
1:B:356:THR:HG21	1:B:373:VAL:HG22	1.83	0.59
1:B:356:THR:HG21	1:B:373:VAL:CG2	2.33	0.59
1:B:539:ALA:O	1:B:540:THR:HG23	2.05	0.57
1:A:114:ARG:O	1:A:114:ARG:HG3	2.04	0.57
1:B:117:ARG:HB2	1:B:123:MET:HE1	1.87	0.56
1:A:408:LEU:HD22	1:A:530:VAL:HG22	1.87	0.56
1:B:222:ARG:HG2	1:B:223:PHE:CE1	2.41	0.55
1:A:348:ARG:HD3	1:A:483:ASN:ND2	2.21	0.55
1:A:387:LEU:HD22	1:A:485:ARG:NH1	2.22	0.55
1:B:563:ASP:OD2	1:B:565:GLU:HA	2.08	0.54
1:A:81:ARG:HG2	1:A:88:ALA:HB3	1.89	0.53
1:A:348:ARG:HD3	1:A:483:ASN:HD21	1.72	0.52
1:B:503[B]:ASN:CG	1:B:506:THR:HB	2.34	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:118:PRO:HD2	1:A:123:MET:HE3	1.92	0.52
1:B:27:PHE:CD1	1:B:74:LEU:HD21	2.44	0.52
1:A:419:SER:OG	1:A:421:PRO:HD2	2.10	0.51
1:B:308:ASP:OD1	1:B:308:ASP:N	2.43	0.51
1:B:538:LYS:O	1:B:539:ALA:C	2.54	0.51
1:A:356:THR:HG21	1:A:373:VAL:HG22	1.92	0.51
1:B:385:GLN:HG2	1:B:446:MET:HE2	1.93	0.50
1:B:67:HIS:NE2	1:B:249:ASP:OD1	2.44	0.50
1:A:59:ALA:HB3	1:A:62:CYS:SG	2.52	0.49
1:A:111:ASN:N	1:A:111:ASN:HD22	2.09	0.49
1:B:333:GLU:OE1	1:B:337:ARG:NH2	2.45	0.49
1:A:19:PHE:O	1:A:23:VAL:HG23	2.13	0.48
1:B:475:LYS:O	1:B:479:GLU:HB2	2.14	0.48
1:A:118:PRO:HD2	1:A:123:MET:CE	2.44	0.48
1:B:492:GLU:HA	1:B:492:GLU:OE1	2.14	0.48
1:B:506:THR:HG22	1:B:506:THR:O	2.14	0.48
1:A:392:CYS:O	1:A:396:GLU:HG2	2.15	0.47
1:B:30:TYR:CD1	1:B:102:PHE:HB3	2.49	0.47
1:A:313:LYS:HA	1:A:367:HIS:NE2	2.30	0.46
1:A:323:LYS:HG3	1:A:324:ASP:N	2.29	0.46
1:A:488:PHE:HA	1:A:491:LEU:HD12	1.96	0.46
1:B:402:LYS:NZ	3:B:600:HOH:O	2.48	0.46
1:A:538:LYS:O	1:A:540:THR:N	2.49	0.46
1:A:122:VAL:O	1:A:125:THR:HB	2.16	0.45
1:B:446:MET:N	1:B:447:PRO:HD2	2.30	0.45
1:A:414:LYS:HD2	1:A:491:LEU:O	2.16	0.45
1:B:345:LEU:O	1:B:349:LEU:HG	2.16	0.45
1:B:422:THR:O	1:B:426:VAL:HG23	2.16	0.45
1:B:408:LEU:HD22	1:B:530:VAL:CG2	2.46	0.45
1:A:198:LEU:O	1:A:202:SER:OG	2.32	0.44
1:B:450:GLU:HG3	1:B:485:ARG:HD3	1.99	0.44
1:B:39:HIS:O	1:B:43:VAL:HG23	2.18	0.44
1:B:196:GLN:NE2	1:B:196:GLN:HA	2.32	0.44
1:A:114:ARG:HH22	1:A:520:GLU:CD	2.25	0.44
1:A:263:TYR:CE1	1:A:267:ASN:OD1	2.70	0.44
1:B:151:ALA:HB3	1:B:152:PRO:HD3	1.99	0.44
1:B:34:CYS:HB3	1:B:39:HIS:HE2	1.83	0.44
1:B:302:LEU:HD23	1:B:303:PRO:HD2	1.99	0.43
1:A:503:ASN:ND2	1:A:505:GLU:HB3	2.29	0.43
1:B:408:LEU:HD11	1:B:526:GLN:HB3	2.00	0.43
1:A:34:CYS:HB3	1:A:39:HIS:NE2	2.34	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:42:LEU:HD22	1:B:73:LYS:HG3	2.00	0.42
1:A:106:LYS:NZ	1:A:147:PRO:O	2.42	0.42
1:A:29:GLN:HG2	1:A:147:PRO:HA	2.02	0.42
1:A:517:SER:O	1:A:518:GLU:C	2.62	0.42
1:B:117:ARG:HH21	1:B:179:LEU:HD22	1.83	0.42
1:B:392:CYS:O	1:B:396:GLU:HG2	2.20	0.42
1:A:420:THR:O	1:A:424:VAL:HG23	2.19	0.42
1:A:536:LYS:O	1:A:540:THR:CG2	2.67	0.42
1:B:398:LEU:HB3	1:B:402:LYS:HB2	2.02	0.41
1:A:383:GLU:HB3	1:A:384:PRO:HD3	2.01	0.41
1:B:45:GLU:OE1	1:B:73:LYS:HE3	2.20	0.41
1:A:383:GLU:HA	1:A:386[B]:ASN:HD22	1.85	0.41
1:B:561:ALA:O	1:B:562:ASP:C	2.64	0.41
1:B:566:THR:C	1:B:568:PHE:H	2.29	0.41
1:A:310:VAL:O	1:A:370:TYR:HE1	2.04	0.41
1:A:356:THR:HG21	1:A:373:VAL:CG2	2.51	0.41
1:A:464:HIS:CE1	1:A:470:SER:HB3	2.55	0.41
1:A:575:LEU:HA	1:A:578:ALA:HB3	2.02	0.41
1:B:137:LYS:O	1:B:141:GLU:HG2	2.21	0.41
1:B:503[A]:ASN:HD22	1:B:506:THR:CB	2.33	0.41
1:A:39:HIS:O	1:A:43:VAL:HG23	2.20	0.41
1:A:464:HIS:CE1	1:A:469:VAL:H	2.39	0.41
1:A:420:THR:N	1:A:421:PRO:CD	2.84	0.40
1:A:117:ARG:NH2	1:A:182:LEU:HB2	2.36	0.40
1:A:485:ARG:HB3	1:A:486:PRO:HD3	2.03	0.40
1:A:513:ILE:HG23	1:A:514:CYS:N	2.36	0.40
1:A:523:ILE:O	1:A:527:THR:OG1	2.29	0.40
1:A:261:ALA:HB1	1:A:286:LYS:HE3	2.03	0.40
1:B:34:CYS:HB3	1:B:39:HIS:NE2	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	578/585 (99%)	533 (92%)	38 (7%)	7 (1%)	10	23
1	B	572/585 (98%)	538 (94%)	28 (5%)	6 (1%)	12	28
All	All	1150/1170 (98%)	1071 (93%)	66 (6%)	13 (1%)	11	25

All (13) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	115	LEU
1	B	501	GLU
1	A	538	LYS
1	A	539	ALA
1	A	364	ALA
1	A	505	GLU
1	A	541	LYS
1	B	80	LEU
1	B	506	THR
1	B	540	THR
1	A	518	GLU
1	B	567	CYS
1	B	364	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	508/511 (99%)	476 (94%)	32 (6%)	16	36
1	B	505/511 (99%)	484 (96%)	21 (4%)	26	52
All	All	1013/1022 (99%)	960 (95%)	53 (5%)	20	44

All (53) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	GLU
1	A	79	THR
1	A	81	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	111	ASN
1	A	117	ARG
1	A	119	GLU
1	A	120	VAL
1	A	129	ASP
1	A	137	LYS
1	A	167	GLU
1	A	236	THR
1	A	245	CYS
1	A	252	GLU
1	A	268	GLN
1	A	292	GLU
1	A	324	ASP
1	A	344	VAL
1	A	367	HIS
1	A	382	GLU
1	A	414	LYS
1	A	419	SER
1	A	420	THR
1	A	469	VAL
1	A	496	THR
1	A	498	VAL
1	A	503	ASN
1	A	508	THR
1	A	509	PHE
1	A	524	LYS
1	A	527	THR
1	A	559	CYS
1	A	571	GLU
1	B	77	VAL
1	B	114	ARG
1	B	115	LEU
1	B	116	VAL
1	B	131	GLU
1	B	166	THR
1	B	245	CYS
1	B	270	SER
1	B	308	ASP
1	B	344	VAL
1	B	387	LEU
1	B	408	LEU
1	B	414	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	442	GLU
1	B	492	GLU
1	B	496	THR
1	B	508	THR
1	B	540	THR
1	B	543	GLN
1	B	554	PHE
1	B	565	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	GLN
1	A	94	GLN
1	A	104	GLN
1	A	109	ASN
1	A	111	ASN
1	A	196	GLN
1	A	440	HIS
1	A	464	HIS
1	A	483	ASN
1	A	503	ASN
1	A	580	GLN
1	B	9	HIS
1	B	196	GLN
1	B	367	HIS
1	B	464	HIS
1	B	543	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	IQX	B	586	-	27,30,30	2.69	7 (25%)	37,45,45	4.55	22 (59%)
2	IQX	A	586	-	27,30,30	2.63	6 (22%)	37,45,45	4.20	20 (54%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	IQX	B	586	-	-	3/19/19/19	0/3/3/3
2	IQX	A	586	-	-	7/19/19/19	0/3/3/3

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	586	IQX	C10-S11	-8.47	1.59	1.73
2	A	586	IQX	C10-S11	-8.43	1.59	1.73
2	B	586	IQX	C21-S23	-6.05	1.60	1.72
2	A	586	IQX	C21-S23	-6.01	1.60	1.72
2	B	586	IQX	C7-S11	-5.54	1.60	1.74
2	A	586	IQX	C7-S11	-5.38	1.61	1.74
2	B	586	IQX	C24-S23	-5.10	1.60	1.73
2	A	586	IQX	C24-S23	-4.94	1.61	1.73
2	B	586	IQX	C10-N13	2.63	1.42	1.38
2	A	586	IQX	C10-N13	2.54	1.42	1.38
2	A	586	IQX	C10-N9	2.35	1.34	1.31
2	B	586	IQX	F28-C2	-2.10	1.32	1.34
2	B	586	IQX	C10-N9	2.07	1.34	1.31

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	586	IQX	C10-S11-C7	17.22	99.07	88.35
2	A	586	IQX	C10-S11-C7	14.88	97.61	88.35
2	A	586	IQX	C3-N1-C2	9.37	120.55	115.88
2	B	586	IQX	C3-N1-C2	8.39	120.06	115.88
2	B	586	IQX	S11-C10-N9	-7.73	108.87	115.78
2	B	586	IQX	S23-C24-N25	-7.60	109.40	114.64
2	A	586	IQX	S11-C10-N9	-6.98	109.53	115.78
2	A	586	IQX	S23-C24-N25	-6.71	110.02	114.64
2	B	586	IQX	C8-C7-S11	-6.66	105.86	109.91
2	B	586	IQX	C21-S23-C24	6.49	98.43	89.70
2	B	586	IQX	O20-S18-O19	-6.18	108.49	119.82
2	A	586	IQX	O20-S18-O19	-6.02	108.78	119.82
2	A	586	IQX	C21-S23-C24	5.77	97.47	89.70
2	B	586	IQX	O20-S18-C21	4.54	113.43	107.36
2	A	586	IQX	C4-C2-N1	-4.43	120.20	128.34
2	B	586	IQX	C4-C2-N1	-4.28	120.48	128.34
2	A	586	IQX	C8-C7-S11	-4.16	107.38	109.91
2	A	586	IQX	C3-C5-C7	-4.10	117.09	120.51
2	A	586	IQX	C15-C14-N13	4.10	121.88	115.27
2	B	586	IQX	C2-C4-N17	-3.94	114.39	120.32
2	A	586	IQX	C10-N9-C8	3.87	112.74	110.38
2	B	586	IQX	C10-N9-C8	3.76	112.67	110.38
2	B	586	IQX	C15-C14-N13	3.72	121.27	115.27
2	A	586	IQX	C7-C8-N9	-3.26	112.73	115.01
2	B	586	IQX	O19-S18-C21	3.16	111.58	107.36
2	A	586	IQX	C5-C7-S11	3.09	125.19	117.32
2	B	586	IQX	S11-C10-N13	2.92	127.63	123.72
2	A	586	IQX	C2-C4-N17	-2.90	115.94	120.32
2	A	586	IQX	S11-C10-N13	2.82	127.50	123.72
2	A	586	IQX	O20-S18-C21	2.81	111.11	107.36
2	A	586	IQX	C6-C5-C7	2.70	123.94	120.36
2	B	586	IQX	C6-C4-N17	2.50	126.26	120.84
2	A	586	IQX	C5-C7-C8	-2.50	127.43	130.81
2	B	586	IQX	F28-C2-N1	2.48	121.19	115.14
2	B	586	IQX	C5-C7-S11	2.45	123.55	117.32
2	A	586	IQX	O19-S18-C21	2.42	110.59	107.36
2	B	586	IQX	C6-C5-C7	2.26	123.37	120.36
2	B	586	IQX	C3-C5-C7	-2.22	118.66	120.51
2	B	586	IQX	C14-N13-C10	2.15	127.59	124.05
2	B	586	IQX	C7-C8-N9	-2.12	113.53	115.01
2	B	586	IQX	O16-C14-C15	-2.12	118.29	122.05
2	A	586	IQX	C5-C3-N1	-2.07	120.55	123.59

There are no chirality outliers.


All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	586	IQX	N9-C10-N13-C14
2	A	586	IQX	S11-C10-N13-C14
2	A	586	IQX	C2-C4-N17-S18
2	B	586	IQX	C2-C4-N17-S18
2	A	586	IQX	C4-N17-S18-O19
2	B	586	IQX	C4-N17-S18-O19
2	A	586	IQX	S23-C21-S18-N17
2	A	586	IQX	C22-C21-S18-O19
2	A	586	IQX	S23-C21-S18-O19
2	B	586	IQX	S23-C21-S18-O19

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	578/585 (98%)	0.08	13 (2%) 62 57	22, 40, 59, 75	15 (2%)
1	B	573/585 (97%)	-0.01	5 (0%) 81 78	16, 41, 58, 66	13 (2%)
All	All	1151/1170 (98%)	0.04	18 (1%) 70 66	16, 41, 59, 75	28 (2%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	83	THR	3.7
1	A	568	PHE	3.4
1	A	574	LYS	3.3
1	A	555	VAL	3.3
1	B	539	ALA	3.2
1	A	575	LEU	3.2
1	A	79	THR	2.7
1	B	549	ASP	2.7
1	A	567	CYS	2.7
1	B	567	CYS	2.5
1	B	87	MET	2.4
1	B	563	ASP	2.4
1	A	111	ASN	2.2
1	A	563	ASP	2.2
1	A	517	SER	2.2
1	A	301	ASP	2.1
1	A	283	LEU	2.1
1	A	80	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	IQX	A	586	28/28	0.89	0.10	47,61,69,77	0
2	IQX	B	586	28/28	0.92	0.09	51,55,65,67	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around IQX A 586:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around IQX B 586:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.