



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 06:24 PM UTC

PDB ID : 4LUS / pdb_00004lus
Title : alanine racemase [Clostridium difficile 630]
Authors : Asojo, O.A.
Deposited on : 2013-07-25
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	385	2% 45% 42% 6% 6%
1	C	385	10% 54% 36% 9% .
1	D	385	11% 55% 36% 7% ..
2	B	385	4% 51% 39% 7% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	401	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 12484 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Alanine racemase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	362	Total	C	N	O	P	S	0	0	0
			2880	1828	481	554	1	16			
1	C	382	Total	C	N	O	P	S	0	0	0
			3029	1923	504	585	1	16			
1	D	380	Total	C	N	O	P	S	0	0	0
			3013	1912	501	583	1	16			

- Molecule 2 is a protein called Alanine racemase.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	374	Total	C	N	O	P	S	0	1	0
			2972	1888	493	575	1	15			

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is 3,3',3''-phosphanetriyltripropanoic acid (CCD ID: TCE) (formula: $C_9H_{15}O_6P$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	O	P	0	0
			16	9	6	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	186	Total	O	0	0
			186	186		
5	B	150	Total	O	0	0
			150	150		
5	C	125	Total	O	0	0
			125	125		
5	D	107	Total	O	0	0
			107	107		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Alanine racemase




• Molecule 1: Alanine racemase






• Molecule 1: Alanine racemase



• Molecule 2: Alanine racemase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	85.15Å 93.30Å 107.09Å 90.00° 91.00° 90.00°	Depositor
Resolution (Å)	54.54 – 2.10 54.54 – 2.10	Depositor EDS
% Data completeness (in resolution range)	95.1 (54.54-2.10) 95.1 (54.54-2.10)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	12.41 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.202 , 0.266 0.215 , 0.264	Depositor DCC
R_{free} test set	4555 reflections (4.50%)	wwPDB-VP
Wilson B-factor (Å ²)	34.9	Xtriage
Anisotropy	0.018	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 37.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.013 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	12484	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: KCX, GOL, TCE, LLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	2.24	149/2881 (5.2%)	1.30	19/3883 (0.5%)
1	C	1.98	67/3034 (2.2%)	1.25	21/4092 (0.5%)
1	D	1.93	64/3017 (2.1%)	1.27	23/4068 (0.6%)
2	B	2.04	98/2988 (3.3%)	1.28	29/4029 (0.7%)
All	All	2.05	378/11920 (3.2%)	1.28	92/16072 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
2	B	0	2
All	All	0	4

The worst 5 of 378 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	362	ILE	C-O	-10.06	1.14	1.24
1	A	85	LEU	C-O	-9.61	1.12	1.24
1	A	78	ILE	C-O	-8.24	1.15	1.24
1	A	186	TYR	C-O	-8.19	1.14	1.24
2	B	51	ALA	C-O	-8.14	1.14	1.24

The worst 5 of 92 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	276	GLY	N-CA-C	-11.44	100.52	112.08
1	A	199	LYS	N-CA-C	11.07	123.09	111.14

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	362	ILE	O-C-N	-10.87	111.59	122.67
1	A	3	LYS	O-C-N	-10.32	106.49	123.00
1	C	162	GLU	N-CA-C	9.58	121.80	111.36

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	3	LYS	Peptide,Mainchain
2	B	218	ARG	Mainchain
2	B	362	ILE	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2880	0	2926	56	0
1	C	3029	0	3076	120	0
1	D	3013	0	3052	101	0
2	B	2972	0	3010	76	0
3	A	6	0	8	17	0
4	B	16	0	12	6	0
5	A	186	0	0	5	0
5	B	150	0	0	2	0
5	C	125	0	0	6	0
5	D	107	0	0	4	0
All	All	12484	0	12084	336	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 336 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:43:TYR:CE1	1:D:315:MET:HE1	1.57	1.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:259:GLN:OE1	1:D:279:THR:HG22	1.49	1.12
2:B:264:VAL:H	2:B:274:THR:HG22	1.01	1.12
2:B:264:VAL:H	2:B:274:THR:CG2	1.61	1.11
1:C:96:SER:CB	1:C:103:MET:HE2	1.85	1.05

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	356/385 (92%)	346 (97%)	9 (2%)	1 (0%)	36	36
1	C	378/385 (98%)	365 (97%)	11 (3%)	2 (0%)	24	22
1	D	374/385 (97%)	365 (98%)	9 (2%)	0	100	100
2	B	368/385 (96%)	360 (98%)	6 (2%)	2 (0%)	24	22
All	All	1476/1540 (96%)	1436 (97%)	35 (2%)	5 (0%)	36	36

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	219	LEU
1	C	267	SER
1	C	58	LYS
2	B	200	ILE
1	A	200	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	313/332 (94%)	302 (96%)	11 (4%)	32	35
1	C	329/332 (99%)	309 (94%)	20 (6%)	17	15
1	D	327/332 (98%)	309 (94%)	18 (6%)	19	18
2	B	324/333 (97%)	312 (96%)	12 (4%)	30	33
All	All	1293/1329 (97%)	1232 (95%)	61 (5%)	23	24

5 of 61 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	127	VAL
1	D	274	THR
1	C	240	ASP
1	D	254	ILE
1	D	316	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	148	GLN
1	D	374	ASN
1	D	256	HIS
1	C	75	GLN
1	C	378	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

7 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	LLP	A	39	1	23,24,25	2.63	9 (39%)	25,32,34	1.50	6 (24%)
1	LLP	D	39	1	23,24,25	2.16	9 (39%)	25,32,34	2.16	10 (40%)
1	KCX	C	130	1	10,11,12	1.28	1 (10%)	6,12,14	0.90	0
1	LLP	C	39	1	23,24,25	2.22	8 (34%)	25,32,34	1.91	8 (32%)
1	KCX	D	130	1	10,11,12	1.26	0	6,12,14	2.14	2 (33%)
1	KCX	A	130	1	10,11,12	1.94	5 (50%)	6,12,14	1.69	1 (16%)
2	LLP	B	39	2	23,24,25	2.44	11 (47%)	25,32,34	1.50	6 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	A	39	1	-	5/16/17/19	0/1/1/1
1	LLP	D	39	1	-	3/16/17/19	0/1/1/1
1	KCX	C	130	1	-	3/9/10/12	-
1	LLP	C	39	1	-	6/16/17/19	0/1/1/1
1	KCX	D	130	1	-	2/9/10/12	-
1	KCX	A	130	1	-	4/9/10/12	-
2	LLP	B	39	2	-	3/16/17/19	0/1/1/1

The worst 5 of 43 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	39	LLP	C3-C2	5.99	1.47	1.41
1	D	39	LLP	C4'-NZ	5.44	1.45	1.27
1	A	39	LLP	C4'-NZ	5.41	1.45	1.27
2	B	39	LLP	C3-C2	4.99	1.46	1.41
1	C	39	LLP	C4'-NZ	4.83	1.43	1.27

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	39	LLP	C4-C4'-NZ	-5.08	100.59	124.04
1	C	39	LLP	CD-CE-NZ	4.10	121.68	110.83

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	39	LLP	C5-C4-C4'	3.85	127.41	121.47
1	C	39	LLP	C4-C4'-NZ	-3.84	106.30	124.04
2	B	39	LLP	C4-C3-C2	-3.76	118.03	120.14

There are no chirality outliers.

5 of 26 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	39	LLP	C4-C4'-NZ-CE
1	A	130	KCX	N-CA-CB-CG
1	A	130	KCX	C-CA-CB-CG
2	B	39	LLP	C4-C4'-NZ-CE
1	C	39	LLP	C4-C4'-NZ-CE

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	C	130	KCX	1	0
1	C	39	LLP	1	0
1	A	130	KCX	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	TCE	B	401	-	12,15,15	1.25	0	12,18,18	3.16	3 (25%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	A	401	-	5,5,5	1.20	0	5,5,5	2.17	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TCE	B	401	-	-	12/15/15/15	-
3	GOL	A	401	-	-	3/4/4/4	-

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	401	TCE	C1-P-C3	5.99	120.03	100.95
4	B	401	TCE	C1-P-C2	5.98	120.00	100.95
4	B	401	TCE	C3-P-C2	5.97	119.97	100.95
3	A	401	GOL	O1-C1-C2	3.41	125.75	110.38
3	A	401	GOL	O3-C3-C2	2.62	122.17	110.38

There are no chirality outliers.

5 of 15 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	401	GOL	O1-C1-C2-C3
4	B	401	TCE	P-C2-C5-C14
4	B	401	TCE	C5-C2-P-C3
4	B	401	TCE	C5-C2-P-C1
4	B	401	TCE	C6-C3-P-C2


There are no ring outliers.

2 monomers are involved in 23 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	401	TCE	6	0
3	A	401	GOL	17	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	360/385 (93%)	-0.02	9 (2%) 58 61	16, 28, 49, 73	0
1	C	380/385 (98%)	0.69	38 (10%) 12 13	23, 42, 63, 85	0
1	D	378/385 (98%)	0.73	44 (11%) 9 10	21, 43, 72, 92	0
2	B	373/385 (96%)	0.24	14 (3%) 44 46	10, 35, 63, 81	1 (0%)
All	All	1491/1540 (96%)	0.42	105 (7%) 22 24	10, 37, 65, 92	1 (0%)

The worst 5 of 105 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	332	ASP	6.9
1	D	188	PHE	5.2
1	D	273	THR	5.1
1	D	216	ASP	4.3
1	C	266	ILE	4.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	KCX	D	130	12/13	0.85	0.14	36,45,56,61	0
1	KCX	C	130	12/13	0.88	0.12	40,45,50,56	0
1	KCX	A	130	12/13	0.93	0.10	20,29,37,37	0
1	LLP	C	39	24/25	0.95	0.08	28,33,42,47	0
2	LLP	B	39	24/25	0.96	0.07	16,28,32,34	0
1	LLP	D	39	24/25	0.96	0.07	25,34,42,43	0
1	LLP	A	39	24/25	0.96	0.07	15,23,29,36	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	TCE	B	401	16/16	0.74	0.17	40,48,58,59	0
3	GOL	A	401	6/6	0.92	0.18	21,29,39,54	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around TCE B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.