



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 13, 2026 – 10:08 AM UTC

PDB ID : 5LUP / pdb_00005lup
Title : Structures of DHBN domain of human BLM helicase
Authors : Shi, J.; Chen, W.-F.; Zhang, B.; Fan, S.-H.; Ai, X.; Liu, N.-N.; Rety, S.; Xi, X.-G.
Deposited on : 2016-09-09
Resolution : 2.03 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2260 (2.04-2.04)
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	54	
1	B	54	
1	C	54	
1	D	54	
1	E	54	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	54	 74% 22% .
1	G	54	 76% 17% . .
1	I	54	 74% 19% . 6%
1	J	54	 74% 15% . 9%
1	K	54	 70% 17% 6% 7%
1	L	54	 81% 11% 7%
2	H	54	 81% 13% . .

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5543 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BLM protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	50	Total 415	C 259	N 78	O 75	S 3	0	0	0
1	B	51	Total 419	C 261	N 79	O 76	S 3	0	0	0
1	C	54	Total 443	C 276	N 82	O 81	S 4	0	0	0
1	D	53	Total 435	C 271	N 81	O 80	S 3	0	0	0
1	E	53	Total 435	C 271	N 81	O 80	S 3	0	0	0
1	F	52	Total 428	C 266	N 80	O 79	S 3	0	0	0
1	G	52	Total 428	C 266	N 80	O 79	S 3	0	0	0
1	I	51	Total 420	C 262	N 79	O 76	S 3	0	0	0
1	J	49	Total 406	C 254	N 77	O 72	S 3	0	0	0
1	K	50	Total 415	C 259	N 78	O 75	S 3	0	0	0
1	L	50	Total 415	C 259	N 78	O 75	S 3	0	0	0

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	361	MET	-	initiating methionine	UNP Q3B7X0
B	361	MET	-	initiating methionine	UNP Q3B7X0
C	361	MET	-	initiating methionine	UNP Q3B7X0
D	361	MET	-	initiating methionine	UNP Q3B7X0
E	361	MET	-	initiating methionine	UNP Q3B7X0
F	361	MET	-	initiating methionine	UNP Q3B7X0
G	361	MET	-	initiating methionine	UNP Q3B7X0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
I	361	MET	-	initiating methionine	UNP Q3B7X0
J	361	MET	-	initiating methionine	UNP Q3B7X0
K	361	MET	-	initiating methionine	UNP Q3B7X0
L	361	MET	-	initiating methionine	UNP Q3B7X0


- Molecule 2 is a protein called BLM protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	52	428	266	80	79	3	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	361	MET	-	initiating methionine	UNP Q3B7X0
H	366	LEU	ILE	conflict	UNP Q3B7X0

- Molecule 3 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
3	A	1	5	4	1	0	0
3	J	1	5	4	1	0	0

- Molecule 4 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total K 1 1	0	0
4	D	1	Total K 1 1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	38	Total O 38 38	0	0
5	B	39	Total O 39 39	0	0
5	C	50	Total O 50 50	0	0
5	D	42	Total O 42 42	0	0
5	E	36	Total O 36 36	0	0
5	F	35	Total O 35 35	0	0
5	G	36	Total O 36 36	0	0
5	H	43	Total O 43 43	0	0
5	I	38	Total O 38 38	0	0
5	J	35	Total O 35 35	0	0
5	K	32	Total O 32 32	0	0
5	L	20	Total O 20 20	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: BLM protein

Chain A:  76% 13% • 7%




- Molecule 1: BLM protein

Chain B:  83% 11% 6%




- Molecule 1: BLM protein

Chain C:  83% 17%




- Molecule 1: BLM protein

Chain D:  81% 15% ••



- Molecule 1: BLM protein

Chain E:  85% 11% ••



- Molecule 1: BLM protein

Chain F:  74% 22% •



- Molecule 1: BLM protein




- Molecule 1: BLM protein




- Molecule 1: BLM protein




- Molecule 1: BLM protein



- Molecule 1: BLM protein



- Molecule 2: BLM protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 62	Depositor
Cell constants a, b, c, α , β , γ	132.79Å 132.79Å 64.98Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.47 – 2.03 43.47 – 2.03	Depositor EDS
% Data completeness (in resolution range)	98.0 (43.47-2.03) 98.0 (43.47-2.03)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.84 (at 2.03Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.210 , 0.263 (Not available) , 0.244	Depositor DCC
R_{free} test set	1987 reflections (4.70%)	wwPDB-VP
Wilson B-factor (Å ²)	31.0	Xtrriage
Anisotropy	0.073	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 35.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.429 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5543	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 32.58 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.1889e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.50	0/417	0.75	0/558
1	B	0.49	0/421	0.74	0/564
1	C	0.48	0/445	0.75	0/596
1	D	0.51	0/437	0.71	0/586
1	E	0.51	0/437	0.75	0/586
1	F	0.47	0/430	0.76	0/576
1	G	0.45	0/430	0.77	0/576
1	I	0.51	0/422	0.70	0/565
1	J	0.46	0/408	0.72	0/546
1	K	0.41	0/417	0.80	0/558
1	L	0.44	0/417	0.75	0/558
2	H	0.51	0/430	0.87	0/576
All	All	0.48	0/5111	0.76	0/6845

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	415	0	444	11	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	419	0	447	3	0
1	C	443	0	471	6	0
1	D	435	0	461	12	0
1	E	435	0	462	8	0
1	F	428	0	453	11	0
1	G	428	0	453	12	0
1	I	420	0	449	12	0
1	J	406	0	438	19	0
1	K	415	0	444	13	0
1	L	415	0	444	10	0
2	H	428	0	453	11	0
3	A	5	0	0	0	0
3	J	5	0	0	1	0
4	B	1	0	0	0	0
4	D	1	0	0	0	0
5	A	38	0	0	10	0
5	B	39	0	0	1	0
5	C	50	0	0	3	0
5	D	42	0	0	11	0
5	E	36	0	0	7	0
5	F	35	0	0	6	0
5	G	36	0	0	7	0
5	H	43	0	0	5	0
5	I	38	0	0	8	0
5	J	35	0	0	17	0
5	K	32	0	0	9	0
5	L	20	0	0	8	0
All	All	5543	0	5419	123	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 123 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:J:364:ARG:N	5:J:601:HOH:O	1.80	1.14
5:D:601:HOH:O	1:G:369:GLN:NE2	1.83	1.10
1:J:398:ASN:ND2	5:J:602:HOH:O	1.86	1.07
1:I:378:HIS:ND1	5:I:501:HOH:O	1.86	1.07
1:G:378:HIS:CG	5:G:502:HOH:O	2.08	1.04

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	48/54 (89%)	48 (100%)	0	0	100	100
1	B	49/54 (91%)	49 (100%)	0	0	100	100
1	C	52/54 (96%)	49 (94%)	2 (4%)	1 (2%)	6	2
1	D	51/54 (94%)	49 (96%)	1 (2%)	1 (2%)	6	1
1	E	51/54 (94%)	51 (100%)	0	0	100	100
1	F	50/54 (93%)	49 (98%)	1 (2%)	0	100	100
1	G	50/54 (93%)	49 (98%)	1 (2%)	0	100	100
1	I	49/54 (91%)	49 (100%)	0	0	100	100
1	J	47/54 (87%)	47 (100%)	0	0	100	100
1	K	48/54 (89%)	47 (98%)	1 (2%)	0	100	100
1	L	48/54 (89%)	47 (98%)	1 (2%)	0	100	100
2	H	50/54 (93%)	50 (100%)	0	0	100	100
All	All	593/648 (92%)	584 (98%)	7 (1%)	2 (0%)	36	30

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	413	GLU
1	D	363	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	49/52 (94%)	45 (92%)	4 (8%)	10	5
1	B	49/52 (94%)	47 (96%)	2 (4%)	27	21
1	C	52/52 (100%)	51 (98%)	1 (2%)	50	50
1	D	51/52 (98%)	51 (100%)	0	100	100
1	E	51/52 (98%)	49 (96%)	2 (4%)	28	23
1	F	50/52 (96%)	48 (96%)	2 (4%)	28	22
1	G	50/52 (96%)	47 (94%)	3 (6%)	17	10
1	I	49/52 (94%)	47 (96%)	2 (4%)	27	21
1	J	48/52 (92%)	47 (98%)	1 (2%)	47	46
1	K	49/52 (94%)	42 (86%)	7 (14%)	3	1
1	L	49/52 (94%)	49 (100%)	0	100	100
2	H	50/52 (96%)	48 (96%)	2 (4%)	28	22
All	All	597/624 (96%)	571 (96%)	26 (4%)	25	19

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	390	LYS
1	I	400	LEU
1	K	372	LEU
1	I	377	GLU
1	J	404	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	I	403	GLN
1	L	403	GLN
1	L	405	ASN
1	C	398	ASN
1	F	369	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PO4	J	501	-	4,4,4	0.82	0	6,6,6	0.73	0
3	PO4	A	501	-	4,4,4	0.91	0	6,6,6	0.53	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	J	501	PO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	50/54 (92%)	-1.14	0 100 100	23, 31, 50, 63	0
1	B	51/54 (94%)	-1.12	0 100 100	20, 30, 50, 71	0
1	C	54/54 (100%)	-1.05	0 100 100	18, 34, 67, 89	0
1	D	53/54 (98%)	-1.17	0 100 100	20, 31, 64, 72	0
1	E	53/54 (98%)	-1.15	0 100 100	25, 36, 54, 67	0
1	F	52/54 (96%)	-1.00	0 100 100	25, 42, 70, 77	0
1	G	52/54 (96%)	-1.12	0 100 100	23, 35, 67, 77	0
1	I	51/54 (94%)	-1.03	0 100 100	25, 39, 61, 76	0
1	J	49/54 (90%)	-1.05	0 100 100	26, 39, 59, 84	0
1	K	50/54 (92%)	-0.92	0 100 100	26, 41, 70, 87	0
1	L	50/54 (92%)	-0.88	0 100 100	32, 44, 69, 92	0
2	H	52/54 (96%)	-1.16	0 100 100	23, 35, 62, 77	0
All	All	617/648 (95%)	-1.07	0 100 100	18, 37, 67, 92	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PO4	A	501	5/5	0.96	0.06	79,83,85,88	0
3	PO4	J	501	5/5	0.98	0.05	56,73,81,82	0
4	K	B	501	1/1	0.99	0.13	76,76,76,76	0
4	K	D	501	1/1	0.99	0.03	52,52,52,52	0

6.5 Other polymers [i](#)

There are no such residues in this entry.