



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 04:53 AM UTC

PDB ID : 1LVU / pdb_00001lvu
Title : Crystal structure of calf spleen purine nucleoside phosphorylase in a new space group with full trimer in the asymmetric unit
Authors : Bzowska, A.; Koellner, G.; Wielgus-Kutrowska, B.; Stroh, A.; Raszewski, G.; Holy, A.; Steiner, T.; Frank, J.
Deposited on : 2002-05-29
Resolution : 2.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtrriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	289	
1	B	289	
1	C	289	
1	D	289	
1	E	289	
1	F	289	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 14809 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Purine nucleoside phosphorylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	272	Total	C	N	O	S	0	0	0
			2131	1352	374	390	15			
1	B	277	Total	C	N	O	S	0	0	0
			2174	1379	381	399	15			
1	C	273	Total	C	N	O	S	0	0	0
			2140	1357	376	391	16			
1	D	279	Total	C	N	O	S	0	0	0
			2189	1387	384	403	15			
1	E	268	Total	C	N	O	S	0	0	0
			2104	1337	370	382	15			
1	F	269	Total	C	N	O	S	0	0	0
			2112	1341	370	385	16			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	144	GLN	GLU	conflict	UNP P55859
B	1144	GLN	GLU	conflict	UNP P55859
C	2144	GLN	GLU	conflict	UNP P55859
D	144	GLN	GLU	conflict	UNP P55859
E	1144	GLN	GLU	conflict	UNP P55859
F	2144	GLN	GLU	conflict	UNP P55859

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Ca	0	0
			2	2		
2	B	3	Total	Ca	0	0
			3	3		
2	C	1	Total	Ca	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	1	Total	Ca	0	0
			1	1		
2	E	1	Total	Ca	0	0
			1	1		
2	F	2	Total	Ca	0	0
			2	2		

- Molecule 3 is 2,6-DIAMINO-(S)-9-[2-(PHOSPHONOMETHOXY)PROPYL]PURINE (CCD ID: 9PP) (formula: C₉H₁₃N₆O₄P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			20	9	6	4	1		
3	B	1	Total	C	N	O	P	0	0
			20	9	6	4	1		
3	C	1	Total	C	N	O	P	0	0
			20	9	6	4	1		
3	D	1	Total	C	N	O	P	0	0
			20	9	6	4	1		
3	E	1	Total	C	N	O	P	0	0
			20	9	6	4	1		
3	F	1	Total	C	N	O	P	0	0
			20	9	6	4	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	251	Total 251	O 251	0	0
4	B	331	Total 331	O 331	0	0
4	C	272	Total 272	O 272	0	0
4	D	350	Total 350	O 350	0	0
4	E	315	Total 315	O 315	0	0
4	F	310	Total 310	O 310	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: Purine nucleoside phosphorylase




• Molecule 1: Purine nucleoside phosphorylase




• Molecule 1: Purine nucleoside phosphorylase







• Molecule 1: Purine nucleoside phosphorylase



• Molecule 1: Purine nucleoside phosphorylase



• Molecule 1: Purine nucleoside phosphorylase



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	79.01Å 132.62Å 177.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.05	Depositor
% Data completeness (in resolution range)	98.7 (20.00-2.05)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.188 , 0.244	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	14809	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 9PP, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	1/2179 (0.0%)	0.94	8/2948 (0.3%)
1	B	0.39	0/2223	0.93	11/3008 (0.4%)
1	C	0.46	3/2188 (0.1%)	1.25	13/2959 (0.4%)
1	D	0.40	0/2238	0.96	11/3028 (0.4%)
1	E	0.39	0/2152	0.93	10/2912 (0.3%)
1	F	0.39	0/2160	0.91	8/2923 (0.3%)
All	All	0.41	4/13140 (0.0%)	0.99	61/17778 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	4	GLY	N-CA	9.48	1.59	1.45
1	C	2265	LYS	C-N	8.89	1.46	1.33
1	C	2004	GLY	N-CA	7.78	1.56	1.45
1	C	2266	GLN	C-N	5.96	1.42	1.33

The worst 5 of 61 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	2266	GLN	N-CA-C	-26.45	54.47	110.80
1	C	2265	LYS	O-C-N	-22.41	93.99	122.00
1	C	2265	LYS	CA-C-N	-18.13	86.91	121.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	2265	LYS	C-N-CA	-18.13	86.91	121.54
1	C	2266	GLN	N-CA-CB	13.30	132.97	110.49

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	2265	LYS	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2131	0	2091	95	0
1	B	2174	0	2130	64	0
1	C	2140	0	2102	79	0
1	D	2189	0	2143	70	0
1	E	2104	0	2071	90	0
1	F	2112	0	2073	83	0
2	A	2	0	0	0	0
2	B	3	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	2	0	0	0	0
3	A	20	0	13	2	0
3	B	20	0	13	1	0
3	C	20	0	13	0	0
3	D	20	0	13	2	0
3	E	20	0	13	2	0
3	F	20	0	13	2	0
4	A	251	0	0	7	0
4	B	331	0	0	12	0
4	C	272	0	0	11	0
4	D	350	0	0	12	0
4	E	315	0	0	20	0
4	F	310	0	0	9	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	14809	0	12688	473	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 473 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:2265:LYS:HB3	1:C:2266:GLN:OE1	1.30	1.29
1:C:2265:LYS:CB	1:C:2266:GLN:OE1	1.86	1.23
1:A:27:VAL:HG13	1:A:78:CYS:HB3	1.35	1.06
1:E:1206:CYS:HB3	4:E:8161:HOH:O	1.57	1.05
1:D:39:VAL:HA	1:D:80:MET:HE1	1.42	1.01

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	268/289 (93%)	256 (96%)	11 (4%)	1 (0%)	30	22
1	B	273/289 (94%)	266 (97%)	7 (3%)	0	100	100
1	C	269/289 (93%)	258 (96%)	9 (3%)	2 (1%)	18	10
1	D	275/289 (95%)	267 (97%)	8 (3%)	0	100	100
1	E	264/289 (91%)	258 (98%)	6 (2%)	0	100	100
1	F	265/289 (92%)	257 (97%)	8 (3%)	0	100	100
All	All	1614/1734 (93%)	1562 (97%)	49 (3%)	3 (0%)	43	37

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	4	GLY
1	C	2266	GLN
1	C	2004	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	226/239 (95%)	216 (96%)	10 (4%)	25	19
1	B	231/239 (97%)	221 (96%)	10 (4%)	26	20
1	C	228/239 (95%)	218 (96%)	10 (4%)	25	19
1	D	233/239 (98%)	220 (94%)	13 (6%)	19	12
1	E	224/239 (94%)	208 (93%)	16 (7%)	13	7
1	F	225/239 (94%)	213 (95%)	12 (5%)	20	13
All	All	1367/1434 (95%)	1296 (95%)	71 (5%)	21	14

5 of 71 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	1266	GLN
1	F	2001	MET
1	F	2187	LEU
1	C	2115	ASN
1	C	2073	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 65 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	2086	HIS
1	F	2123	ASN
1	C	2115	ASN
1	C	2086	HIS
1	F	2137	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 16 ligands modelled in this entry, 10 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	9PP	C	6300	-	20,21,21	3.63	8 (40%)	27,31,31	2.12	5 (18%)
3	9PP	E	6500	-	20,21,21	3.56	9 (45%)	27,31,31	2.13	5 (18%)
3	9PP	F	6600	-	20,21,21	3.66	8 (40%)	27,31,31	2.12	5 (18%)
3	9PP	D	6400	-	20,21,21	3.50	8 (40%)	27,31,31	2.11	6 (22%)
3	9PP	A	6100	-	20,21,21	3.68	9 (45%)	27,31,31	2.12	6 (22%)
3	9PP	B	6200	-	20,21,21	3.61	9 (45%)	27,31,31	2.08	5 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	9PP	C	6300	-	-	1/10/10/10	0/2/2/2
3	9PP	E	6500	-	-	2/10/10/10	0/2/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	9PP	F	6600	-	-	2/10/10/10	0/2/2/2
3	9PP	D	6400	-	-	1/10/10/10	0/2/2/2
3	9PP	A	6100	-	-	2/10/10/10	0/2/2/2
3	9PP	B	6200	-	-	3/10/10/10	0/2/2/2

The worst 5 of 51 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	6300	9PP	P-C14	-11.09	1.52	1.80
3	F	6600	9PP	P-C14	-11.01	1.53	1.80
3	A	6100	9PP	P-C14	-10.96	1.53	1.80
3	B	6200	9PP	P-C14	-10.86	1.53	1.80
3	E	6500	9PP	P-C14	-10.62	1.53	1.80

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	6300	9PP	C2-N1-C6	6.13	124.64	117.28
3	D	6400	9PP	C2-N1-C6	6.13	124.64	117.28
3	F	6600	9PP	C2-N1-C6	6.05	124.53	117.28
3	E	6500	9PP	C2-N1-C6	6.00	124.47	117.28
3	A	6100	9PP	C2-N1-C6	5.94	124.41	117.28

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	6100	9PP	N9-C10-C11-O13
3	B	6200	9PP	N9-C10-C11-O13
3	C	6300	9PP	N9-C10-C11-O13
3	D	6400	9PP	N9-C10-C11-O13
3	E	6500	9PP	N9-C10-C11-O13

There are no ring outliers.

5 monomers are involved in 9 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	6500	9PP	2	0
3	F	6600	9PP	2	0
3	D	6400	9PP	2	0



Continued on next page...



Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	6100	9PP	2	0
3	B	6200	9PP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.