



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 5, 2026 – 01:18 PM UTC

PDB ID : 3LVA / pdb_00003lva
Title : Crystal structure of colorless GFP-like protein from Aequorea coerulescens
Authors : Pletneva, N.V.; Pletnev, V.Z.; Pletnev, S.V.
Deposited on : 2010-02-19
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	236	
1	B	236	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8001 atoms, of which 3711 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Green fluorescent protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	235	Total	C	H	N	O	S	0	6	0
			3773	1222	1847	327	369	8			
1	B	235	Total	C	H	N	O	S	0	6	0
			3748	1216	1832	321	371	8			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	11	ILE	VAL	engineered mutation	UNP Q6YGZ0
A	64	LEU	TYR	engineered mutation	UNP Q6YGZ0
A	101	GLU	LYS	engineered mutation	UNP Q6YGZ0
A	206	ALA	THR	engineered mutation	UNP Q6YGZ0
A	222	GLY	GLU	engineered mutation	UNP Q6YGZ0
B	11	ILE	VAL	engineered mutation	UNP Q6YGZ0
B	64	LEU	TYR	engineered mutation	UNP Q6YGZ0
B	101	GLU	LYS	engineered mutation	UNP Q6YGZ0
B	206	ALA	THR	engineered mutation	UNP Q6YGZ0
B	222	GLY	GLU	engineered mutation	UNP Q6YGZ0

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			14	3	8	3		
2	A	1	Total	C	H	O	0	0
			14	3	8	3		
2	A	1	Total	C	H	O	0	0
			14	3	8	3		
2	B	1	Total	C	H	O	0	0
			14	3	8	3		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	211	Total	O	0	0
			211	211		
4	B	208	Total	O	0	0
			208	208		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Green fluorescent protein



- Molecule 1: Green fluorescent protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	74.50Å 74.50Å 325.92Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.09 – 1.50 29.09 – 1.50	Depositor EDS
% Data completeness (in resolution range)	96.5 (29.09-1.50) 99.2 (29.09-1.50)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtrriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.144 , 0.201 0.154 , 0.194	Depositor DCC
R_{free} test set	1724 reflections (2.00%)	wwPDB-VP
Wilson B-factor (Å ²)	(Not available)	Xtrriage
Anisotropy	(Not available)	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 42.2	EDS
L-test for twinning ¹	$\langle L \rangle =$ (Not available), $\langle L^2 \rangle =$ (Not available)	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8001	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *(Not available)*

¹Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, GYS, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/1954	0.69	0/2637
1	B	0.48	0/1947	0.67	0/2631
All	All	0.48	0/3901	0.68	0/5268

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1926	1847	1860	31	1
1	B	1916	1832	1846	27	1
2	A	18	24	24	4	0
2	B	6	8	8	1	0
3	B	5	0	0	1	0
4	A	211	0	0	2	0
4	B	208	0	0	2	0
All	All	4290	3711	3738	52	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:239:SO4:O4	4:B:419:HOH:O	1.80	0.96
1:A:80:GLN:HE21	1:A:80:GLN:H	1.24	0.86
1:B:151:TYR:CD2	1:B:233:MET:HG2	2.14	0.81
1:A:93:ILE:HD13	1:A:111:GLU:HG2	1.63	0.80
1:A:164:ASN:HB2	1:A:233:MET:HE1	1.71	0.73
1:A:214:LYS:N	1:A:214:LYS:HD3	2.03	0.73
1:A:93:ILE:CD1	1:A:111:GLU:HG2	2.21	0.70
1:A:182:TYR:HB3	2:A:240:GOL:H2	1.75	0.69
1:B:213:GLU:OE2	1:B:215:ARG:HD3	1.94	0.68
1:A:213:GLU:OE2	1:A:215:ARG:HD3	1.95	0.67
1:A:214:LYS:HD3	1:A:214:LYS:H	1.60	0.65
1:A:231:HIS:HE1	1:B:142:GLU:OE2	1.79	0.65
1:B:214:LYS:H	1:B:214:LYS:HE2	1.63	0.63
1:B:231:HIS:HD2	1:B:232:GLY:O	1.81	0.63
1:A:231:HIS:HD2	1:A:232:GLY:O	1.82	0.61
1:A:214:LYS:H	1:A:214:LYS:CD	2.14	0.60
1:A:142:GLU:OE2	1:B:231:HIS:HE1	1.84	0.59
1:A:204:GLN:HE22	1:B:207:LEU:H	1.50	0.58
1:A:23:ASN:HD21	1:A:130:PHE:H	1.52	0.57
1:B:81:HIS:CD2	1:B:197:ASP:H	2.23	0.56
1:A:122[A]:ARG:NH1	1:A:122[A]:ARG:HB3	2.20	0.56
1:A:207:LEU:H	1:B:204:GLN:HE22	1.52	0.56
1:A:81:HIS:CD2	1:A:197:ASP:H	2.26	0.54
1:A:162:LYS:CD	2:A:240:GOL:H12	2.38	0.53
1:B:214:LYS:H	1:B:214:LYS:CD	2.21	0.53
1:A:169:HIS:HD2	4:A:248:HOH:O	1.93	0.51
1:A:213:GLU:OE2	1:A:214:LYS:HE2	2.11	0.51
1:B:214:LYS:H	1:B:214:LYS:CE	2.23	0.51
1:B:76:ASP:H	2:B:240:GOL:H11	1.76	0.50
1:A:207:LEU:H	1:B:204:GLN:NE2	2.10	0.50
1:B:203[B]:THR:HG22	1:B:224:VAL:HG13	1.94	0.48
1:A:135:ASN:HB2	1:A:177:GLN:HE22	1.78	0.48
1:B:169:HIS:HD2	4:B:259:HOH:O	1.96	0.48
1:A:162:LYS:HD3	2:A:240:GOL:H12	1.96	0.47
1:A:143:TYR:HB3	2:A:239:GOL:H11	1.96	0.47
1:B:166:LYS:HE3	1:B:234:ASP:OD1	2.14	0.46
1:B:135:ASN:HB2	1:B:177:GLN:HE22	1.81	0.46
1:A:238:LYS:HE3	1:B:173:ASP:OD2	2.15	0.45
1:B:93:ILE:CD1	1:B:111:GLU:HG2	2.46	0.45
1:A:204:GLN:NE2	1:B:207:LEU:H	2.15	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:203[A]:THR:HG23	1:B:224:VAL:HG22	1.97	0.45
1:B:163:VAL:HB	1:B:183:GLN:HB3	1.97	0.44
1:B:23:ASN:HD21	1:B:130:PHE:H	1.65	0.44
1:A:203:THR:HG23	1:A:224:VAL:HG22	1.99	0.44
1:A:93:ILE:O	1:A:185:ASN:HA	2.18	0.43
1:A:164:ASN:CB	1:A:233:MET:HE1	2.47	0.43
1:B:18:LEU:C	1:B:18:LEU:HD23	2.44	0.43
1:B:52:LYS:HD2	1:B:216:ASP:OD2	2.19	0.43
1:B:213:GLU:HA	1:B:214:LYS:HE2	2.02	0.41
1:A:79:LYS:HE3	4:A:247:HOH:O	2.19	0.41
1:B:93:ILE:O	1:B:185:ASN:HA	2.21	0.40
1:A:122[A]:ARG:HB3	1:A:122[A]:ARG:HH11	1.84	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:198:ASN:HD22	1:B:233:MET:H[8_445]	1.21	0.39

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	236/236 (100%)	232 (98%)	4 (2%)	0	100	100
1	B	236/236 (100%)	233 (99%)	3 (1%)	0	100	100
All	All	472/472 (100%)	465 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	208/203 (102%)	201 (97%)	7 (3%)	32 7
1	B	208/203 (102%)	205 (99%)	3 (1%)	59 33
All	All	416/406 (102%)	406 (98%)	10 (2%)	42 15

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	LYS
1	A	80	GLN
1	A	123	ILE
1	A	166	LYS
1	A	190	ASP
1	A	214	LYS
1	A	215	ARG
1	B	3	LYS
1	B	15	LEU
1	B	214	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	23	ASN
1	A	25	HIS
1	A	80	GLN
1	A	121	ASN
1	A	135	ASN
1	A	149	ASN
1	A	169	HIS
1	A	170	ASN
1	A	177	GLN
1	A	198	ASN
1	A	204	GLN
1	A	231	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	23	ASN
1	B	81	HIS
1	B	121	ASN
1	B	135	ASN
1	B	149	ASN
1	B	169	HIS
1	B	170	ASN
1	B	177	GLN
1	B	204	GLN
1	B	231	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	GYS	A	65	1	20,22,23	0.85	0	25,30,32	1.23	2 (8%)
1	GYS	B	65	1	20,22,23	0.96	0	25,30,32	1.31	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	GYS	A	65	1	-	1/10/29/30	0/2/2/2
1	GYS	B	65	1	-	1/10/29/30	0/2/2/2

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	65	GYS	N3-C1-N2	3.85	114.51	111.48
1	A	65	GYS	CA1-C1-N3	-3.09	120.82	124.84
1	B	65	GYS	CA1-C1-N3	-3.01	120.91	124.84
1	B	65	GYS	C2-N3-C1	-2.18	107.06	108.07
1	A	65	GYS	N3-C1-N2	2.08	113.12	111.48

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	65	GYS	N1-CA1-CB1-OG1
1	B	65	GYS	N1-CA1-CB1-OG1

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	240	-	5,5,5	0.25	0	5,5,5	0.46	0
2	GOL	B	240	-	5,5,5	0.30	0	5,5,5	0.45	0
3	SO4	B	239	-	4,4,4	0.22	0	6,6,6	0.09	0
2	GOL	A	239	-	5,5,5	0.35	0	5,5,5	0.15	0
2	GOL	A	241	-	5,5,5	0.32	0	5,5,5	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	240	-	-	2/4/4/4	-
2	GOL	A	241	-	-	4/4/4/4	-
2	GOL	A	239	-	-	2/4/4/4	-
2	GOL	A	240	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	240	GOL	O1-C1-C2-O2
2	A	240	GOL	O1-C1-C2-C3
2	A	241	GOL	O1-C1-C2-C3
2	A	239	GOL	C1-C2-C3-O3
2	A	241	GOL	C1-C2-C3-O3
2	B	240	GOL	O1-C1-C2-C3
2	B	240	GOL	O1-C1-C2-O2
2	A	241	GOL	O1-C1-C2-O2
2	A	241	GOL	O2-C2-C3-O3
2	A	239	GOL	O2-C2-C3-O3

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	240	GOL	3	0
2	B	240	GOL	1	0
3	B	239	SO4	1	0
2	A	239	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	234/236 (99%)	-0.55	1 (0%) 88 90	12, 26, 45, 58	6 (2%)
1	B	234/236 (99%)	-0.58	0 100 100	12, 26, 42, 56	6 (2%)
All	All	468/472 (99%)	-0.56	1 (0%) 91 92	12, 26, 44, 58	12 (2%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	117	ASP	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	GYS	A	65	21/22	0.98	0.04	15,17,21,21	0
1	GYS	B	65	21/22	0.99	0.04	15,18,21,23	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	A	241	6/6	0.66	0.18	35,53,63,64	0
2	GOL	A	240	6/6	0.69	0.21	41,50,60,63	0
2	GOL	B	240	6/6	0.79	0.15	44,53,63,64	0
2	GOL	A	239	6/6	0.80	0.18	25,39,50,50	14
3	SO4	B	239	5/5	0.90	0.09	52,53,53,54	0

6.5 Other polymers [i](#)

There are no such residues in this entry.