



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 09:18 AM UTC

PDB ID : 4LV2 / pdb_00004lv2
Title : AmpC beta-lactamase in complex with [1-(6-chloropyrimidin-4-yl)-1H-pyrazo
l-4-yl] boronic acid
Authors : London, N.; Eidam, O.; Shoichet, B.K.
Deposited on : 2013-07-25
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.66-1.66)
Clashscore	190562	2662 (1.66-1.66)
Ramachandran outliers	187476	2621 (1.66-1.66)
Sidechain outliers	187428	2621 (1.66-1.66)
RSRZ outliers	180081	2564 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	358	91% 9%
1	B	358	2% 92% 8%

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 6027 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-lactamase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	358	Total	C	N	O	S	0	4	0
			2766	1792	460	508	6			
1	B	358	Total	C	N	O	S	0	7	0
			2772	1799	461	505	7			

- Molecule 2 is [1-(6-chloropyrimidin-4-yl)-1H-pyrazol-4-yl]boronic acid (CCD ID: N95) (formula: C₇H₆BClN₄O₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	B	C	Cl	N	O	0
			15	1	7	1	4	2	
2	B	1	Total	B	C	Cl	N	O	0
			15	1	7	1	4	2	
2	B	1	Total	B	C	Cl	N	O	0
			15	1	7	1	4	2	

- Molecule 3 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	240	Total	O	0	6
			245	245		
4	B	188	Total	O	0	5
			189	189		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-lactamase



- Molecule 1: Beta-lactamase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	117.67Å 78.14Å 97.25Å 90.00° 116.28° 90.00°	Depositor
Resolution (Å)	45.58 – 1.65 45.58 – 1.65	Depositor EDS
% Data completeness (in resolution range)	99.6 (45.58-1.65) 99.6 (45.58-1.65)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.82 (at 1.65Å)	Xtriage
Refinement program	PHENIX 1.8_1069	Depositor
R, R_{free}	0.172 , 0.193 (Not available) , 0.205	Depositor DCC
R_{free} test set	3424 reflections (3.62%)	wwPDB-VP
Wilson B-factor (Å ²)	20.9	Xtriage
Anisotropy	0.209	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 40.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6027	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, N95

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.79	0/2858	0.92	0/3916
1	B	0.71	0/2873	0.86	4/3936 (0.1%)
All	All	0.75	0/5731	0.89	4/7852 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	346	ASN	CA-C-N	-5.33	113.42	119.28
1	B	346	ASN	C-N-CA	-5.33	113.42	119.28
1	B	164	LYS	CA-C-N	-5.06	114.45	119.56
1	B	164	LYS	C-N-CA	-5.06	114.45	119.56

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2766	0	2675	20	0
1	B	2772	0	2694	18	0
2	A	15	0	6	0	0
2	B	30	0	12	2	0
3	A	5	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	5	0	0	0	0
4	A	245	0	0	3	0
4	B	189	0	0	0	0
All	All	6027	0	5387	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:175[B]:GLN:OE1	1:A:180:GLN:NE2	2.09	0.85
1:A:87:ASP:OD2	1:A:92:TYR:OH	2.02	0.77
1:A:309:ARG:NH1	4:A:697:HOH:O	2.21	0.68
1:B:240:PRO:HA	1:B:243:ILE:HD12	1.85	0.58
1:A:258:ARG:NH1	4:A:707:HOH:O	2.40	0.55
1:A:119:LEU:HA	1:A:151:ALA:HA	1.92	0.52
1:B:344:TYR:CE1	1:B:349:ARG:HG2	2.45	0.52
1:A:276:TRP:CD2	1:A:277:PRO:HA	2.46	0.51
1:A:344:TYR:CZ	1:A:349:ARG:HG2	2.47	0.49
1:A:243:ILE:CD1	1:A:252:ILE:HD12	2.44	0.47
1:A:125:VAL:HG11	1:A:131:LEU:HD23	1.95	0.47
1:A:280:PRO:HG3	1:A:354:TRP:CZ2	2.49	0.47
1:B:326[B]:VAL:HG12	1:B:337:VAL:HG13	1.96	0.47
1:B:87:ASP:OD2	1:B:92:TYR:OH	2.31	0.46
1:B:92:TYR:CE1	1:B:106:LEU:HD11	2.51	0.45
1:A:93:TRP:CD1	1:A:95:GLU:HG2	2.51	0.45
1:B:56[B]:GLN:OE1	2:B:403:N95:CL1	2.71	0.45
1:B:243:ILE:HD13	1:B:252:ILE:HD12	1.98	0.45
1:A:264:ASP:OD1	1:A:282:SER:OG	2.33	0.44
1:A:334:LEU:HG	1:A:357:LEU:HD22	1.99	0.44
1:B:59[A]:LEU:HB2	1:B:199:TYR:HA	1.98	0.44
1:A:11:ILE:HB	4:A:695:HOH:O	2.18	0.44
1:A:344:TYR:CE1	1:A:349:ARG:HG2	2.54	0.43
1:B:26:PRO:HB3	1:B:48:ILE:HD11	2.01	0.43
1:B:119:LEU:HA	1:B:151:ALA:HA	2.00	0.43
1:B:344:TYR:CZ	1:B:349:ARG:HG2	2.55	0.42
1:B:245:GLU:OE1	1:B:245:GLU:N	2.50	0.42
1:A:189[A]:ILE:HD13	1:A:223:VAL:HG22	2.02	0.42
1:A:164[B]:LYS:NZ	3:A:402:PO4:O3	2.43	0.41
1:B:41:PHE:HA	2:B:403:N95:C11	2.51	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:262[B]:THR:CG2	1:B:298:VAL:HG12	2.51	0.41
1:B:240:PRO:O	1:B:249:GLN:HG3	2.20	0.41
1:A:250:GLN:HA	1:A:253:GLN:HE21	1.85	0.41
1:B:276:TRP:CD2	1:B:277:PRO:HA	2.55	0.41
1:A:107:LEU:HD22	1:B:303:PRO:HD3	2.03	0.40
1:B:351:ASP:O	1:B:355:GLN:HG2	2.21	0.40
1:A:130:ASP:OD1	1:A:133:ARG:NH2	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	360/358 (101%)	353 (98%)	7 (2%)	0	100	100
1	B	363/358 (101%)	356 (98%)	7 (2%)	0	100	100
All	All	723/716 (101%)	709 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	150/292 (51%)	148 (99%)	2 (1%)	61	42

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	262[A]	THR
1	B	262[B]	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	253	GLN
1	B	346	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	PO4	B	402	-	4,4,4	0.72	0	6,6,6	0.78	0
2	N95	B	403	-	16,16,16	3.05	5 (31%)	20,22,22	3.86	10 (50%)
3	PO4	A	402	-	4,4,4	0.93	0	6,6,6	0.43	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	N95	A	401	1	16,16,16	2.91	3 (18%)	20,22,22	4.57	14 (70%)
2	N95	B	401	1	16,16,16	2.62	4 (25%)	20,22,22	4.67	14 (70%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	N95	B	401	1	-	0/6/8/8	0/2/2/2
2	N95	A	401	1	-	0/6/8/8	0/2/2/2
2	N95	B	403	-	-	0/6/8/8	0/2/2/2

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	N95	B03-O05	8.17	1.48	1.36
2	B	403	N95	B03-C06	7.66	1.64	1.57
2	A	401	N95	B03-O04	7.35	1.47	1.36
2	B	401	N95	B03-O04	6.58	1.45	1.36
2	B	401	N95	B03-O05	6.58	1.45	1.36
2	B	403	N95	B03-O05	5.52	1.44	1.36
2	B	403	N95	C07-C06	-4.45	1.38	1.41
2	B	403	N95	C10-N09	-4.10	1.34	1.42
2	B	403	N95	B03-O04	3.66	1.41	1.36
2	A	401	N95	C10-N09	-2.98	1.36	1.42
2	B	401	N95	C10-N09	-2.93	1.36	1.42
2	B	401	N95	C07-C06	-2.08	1.40	1.41

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	N95	C15-N14-C12	10.34	121.61	113.85
2	B	403	N95	C15-N14-C12	9.08	120.67	113.85
2	B	401	N95	C15-N16-C10	8.65	122.21	114.95
2	A	401	N95	C07-N08-N09	8.54	110.99	104.19
2	A	401	N95	C15-N16-C10	8.33	121.94	114.95
2	A	401	N95	C15-N14-C12	7.64	119.58	113.85
2	B	403	N95	C11-C12-N14	-7.54	119.03	125.59
2	B	401	N95	C07-N08-N09	6.72	109.54	104.19
2	B	403	N95	C12-C11-C10	6.62	118.83	115.02
2	B	401	N95	N16-C15-N14	-6.48	118.77	128.58

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	N95	C11-C12-N14	-6.22	120.18	125.59
2	A	401	N95	C12-C11-C10	6.18	118.58	115.02
2	B	403	N95	C15-N16-C10	5.46	119.53	114.95
2	A	401	N95	C11-C12-N14	-5.41	120.88	125.59
2	A	401	N95	C17-N09-N08	-5.08	106.69	112.14
2	A	401	N95	N16-C15-N14	-5.05	120.93	128.58
2	B	403	N95	N16-C15-N14	-4.72	121.44	128.58
2	B	401	N95	C12-C11-C10	4.44	117.57	115.02
2	B	401	N95	C10-N09-N08	4.28	127.76	120.71
2	A	401	N95	C06-C17-N09	4.14	109.71	106.65
2	A	401	N95	O05-B03-O04	-4.11	108.54	120.77
2	A	401	N95	O05-B03-C06	-4.11	108.41	120.19
2	B	401	N95	C17-N09-N08	-4.03	107.82	112.14
2	B	401	N95	O05-B03-O04	-4.01	108.84	120.77
2	B	403	N95	C07-N08-N09	3.83	107.24	104.19
2	A	401	N95	C10-N09-N08	3.75	126.90	120.71
2	B	403	N95	N16-C10-N09	3.70	120.36	114.84
2	B	401	N95	O05-B03-C06	-3.62	109.82	120.19
2	B	401	N95	O04-B03-C06	-3.47	110.23	120.19
2	B	401	N95	N16-C10-N09	3.35	119.83	114.84
2	A	401	N95	O04-B03-C06	-3.02	111.53	120.19
2	B	403	N95	CL1-C12-N14	2.99	121.59	115.98
2	A	401	N95	N16-C10-N09	2.95	119.24	114.84
2	B	403	N95	C10-N09-N08	2.88	125.45	120.71
2	A	401	N95	C11-C10-N16	-2.87	118.05	122.82
2	B	401	N95	CL1-C12-N14	2.68	121.00	115.98
2	B	403	N95	C17-N09-N08	-2.55	109.40	112.14
2	B	401	N95	C06-C17-N09	2.48	108.48	106.65

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	403	N95	2	0
3	A	402	PO4	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	358/358 (100%)	-0.20	2 (0%) 85 90	10, 21, 34, 47	4 (1%)
1	B	358/358 (100%)	-0.01	6 (1%) 69 74	10, 24, 37, 46	7 (1%)
All	All	716/716 (100%)	-0.10	8 (1%) 78 83	10, 23, 35, 47	11 (1%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	4	ALA	3.7
1	A	4	ALA	3.0
1	A	264	ASP	2.7
1	B	5	PRO	2.4
1	B	244	ASN	2.3
1	B	245	GLU	2.2
1	B	242	ASP	2.1
1	B	128	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PO4	B	402	5/5	0.80	0.10	43,45,51,57	0
3	PO4	A	402	5/5	0.90	0.09	42,45,52,57	0
2	N95	B	403	15/15	0.90	0.10	22,27,35,43	0
2	N95	B	401	15/15	0.94	0.07	17,23,31,45	0
2	N95	A	401	15/15	0.95	0.06	14,19,25,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.