



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 14, 2026 – 03:19 PM UTC

PDB ID : 4LVH / pdb_00004lvh
Title : Insight into highly conserved H1 subtype-specific epitopes in influenza virus hemagglutinin
Authors : Kim, K.H.; Cho, K.J.; Kim, S.; Seok, J.H.; Lee, J.-H.
Deposited on : 2013-07-26
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	518	3% 15%22%8%54%
1	D	518	2% 15%20%7%57%
1	G	518	0% 15%21%6%57%
1	J	518	2% 15%18%9%57%
2	B	222	3% 31%42%21%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	E	222	5% 35% 39% 20% 5%
2	H	222	5% 29% 42% 23% 5%
2	K	222	7% 29% 46% 20%
3	C	211	5% 26% 46% 20% 7%
3	F	211	2% 27% 41% 26% 5%
3	I	211	6% 29% 49% 18%
3	L	211	5% 30% 39% 26%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 20106 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	236	Total	C	N	O	S	0	1	0
			1831	1162	311	352	6			
1	D	222	Total	C	N	O	S	0	1	0
			1762	1121	297	338	6			
1	G	222	Total	C	N	O	S	0	1	0
			1762	1121	297	338	6			
1	J	222	Total	C	N	O	S	0	1	0
			1762	1121	297	338	6			

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-8	ALA	-	expression tag	UNP C5MQE6
A	-7	ASP	-	expression tag	UNP C5MQE6
A	-6	PRO	-	expression tag	UNP C5MQE6
A	-5	GLY	-	expression tag	UNP C5MQE6
A	-4	TYR	-	expression tag	UNP C5MQE6
A	-3	LEU	-	expression tag	UNP C5MQE6
A	-2	LEU	-	expression tag	UNP C5MQE6
A	-1	GLU	-	expression tag	UNP C5MQE6
A	0	PHE	-	expression tag	UNP C5MQE6
A	507	ARG	-	expression tag	UNP C5MQE6
A	508	SER	-	expression tag	UNP C5MQE6
A	509	LEU	-	expression tag	UNP C5MQE6
A	510	VAL	-	expression tag	UNP C5MQE6
A	511	PRO	-	expression tag	UNP C5MQE6
A	512	ARG	-	expression tag	UNP C5MQE6
D	-8	ALA	-	expression tag	UNP C5MQE6
D	-7	ASP	-	expression tag	UNP C5MQE6
D	-6	PRO	-	expression tag	UNP C5MQE6
D	-5	GLY	-	expression tag	UNP C5MQE6
D	-4	TYR	-	expression tag	UNP C5MQE6
D	-3	LEU	-	expression tag	UNP C5MQE6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-2	LEU	-	expression tag	UNP C5MQE6
D	-1	GLU	-	expression tag	UNP C5MQE6
D	0	PHE	-	expression tag	UNP C5MQE6
D	504	ARG	-	expression tag	UNP C5MQE6
D	505	SER	-	expression tag	UNP C5MQE6
D	506	LEU	-	expression tag	UNP C5MQE6
D	507	VAL	-	expression tag	UNP C5MQE6
D	508	PRO	-	expression tag	UNP C5MQE6
D	509	ARG	-	expression tag	UNP C5MQE6
G	-8	ALA	-	expression tag	UNP C5MQE6
G	-7	ASP	-	expression tag	UNP C5MQE6
G	-6	PRO	-	expression tag	UNP C5MQE6
G	-5	GLY	-	expression tag	UNP C5MQE6
G	-4	TYR	-	expression tag	UNP C5MQE6
G	-3	LEU	-	expression tag	UNP C5MQE6
G	-2	LEU	-	expression tag	UNP C5MQE6
G	-1	GLU	-	expression tag	UNP C5MQE6
G	0	PHE	-	expression tag	UNP C5MQE6
G	504	ARG	-	expression tag	UNP C5MQE6
G	505	SER	-	expression tag	UNP C5MQE6
G	506	LEU	-	expression tag	UNP C5MQE6
G	507	VAL	-	expression tag	UNP C5MQE6
G	508	PRO	-	expression tag	UNP C5MQE6
G	509	ARG	-	expression tag	UNP C5MQE6
J	-8	ALA	-	expression tag	UNP C5MQE6
J	-7	ASP	-	expression tag	UNP C5MQE6
J	-6	PRO	-	expression tag	UNP C5MQE6
J	-5	GLY	-	expression tag	UNP C5MQE6
J	-4	TYR	-	expression tag	UNP C5MQE6
J	-3	LEU	-	expression tag	UNP C5MQE6
J	-2	LEU	-	expression tag	UNP C5MQE6
J	-1	GLU	-	expression tag	UNP C5MQE6
J	0	PHE	-	expression tag	UNP C5MQE6
J	504	ARG	-	expression tag	UNP C5MQE6
J	505	SER	-	expression tag	UNP C5MQE6
J	506	LEU	-	expression tag	UNP C5MQE6
J	507	VAL	-	expression tag	UNP C5MQE6
J	508	PRO	-	expression tag	UNP C5MQE6
J	509	ARG	-	expression tag	UNP C5MQE6

- Molecule 2 is a protein called MONOCLONAL ANTIBODY H-CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	218	Total	C	N	O	S	0	0	0
			1635	1028	280	319	8			
2	E	218	Total	C	N	O	S	0	0	0
			1635	1028	280	319	8			
2	H	218	Total	C	N	O	S	0	0	0
			1635	1028	280	319	8			
2	K	218	Total	C	N	O	S	0	0	0
			1635	1028	280	319	8			

- Molecule 3 is a protein called MONOCLONAL ANTIBODY L-CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	210	Total	C	N	O	S	0	0	0
			1610	1007	271	326	6			
3	F	210	Total	C	N	O	S	0	0	0
			1610	1007	271	326	6			
3	I	210	Total	C	N	O	S	0	0	0
			1610	1007	271	326	6			
3	L	209	Total	C	N	O	S	0	0	0
			1604	1004	270	324	6			

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ca	0	0
			1	1		
4	D	1	Total	Ca	0	0
			1	1		
4	G	1	Total	Ca	0	0
			1	1		
4	J	1	Total	Ca	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	O	0	0
			1	1		
5	B	1	Total	O	0	0
			1	1		
5	F	1	Total	O	0	0
			1	1		


Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	G	1	Total 1	O 1	0	0
5	H	3	Total 3	O 3	0	0
5	I	2	Total 2	O 2	0	0
5	K	1	Total 1	O 1	0	0
5	L	1	Total 1	O 1	0	0




• Molecule 1: Hemagglutinin







• Molecule 2: MONOCLONAL ANTIBODY H-CHAIN



• Molecule 2: MONOCLONAL ANTIBODY H-CHAIN




• Molecule 2: MONOCLONAL ANTIBODY H-CHAIN









• Molecule 3: MONOCLONAL ANTIBODY L-CHAIN




• Molecule 3: MONOCLONAL ANTIBODY L-CHAIN




• Molecule 3: MONOCLONAL ANTIBODY L-CHAIN





● Molecule 3: MONOCLONAL ANTIBODY L-CHAIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	72.75Å 237.81Å 94.23Å 90.00° 110.31° 90.00°	Depositor
Resolution (Å)	43.49 – 2.80 43.49 – 2.80	Depositor EDS
% Data completeness (in resolution range)	76.8 (43.49-2.80) 92.6 (43.49-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.05 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.232 , 0.274 0.228 , 0.233	Depositor DCC
R_{free} test set	3559 reflections (4.31%)	wwPDB-VP
Wilson B-factor (Å ²)	36.9	Xtriage
Anisotropy	0.160	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.25 , 38.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.004 for h,-k,-h-l	Xtriage
Reported twinning fraction	0.844 for H, K, L 0.156 for -H, -K, H+L	Depositor
Outliers	0 of 69841 reflections	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	20106	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 22.72 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.3778e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.74	1/1879 (0.1%)	1.37	26/2549 (1.0%)
1	D	0.69	0/1811	1.28	25/2456 (1.0%)
1	G	0.74	0/1811	1.29	26/2456 (1.1%)
1	J	0.67	0/1811	1.30	24/2456 (1.0%)
2	B	0.76	0/1675	1.40	27/2277 (1.2%)
2	E	0.80	1/1675 (0.1%)	1.35	25/2277 (1.1%)
2	H	0.79	0/1675	1.33	24/2277 (1.1%)
2	K	0.75	0/1675	1.37	18/2277 (0.8%)
3	C	0.81	1/1648 (0.1%)	1.57	38/2242 (1.7%)
3	F	0.83	1/1648 (0.1%)	1.58	37/2242 (1.7%)
3	I	0.79	0/1648	1.53	40/2242 (1.8%)
3	L	0.81	1/1641 (0.1%)	1.56	38/2231 (1.7%)
All	All	0.76	5/20597 (0.0%)	1.41	348/27982 (1.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	6
1	D	0	1
1	G	0	1
1	J	0	3
2	B	0	7
2	E	0	2
2	H	0	5
2	K	0	6
3	C	0	7
3	F	0	6
3	I	0	4

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
3	L	0	7
All	All	0	55

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	45	GLN	CA-C	6.52	1.61	1.52
1	A	70	LEU	CA-C	5.78	1.58	1.53
3	L	45	GLN	CA-C	5.46	1.59	1.52
3	F	202	THR	CA-CB	5.03	1.61	1.53
2	E	45	GLU	CA-C	5.01	1.58	1.52

The worst 5 of 348 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	L	97	VAL	CA-C-N	-14.77	104.04	119.99
3	L	97	VAL	C-N-CA	-14.77	104.04	119.99
1	D	235	GLU	CA-C-N	-11.85	109.68	121.65
1	D	235	GLU	C-N-CA	-11.85	109.68	121.65
3	F	97	VAL	CA-C-N	-11.79	107.53	119.90

There are no chirality outliers.

5 of 55 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	156	ASN	Peptide
1	A	203	SER	Peptide
1	A	221	ARG	Sidechain
1	A	65	PRO	Peptide
1	A	74	SER	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1831	0	1725	142	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	1762	0	1694	110	0
1	G	1762	0	1694	114	0
1	J	1762	0	1694	114	0
2	B	1635	0	1587	138	0
2	E	1635	0	1587	153	0
2	H	1635	0	1587	153	0
2	K	1635	0	1587	142	0
3	C	1610	0	1549	173	0
3	F	1610	0	1549	165	0
3	I	1610	0	1549	140	0
3	L	1604	0	1543	127	0
4	A	1	0	0	0	0
4	D	1	0	0	0	0
4	G	1	0	0	0	0
4	J	1	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	F	1	0	0	0	0
5	G	1	0	0	0	0
5	H	3	0	0	0	0
5	I	2	0	0	0	0
5	K	1	0	0	0	0
5	L	1	0	0	0	0
All	All	20106	0	19345	1554	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 39.

The worst 5 of 1554 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:115:GLU:HB3	3:I:96:GLU:HB3	1.37	1.07
2:H:216:ASP:O	2:H:217:LYS:HG3	1.56	1.06
2:K:63:VAL:HA	2:K:66:ARG:NH2	1.71	1.02
2:K:63:VAL:HA	2:K:66:ARG:HH21	1.18	1.02
2:E:39:ALA:HB2	2:E:44:LEU:HB2	1.42	1.01

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	233/518 (45%)	191 (82%)	40 (17%)	2 (1%)	14	41
1	D	221/518 (43%)	183 (83%)	38 (17%)	0	100	100
1	G	221/518 (43%)	189 (86%)	30 (14%)	2 (1%)	14	41
1	J	221/518 (43%)	187 (85%)	32 (14%)	2 (1%)	14	41
2	B	214/222 (96%)	180 (84%)	29 (14%)	5 (2%)	5	18
2	E	214/222 (96%)	176 (82%)	35 (16%)	3 (1%)	9	30
2	H	214/222 (96%)	173 (81%)	35 (16%)	6 (3%)	4	14
2	K	214/222 (96%)	164 (77%)	47 (22%)	3 (1%)	9	30
3	C	208/211 (99%)	147 (71%)	50 (24%)	11 (5%)	1	5
3	F	208/211 (99%)	150 (72%)	47 (23%)	11 (5%)	1	5
3	I	208/211 (99%)	157 (76%)	41 (20%)	10 (5%)	2	6
3	L	205/211 (97%)	152 (74%)	44 (22%)	9 (4%)	2	7
All	All	2581/3804 (68%)	2049 (79%)	468 (18%)	64 (2%)	4	16

5 of 64 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	F	34	ASN
2	K	48	SER
3	L	5	GLN
3	L	45	GLN
3	L	79	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	193/451 (43%)	141 (73%)	52 (27%)	0	1
1	D	193/451 (43%)	145 (75%)	48 (25%)	0	2
1	G	193/451 (43%)	148 (77%)	45 (23%)	1	3
1	J	193/451 (43%)	139 (72%)	54 (28%)	0	1
2	B	180/184 (98%)	124 (69%)	56 (31%)	0	1
2	E	180/184 (98%)	125 (69%)	55 (31%)	0	1
2	H	180/184 (98%)	115 (64%)	65 (36%)	0	0
2	K	180/184 (98%)	129 (72%)	51 (28%)	0	1
3	C	182/183 (100%)	134 (74%)	48 (26%)	0	2
3	F	182/183 (100%)	130 (71%)	52 (29%)	0	1
3	I	182/183 (100%)	133 (73%)	49 (27%)	0	1
3	L	181/183 (99%)	126 (70%)	55 (30%)	0	1
All	All	2219/3272 (68%)	1589 (72%)	630 (28%)	0	1

5 of 630 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	I	199	LYS
2	K	217	LYS
1	J	68	GLU
3	I	195	GLU
1	J	266	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 36 such sidechains are listed below:

Mol	Chain	Res	Type
3	I	198	HIS
3	L	45	GLN
1	J	87	ASN
2	K	12	GLN
1	D	193	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	236/518 (45%)	0.18	14 (5%) 28 21	6, 15, 85, 123	1 (0%)
1	D	222/518 (42%)	0.15	9 (4%) 41 33	8, 18, 41, 108	1 (0%)
1	G	222/518 (42%)	0.11	3 (1%) 73 64	7, 16, 47, 88	1 (0%)
1	J	222/518 (42%)	0.32	9 (4%) 41 33	6, 19, 50, 110	1 (0%)
2	B	218/222 (98%)	0.14	6 (2%) 55 45	8, 16, 30, 47	0
2	E	218/222 (98%)	0.32	11 (5%) 34 26	7, 17, 37, 50	0
2	H	218/222 (98%)	0.29	10 (4%) 37 29	6, 19, 37, 48	0
2	K	218/222 (98%)	0.48	15 (6%) 23 16	10, 23, 40, 64	0
3	C	210/211 (99%)	0.37	10 (4%) 35 28	10, 20, 47, 78	0
3	F	210/211 (99%)	0.33	4 (1%) 66 57	8, 21, 41, 56	0
3	I	210/211 (99%)	0.40	12 (5%) 29 22	11, 23, 49, 69	0
3	L	209/211 (99%)	0.37	10 (4%) 35 28	11, 24, 44, 82	0
All	All	2613/3804 (68%)	0.29	113 (4%) 40 31	6, 19, 44, 123	4 (0%)

The worst 5 of 113 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	427	GLU	6.4
1	D	262	GLY	5.5
1	A	425	ASN	5.5
2	E	107	ALA	5.4
2	K	202	TYR	5.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	CA	A	601	1/1	0.96	0.15	9,9,9,9	0
4	CA	D	601	1/1	0.96	0.10	7,7,7,7	0
4	CA	G	601	1/1	0.99	0.08	10,10,10,10	0
4	CA	J	601	1/1	0.99	0.09	3,3,3,3	0

6.5 Other polymers [i](#)

There are no such residues in this entry.