



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 11:37 AM UTC

PDB ID : 6LV0 / pdb_00006lv0
Title : Crystal structure of the soluble domain of the multiple peptide resistance factor (MprF) from *Rhizobium tropici*
Authors : Jiao, H.Z.; Song, D.F.; Liu, Z.F.
Deposited on : 2020-02-02
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	335	0% 92% 5%
1	B	335	2% 89% 6% 5%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5683 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Low pH-inducible protein LpiA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	319	Total	C	N	O	S	0	0	0
			2478	1590	428	448	12			
1	B	319	Total	C	N	O	S	0	0	0
			2478	1590	428	448	12			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	541	MET	-	initiating methionine	UNP L0LM43
A	863	LYS	-	expression tag	UNP L0LM43
A	864	LEU	-	expression tag	UNP L0LM43
A	865	ALA	-	expression tag	UNP L0LM43
A	866	ALA	-	expression tag	UNP L0LM43
A	867	ALA	-	expression tag	UNP L0LM43
A	868	LEU	-	expression tag	UNP L0LM43
A	869	GLU	-	expression tag	UNP L0LM43
A	870	HIS	-	expression tag	UNP L0LM43
A	871	HIS	-	expression tag	UNP L0LM43
A	872	HIS	-	expression tag	UNP L0LM43
A	873	HIS	-	expression tag	UNP L0LM43
A	874	HIS	-	expression tag	UNP L0LM43
A	875	HIS	-	expression tag	UNP L0LM43
B	541	MET	-	initiating methionine	UNP L0LM43
B	863	LYS	-	expression tag	UNP L0LM43
B	864	LEU	-	expression tag	UNP L0LM43
B	865	ALA	-	expression tag	UNP L0LM43
B	866	ALA	-	expression tag	UNP L0LM43
B	867	ALA	-	expression tag	UNP L0LM43
B	868	LEU	-	expression tag	UNP L0LM43
B	869	GLU	-	expression tag	UNP L0LM43
B	870	HIS	-	expression tag	UNP L0LM43
B	871	HIS	-	expression tag	UNP L0LM43
B	872	HIS	-	expression tag	UNP L0LM43

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	873	HIS	-	expression tag	UNP L0LM43
B	874	HIS	-	expression tag	UNP L0LM43
B	875	HIS	-	expression tag	UNP L0LM43


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	378	Total 378	O 378	0	0
2	B	349	Total 349	O 349	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Low pH-inducible protein LpiA



- Molecule 1: Low pH-inducible protein LpiA



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	83.59Å 95.79Å 211.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.21 – 2.00 35.21 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (35.21-2.00) 99.7 (35.21-2.00)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.15 (at 2.00Å)	Xtriage
Refinement program	PHENIX 1.11.1 _2575	Depositor
R, R_{free}	0.168 , 0.198 0.169 , 0.200	Depositor DCC
R_{free} test set	2921 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	24.2	Xtriage
Anisotropy	0.077	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 42.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5683	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.33	0/2535	0.49	0/3421
1	B	0.33	0/2535	0.51	0/3421
All	All	0.33	0/5070	0.50	0/6842

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2478	0	2458	6	0
1	B	2478	0	2458	13	0
2	A	378	0	0	0	0
2	B	349	0	0	0	0
All	All	5683	0	4916	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:673:ARG:HH11	1:A:673:ARG:HG2	1.59	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:550:GLU:OE2	1:B:553:ARG:NH1	2.29	0.66
1:A:642:LYS:HE3	1:A:837:PRO:HB3	1.82	0.62
1:B:654:PHE:O	1:B:778:LYS:NZ	2.37	0.58
1:A:673:ARG:HG2	1:A:673:ARG:NH1	2.26	0.50
1:B:660:LYS:HE2	1:B:832:HIS:CD2	2.47	0.50
1:B:547:ARG:HH11	1:B:547:ARG:HG3	1.78	0.49
1:B:598:PRO:O	1:B:602:LEU:HD13	2.13	0.48
1:B:725:PRO:HG2	1:B:743:MET:CE	2.44	0.47
1:B:657:LYS:HB3	1:B:665:ARG:CZ	2.45	0.47
1:A:725:PRO:HG2	1:A:743:MET:CE	2.46	0.46
1:A:676:LEU:HD11	1:A:762:PRO:HG2	1.97	0.45
1:A:852:LEU:HD12	1:A:852:LEU:HA	1.74	0.45
1:B:683:PRO:HA	1:B:686:ILE:HD12	1.98	0.45
1:B:639:ARG:HG2	1:B:641:PHE:CE1	2.52	0.44
1:B:725:PRO:HG2	1:B:743:MET:HE1	2.00	0.43
1:B:729:LEU:HB2	1:B:737:ALA:HB3	2.03	0.41
1:B:599:ARG:HA	1:B:602:LEU:HD22	2.03	0.41
1:B:657:LYS:HB3	1:B:665:ARG:NH1	2.36	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	315/335 (94%)	309 (98%)	5 (2%)	1 (0%)	36	35
1	B	315/335 (94%)	306 (97%)	8 (2%)	1 (0%)	36	35
All	All	630/670 (94%)	615 (98%)	13 (2%)	2 (0%)	36	35

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	711	PHE
1	B	711	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	254/268 (95%)	253 (100%)	1 (0%)	84	89
1	B	254/268 (95%)	254 (100%)	0	100	100
All	All	508/536 (95%)	507 (100%)	1 (0%)	87	92

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	568	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	684	GLN
1	A	772	GLN
1	B	585	GLN
1	B	663	ASN
1	B	832	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	319/335 (95%)	-0.19	3 (0%) 81 80	14, 24, 41, 57	0
1	B	319/335 (95%)	-0.14	6 (1%) 66 66	13, 24, 47, 58	0
All	All	638/670 (95%)	-0.17	9 (1%) 73 73	13, 24, 45, 58	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	846	VAL	3.6
1	B	844	GLY	3.0
1	B	543	ASP	2.8
1	B	614	ARG	2.5
1	A	633	CYS	2.4
1	B	607	TRP	2.4
1	A	793	SER	2.2
1	B	734	LYS	2.2
1	B	847	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.