



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 03:12 PM UTC

PDB ID : 3LWL / pdb_00003lwl
Title : Structure of Klenow fragment of Taq polymerase in complex with an abasic site
Authors : Marx, A.; Diederichs, K.; Obeid, S.
Deposited on : 2010-02-24
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1898 (2.26-2.26)
Clashscore	190562	2005 (2.26-2.26)
Ramachandran outliers	187476	1965 (2.26-2.26)
Sidechain outliers	187428	1966 (2.26-2.26)
RSRZ outliers	180081	1898 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	540	5% 74% 23% ..
2	B	12	83% 17%
3	C	16	50% 31% 6% 12%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	GOL	A	6	-	-	X	-
7	GOL	B	8	-	-	X	-

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 5013 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase I, thermostable.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	534	Total	C	N	O	S	0	0	0
			4236	2693	765	765	13			


- Molecule 2 is a DNA chain called DNA (5'-D(*GP*AP*CP*CP*AP*CP*GP*GP*CP*GP*CP*(2DA))-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	12	Total	C	N	O	P	0	0	0
			242	115	50	66	11			

- Molecule 3 is a DNA chain called DNA (5'-D(*AP*AP*AP*(3DR)P*TP*GP*CP*GP*CP*CP*GP*TP*GP*GP*TP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	14	Total	C	N	O	P	10	0	0
			278	131	48	85	14			

- Molecule 4 is 2',3'-dideoxyadenosine triphosphate (CCD ID: DDS) (formula: C₁₀H₁₆N₅O₁₁P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			29	10	5	11	3		

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		

- Molecule 6 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	C	1	Total C O 4 2 2	0	0
6	C	1	Total C O 4 2 2	0	0

- Molecule 7 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C O 6 3 3	0	0
7	A	1	Total C O 6 3 3	0	0
7	A	1	Total C O 6 3 3	0	0
7	A	1	Total C O 6 3 3	0	0
7	A	1	Total C O 6 3 3	0	0
7	B	1	Total C O 6 3 3	0	0
7	B	1	Total C O 6 3 3	0	0

- Molecule 8 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	4	Total 4	Na 4	0	0


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	112	Total 112	O 112	0	0
9	B	23	Total 23	O 23	0	0
9	C	26	Total 26	O 26	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA polymerase I, thermostable



- Molecule 2: DNA (5'-D(*GP*AP*CP*CP*AP*CP*GP*GP*CP*GP*CP*(2DA))-3')



- Molecule 3: DNA (5'-D(*AP*AP*AP*(3DR)P*TP*GP*CP*GP*CP*CP*GP*TP*GP*GP*TP*C)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	110.23Å 110.23Å 91.27Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.73 – 2.25 47.73 – 2.25	Depositor EDS
% Data completeness (in resolution range)	99.8 (47.73-2.25) 99.8 (47.73-2.25)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.34 (at 2.24Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6_289)	Depositor
R, R_{free}	0.188 , 0.234 0.188 , 0.230	Depositor DCC
R_{free} test set	1560 reflections (4.31%)	wwPDB-VP
Wilson B-factor (Å ²)	34.1	Xtriage
Anisotropy	0.206	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 57.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.026 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5013	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, GOL, NA, 3DR, DDS, ACT, 2DA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.33	0/4324	0.75	5/5854 (0.1%)
2	B	0.18	0/249	0.79	0/382
3	C	0.19	0/297	0.87	0/454
All	All	0.32	0/4870	0.76	5/6690 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	337	GLU	CA-C-N	6.07	125.55	119.24
1	A	337	GLU	C-N-CA	6.07	125.55	119.24
1	A	700	PHE	CA-C-N	5.92	125.12	118.97
1	A	700	PHE	C-N-CA	5.92	125.12	118.97
1	A	646	MET	N-CA-C	-5.19	101.83	109.62

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4236	0	4277	97	0
2	B	242	0	134	2	0
3	C	278	0	155	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	29	0	12	1	0
5	A	1	0	0	0	0
6	A	12	0	9	0	0
6	C	8	0	6	0	0
7	A	30	0	40	11	0
7	B	12	0	16	6	0
8	A	4	0	0	0	0
9	A	112	0	0	4	0
9	B	23	0	0	2	0
9	C	26	0	0	1	0
All	All	5013	0	4649	108	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (108) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:636:ARG:HH11	1:A:651:ARG:HH12	1.18	0.90
1:A:650:PRO:HB2	1:A:651:ARG:HE	1.34	0.90
4:A:1:DDS:O2B	9:A:846:HOH:O	2.00	0.79
1:A:317:MET:HB2	7:A:6:GOL:H2	1.67	0.75
1:A:649:VAL:HG11	1:A:652:GLU:HB3	1.72	0.72
7:B:4:GOL:H12	9:B:113:HOH:O	1.90	0.72
1:A:503:ILE:HD11	1:A:522:LEU:HD13	1.72	0.70
1:A:643:ALA:HB2	1:A:662:ALA:HB2	1.74	0.69
1:A:557:THR:OG1	1:A:561:HIS:HE1	1.75	0.68
1:A:636:ARG:HH11	1:A:651:ARG:NH1	1.92	0.66
1:A:563:ARG:HG3	7:A:6:GOL:H31	1.77	0.65
2:B:107:DG:H21	7:B:8:GOL:H32	1.62	0.65
1:A:771:ARG:O	1:A:774:GLU:HG2	1.96	0.65
7:B:4:GOL:H32	9:B:161:HOH:O	1.97	0.64
1:A:549:LEU:HD22	1:A:560:LEU:HD21	1.80	0.64
1:A:650:PRO:O	1:A:651:ARG:HG3	1.98	0.63
1:A:302:PRO:HB2	1:A:328:ARG:HD3	1.83	0.60
1:A:641:GLU:O	1:A:645:TRP:CD1	2.55	0.60
1:A:380:LEU:HD12	1:A:416:LEU:HD12	1.82	0.60
1:A:627:ASN:O	1:A:631:VAL:HG23	2.01	0.60
3:C:210:DC:H2'	3:C:211:DG:C8	2.37	0.60
1:A:636:ARG:NH1	1:A:651:ARG:HH12	1.96	0.59
1:A:557:THR:HG1	1:A:561:HIS:HE1	1.49	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:563:ARG:CG	7:A:6:GOL:H31	2.34	0.58
1:A:315:GLU:HB3	7:A:6:GOL:O2	2.04	0.57
1:A:380:LEU:CD1	1:A:416:LEU:HD12	2.33	0.57
1:A:609:LEU:HG	1:A:821:VAL:HG23	1.87	0.56
1:A:728:ARG:NH2	1:A:747:MET:HE3	2.19	0.56
1:A:650:PRO:HB2	1:A:651:ARG:NE	2.14	0.56
1:A:500:LEU:HD21	1:A:526:HIS:HB2	1.88	0.56
1:A:297:ALA:HB2	1:A:331:ARG:HD3	1.89	0.55
1:A:737:VAL:HG11	7:A:7:GOL:H32	1.88	0.55
7:B:8:GOL:H12	9:C:81:HOH:O	2.07	0.54
1:A:561:HIS:O	7:A:6:GOL:H32	2.08	0.54
1:A:780:LEU:HD11	1:A:790:GLU:HB2	1.92	0.51
1:A:354:LYS:O	1:A:358:VAL:HG23	2.10	0.51
1:A:801:ARG:O	1:A:805:GLU:HG2	2.10	0.51
1:A:470:LEU:O	1:A:474:VAL:HG23	2.10	0.51
1:A:317:MET:CB	7:A:6:GOL:H2	2.39	0.51
1:A:822:GLY:HA3	1:A:830:ALA:O	2.12	0.50
1:A:650:PRO:CB	1:A:651:ARG:HE	2.17	0.50
1:A:785:ASP:OD1	1:A:786:GLU:HG3	2.12	0.50
1:A:297:ALA:CB	1:A:331:ARG:HD3	2.42	0.49
1:A:435:ARG:HB3	1:A:436:PRO:HD3	1.95	0.49
1:A:613:GLN:HG2	1:A:638:ILE:HD11	1.93	0.49
1:A:299:TRP:CG	1:A:300:PRO:HA	2.47	0.48
3:C:204:3DR:H2"	3:C:205:DT:H73	1.96	0.48
1:A:585:PRO:HB2	1:A:591:GLY:HA2	1.96	0.48
1:A:526:HIS:O	1:A:529:VAL:HG13	2.14	0.48
1:A:678:LEU:O	1:A:682:LEU:HB2	2.12	0.48
1:A:401:GLU:CD	1:A:403:GLY:H	2.22	0.47
1:A:649:VAL:CG1	1:A:652:GLU:HB3	2.43	0.47
1:A:649:VAL:HG21	1:A:657:LEU:CD1	2.44	0.47
1:A:694:GLU:OE2	1:A:694:GLU:HA	2.15	0.47
1:A:563:ARG:HG2	7:A:6:GOL:H12	1.96	0.47
1:A:445:GLU:O	1:A:561:HIS:HD2	1.97	0.47
1:A:643:ALA:HB2	1:A:662:ALA:CB	2.45	0.46
1:A:585:PRO:O	1:A:595:ARG:HG2	2.16	0.46
1:A:658:MET:HA	1:A:658:MET:HE2	1.98	0.46
1:A:492:ARG:O	1:A:496:ASP:HB2	2.16	0.45
1:A:639:HIS:HB2	1:A:659:ARG:NE	2.30	0.45
1:A:613:GLN:CG	1:A:638:ILE:HD11	2.47	0.45
1:A:631:VAL:HG13	1:A:636:ARG:HB2	1.99	0.45
1:A:721:GLU:HA	1:A:726:ARG:O	2.17	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:661:ALA:O	1:A:665:ILE:HG13	2.16	0.44
1:A:637:ASP:HB3	1:A:640:THR:OG1	2.17	0.44
9:A:123:HOH:O	7:B:8:GOL:H31	2.17	0.44
1:A:347:GLU:HA	1:A:368:PRO:HD2	1.98	0.44
1:A:392:ARG:HA	1:A:392:ARG:HD3	1.71	0.44
1:A:579:PRO:HB2	7:A:834:GOL:H2	1.99	0.44
1:A:658:MET:HA	1:A:658:MET:CE	2.47	0.44
1:A:483:ASN:H	1:A:489:GLN:HE22	1.66	0.44
1:A:669:VAL:HG21	1:A:696:TYR:CE2	2.53	0.44
1:A:520:GLU:O	1:A:523:ARG:HB2	2.18	0.43
1:A:329:GLY:C	1:A:331:ARG:H	2.26	0.43
1:A:576:SER:O	3:C:208:DG:H4'	2.17	0.43
1:A:673:MET:HG3	1:A:677:ARG:HB3	1.99	0.43
1:A:616:LEU:HD11	1:A:666:ASN:HB3	2.01	0.43
1:A:636:ARG:NH1	1:A:651:ARG:NH1	2.62	0.43
1:A:743:ALA:HA	1:A:746:ARG:NH2	2.33	0.43
1:A:346:LYS:C	1:A:346:LYS:HD2	2.43	0.43
1:A:751:MET:HB3	1:A:752:PRO:HD3	1.99	0.43
1:A:382:PRO:HD2	9:A:151:HOH:O	2.18	0.43
3:C:205:DT:H2''	3:C:206:DG:H5'	2.01	0.43
1:A:665:ILE:O	1:A:669:VAL:HG23	2.19	0.43
1:A:800:ALA:HB2	1:A:823:ILE:HD11	2.01	0.43
1:A:335:ALA:HA	1:A:336:PRO:HD3	1.89	0.42
1:A:423:GLU:OE1	1:A:727:ARG:NH2	2.51	0.42
1:A:650:PRO:HG2	1:A:651:ARG:H	1.84	0.42
1:A:397:GLU:H	1:A:408:LEU:HD13	1.85	0.42
1:A:526:HIS:HA	1:A:527:PRO:HD3	1.88	0.41
1:A:367:LEU:HA	1:A:368:PRO:HD3	1.84	0.41
1:A:712:GLU:HG2	1:A:715:ARG:HH21	1.85	0.41
7:A:836:GOL:H2	9:A:100:HOH:O	2.20	0.41
1:A:542:LYS:HA	1:A:546:ILE:HB	2.01	0.41
1:A:611:TYR:HB3	1:A:614:ILE:HB	2.01	0.41
1:A:628:LEU:HD12	1:A:628:LEU:HA	1.91	0.41
1:A:787:LEU:HD12	1:A:787:LEU:HA	1.89	0.41
2:B:107:DG:N2	7:B:8:GOL:H32	2.32	0.41
1:A:301:PRO:HG3	1:A:326:ALA:HB3	2.02	0.41
1:A:327:ALA:HA	1:A:331:ARG:O	2.20	0.41
1:A:473:GLU:OE1	1:A:473:GLU:HA	2.21	0.41
1:A:666:ASN:O	1:A:670:LEU:HD13	2.21	0.40
1:A:669:VAL:HG21	1:A:696:TYR:CZ	2.56	0.40
1:A:638:ILE:O	1:A:642:THR:N	2.31	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:763:LEU:HD11	1:A:767:LYS:HE2	2.03	0.40
1:A:782:GLN:CD	1:A:784:HIS:H	2.29	0.40
1:A:579:PRO:CB	7:A:834:GOL:H2	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	530/540 (98%)	509 (96%)	19 (4%)	2 (0%)	30 31

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	647	PHE
1	A	586	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	432/441 (98%)	417 (96%)	15 (4%)	32 39

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	295	GLU
1	A	392	ARG
1	A	416	LEU
1	A	487	ARG
1	A	522	LEU
1	A	529	VAL
1	A	638	ILE
1	A	646	MET
1	A	649	VAL
1	A	651	ARG
1	A	657	LEU
1	A	658	MET
1	A	708	GLU
1	A	787	LEU
1	A	832	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	384	ASN
1	A	489	GLN
1	A	561	HIS
1	A	582	GLN
1	A	754	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	3DR	C	204	3	8,11,12	0.41	0	7,14,17	1.39	1 (14%)
2	2DA	B	112	2,3	19,22,23	1.61	4 (21%)	25,31,34	2.31	10 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	3DR	C	204	3	-	2/3/15/16	0/1/1/1
2	2DA	B	112	2,3	-	0/7/18/19	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	112	2DA	C5-C4	4.67	1.47	1.39
2	B	112	2DA	C5-C6	2.69	1.48	1.41
2	B	112	2DA	C8-N7	2.32	1.36	1.31
2	B	112	2DA	C5-N7	-2.30	1.34	1.39

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	112	2DA	C5-C4-N3	-5.41	119.27	126.72
2	B	112	2DA	N3-C4-N9	4.62	135.02	127.17
2	B	112	2DA	N3-C2-N1	-3.74	122.92	128.58
2	B	112	2DA	C2-N3-C4	3.71	120.89	111.83
2	B	112	2DA	C3'-C2'-C1'	2.92	106.24	102.87
2	B	112	2DA	C4-N9-C8	2.92	108.80	105.74
2	B	112	2DA	C4-C5-N7	-2.90	107.26	110.58
3	C	204	3DR	C1'-C2'-C3'	2.38	105.80	103.26
2	B	112	2DA	C5-N7-C8	2.35	107.14	103.45
2	B	112	2DA	C2-N1-C6	2.30	122.51	118.73
2	B	112	2DA	N9-C8-N7	-2.01	111.09	113.94

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	204	3DR	O4'-C4'-C5'-O5'
3	C	204	3DR	C3'-C4'-C5'-O5'

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	204	3DR	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 5 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	ACT	A	2	-	3,3,3	0.82	0	3,3,3	1.31	0
7	GOL	A	835	-	5,5,5	0.35	0	5,5,5	0.33	0
6	ACT	C	6	-	3,3,3	0.78	0	3,3,3	1.59	1 (33%)
7	GOL	A	6	-	5,5,5	0.43	0	5,5,5	0.27	0
4	DDS	A	1	5	30,31,31	1.51	6 (20%)	42,48,48	1.90	9 (21%)
6	ACT	A	3	-	3,3,3	0.83	0	3,3,3	1.40	0
7	GOL	A	836	-	5,5,5	0.39	0	5,5,5	0.25	0
7	GOL	B	4	-	5,5,5	0.32	0	5,5,5	0.52	0
7	GOL	B	8	-	5,5,5	0.39	0	5,5,5	0.30	0
6	ACT	A	5	-	3,3,3	0.82	0	3,3,3	1.28	0
7	GOL	A	834	-	5,5,5	0.37	0	5,5,5	0.23	0
7	GOL	A	7	-	5,5,5	0.40	0	5,5,5	0.20	0
6	ACT	C	1	-	3,3,3	0.84	0	3,3,3	1.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	A	835	-	-	1/4/4/4	-
7	GOL	A	6	-	-	1/4/4/4	-
4	DDS	A	1	5	-	5/22/31/31	0/3/3/3
7	GOL	A	836	-	-	4/4/4/4	-
7	GOL	B	4	-	-	2/4/4/4	-
7	GOL	B	8	-	-	2/4/4/4	-
7	GOL	A	834	-	-	2/4/4/4	-
7	GOL	A	7	-	-	3/4/4/4	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1	DDS	C5-C4	4.84	1.47	1.39
4	A	1	DDS	C5-C6	2.82	1.48	1.41
4	A	1	DDS	C8-N7	2.41	1.36	1.31
4	A	1	DDS	C5-N7	-2.21	1.35	1.39
4	A	1	DDS	PB-O3B	2.04	1.61	1.59
4	A	1	DDS	PA-O3A	2.01	1.61	1.59

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1	DDS	C5-C4-N3	-6.01	118.44	126.72
4	A	1	DDS	N3-C4-N9	4.74	135.24	127.17
4	A	1	DDS	C2-N3-C4	3.77	121.03	111.83
4	A	1	DDS	C4-C5-N7	-3.35	106.75	110.58
4	A	1	DDS	N3-C2-N1	-3.21	123.72	128.58
4	A	1	DDS	C3'-C2'-C1'	2.81	106.10	102.87
4	A	1	DDS	C5-N7-C8	2.43	107.27	103.45
4	A	1	DDS	C4-N9-C8	2.32	108.17	105.74
4	A	1	DDS	C4'-O4'-C1'	2.21	111.90	109.81
6	C	6	ACT	OXT-C-CH3	2.02	123.53	115.05

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	834	GOL	O1-C1-C2-C3
7	A	7	GOL	O1-C1-C2-C3
7	B	8	GOL	O1-C1-C2-O2
7	B	8	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
7	A	7	GOL	O1-C1-C2-O2
7	B	4	GOL	O1-C1-C2-O2
7	A	836	GOL	O1-C1-C2-C3
7	B	4	GOL	O1-C1-C2-C3
7	A	834	GOL	O1-C1-C2-O2
7	A	836	GOL	O1-C1-C2-O2
4	A	1	DDS	PB-O3B-PG-O1G
7	A	835	GOL	O1-C1-C2-O2
7	A	6	GOL	O2-C2-C3-O3
7	A	836	GOL	O2-C2-C3-O3
4	A	1	DDS	PB-O3B-PG-O3G
4	A	1	DDS	C5'-O5'-PA-O1A
7	A	836	GOL	C1-C2-C3-O3
7	A	7	GOL	C1-C2-C3-O3
4	A	1	DDS	PA-O3A-PB-O1B
4	A	1	DDS	O4'-C1'-N9-C4

There are no ring outliers.

7 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	6	GOL	7	0
4	A	1	DDS	1	0
7	A	836	GOL	1	0
7	B	4	GOL	2	0
7	B	8	GOL	4	0
7	A	834	GOL	2	0
7	A	7	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	534/540 (98%)	0.00	28 (5%) 33 31	20, 47, 111, 167	0
2	B	11/12 (91%)	-1.05	0 100 100	21, 27, 70, 73	0
3	C	13/16 (81%)	-0.95	0 100 100	22, 28, 49, 63	1 (7%)
All	All	558/568 (98%)	-0.04	28 (5%) 34 33	20, 46, 110, 167	1 (0%)

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	647	PHE	4.4
1	A	648	GLY	4.1
1	A	658	MET	3.5
1	A	657	LEU	3.5
1	A	649	VAL	3.2
1	A	646	MET	3.2
1	A	650	PRO	3.1
1	A	645	TRP	3.0
1	A	298	PRO	3.0
1	A	642	THR	2.9
1	A	329	GLY	2.8
1	A	696	TYR	2.7
1	A	326	ALA	2.6
1	A	685	PRO	2.6
1	A	643	ALA	2.5
1	A	673	MET	2.4
1	A	640	THR	2.4
1	A	695	ARG	2.3
1	A	692	PHE	2.3
1	A	680	GLN	2.3
1	A	681	GLU	2.2
1	A	330	GLY	2.2
1	A	503	ILE	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	332	VAL	2.1
1	A	499	GLY	2.1
1	A	667	PHE	2.1
1	A	832	GLU	2.1
1	A	638	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	3DR	C	204	11/12	0.92	0.10	44,53,81,84	0
2	2DA	B	112	20/21	0.97	0.06	24,30,35,35	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
8	NA	A	838	1/1	0.43	0.30	86,86,86,86	0
8	NA	A	4	1/1	0.61	0.27	69,69,69,69	0
6	ACT	A	5	4/4	0.67	0.26	69,69,70,71	0
7	GOL	A	7	6/6	0.69	0.23	35,38,39,40	6
6	ACT	C	6	4/4	0.72	0.19	44,46,47,48	0
6	ACT	A	2	4/4	0.77	0.20	59,60,60,60	4
6	ACT	C	1	4/4	0.77	0.17	55,58,59,60	0
7	GOL	A	835	6/6	0.78	0.19	63,68,70,70	0
6	ACT	A	3	4/4	0.78	0.14	40,49,51,52	0
8	NA	A	839	1/1	0.82	0.16	71,71,71,71	0
7	GOL	B	8	6/6	0.82	0.17	56,65,66,67	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	GOL	A	836	6/6	0.85	0.13	47,60,63,65	0
7	GOL	B	4	6/6	0.85	0.14	37,46,52,56	0
4	DDS	A	1	29/29	0.87	0.12	49,72,94,271	0
7	GOL	A	6	6/6	0.91	0.20	34,37,38,43	0
5	MG	A	833	1/1	0.92	0.07	73,73,73,73	0
8	NA	A	837	1/1	0.93	0.10	39,39,39,39	0
7	GOL	A	834	6/6	0.96	0.07	31,31,33,34	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.