



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 08:47 AM UTC

PDB ID : 5LW9 / pdb_00005lw9
Title : Crystal structure of human JARID1B in complex with S40563a
Authors : Srikannathasan, V.; Le Bihan, Y.V.; Szykowska, A.; Johansson, C.; Gileadi, C.; Arrowsmith, C.H.; Edwards, A.M.; Bountra, C.; von Delft, F.; Oppermann, U.; Huber, K.
Deposited on : 2016-09-15
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	481	12% 81% 13% • 5%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 3854 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lysine-specific demethylase 5B, Lysine-specific demethylase 5B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	456	Total	C	N	O	S	0	7	0
			3689	2372	613	666	38			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	24	SER	-	expression tag	UNP Q9UGL1
A	25	MET	-	expression tag	UNP Q9UGL1
A	100	LYS	-	linker	UNP Q9UGL1
A	369	LEU	-	linker	UNP Q9UGL1
A	370	GLY	-	linker	UNP Q9UGL1
A	371	GLY	-	linker	UNP Q9UGL1
A	372	GLY	-	linker	UNP Q9UGL1
A	373	GLY	-	linker	UNP Q9UGL1

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Mn	0	0
			2	2		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C_2H_6OS).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	O	S	0	0
			4	2	1	1		
5	A	1	Total	C	O	S	0	0
			4	2	1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 6 is 8-[4-[2-[4-[3,5-bis(chloranyl)phenyl]piperidin-1-yl]ethyl]pyrazol-1-yl]-3 {H}-pyrido[3,4-d]pyrimidin-4-one (CCD ID: 5UJ) (formula: C₂₃H₂₂Cl₂N₆O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	Cl	N	O	0	0
			32	23	2	6	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	110	Total	O	0	0
			110	110		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Lysine-specific demethylase 5B, Lysine-specific demethylase 5B



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	142.96Å 142.96Å 152.47Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	64.92 – 2.30 64.92 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.7 (64.92-2.30) 99.7 (64.92-2.30)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.21 (at 2.29Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.185 , 0.218 0.189 , 0.218	Depositor DCC
R_{free} test set	2092 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	51.5	Xtriage
Anisotropy	0.270	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 55.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3854	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MN, 5UJ, EDO, ZN, DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.56	0/3798	0.85	0/5171

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3689	0	3469	47	0
2	A	1	0	0	0	0
3	A	2	0	0	0	0
4	A	8	0	12	1	0
5	A	12	0	18	4	0
6	A	32	0	22	2	0
7	A	110	0	0	3	1
All	All	3854	0	3521	49	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:806:DMS:H13	6:A:808:5UJ:H13	1.60	0.84
1:A:93:LEU:HD21	1:A:556:MET:HE3	1.67	0.74
1:A:509[B]:ASN:ND2	7:A:902:HOH:O	2.22	0.73
1:A:398:VAL:HG23	1:A:399:HIS:HD2	1.56	0.70
1:A:601:ASN:OD1	5:A:806:DMS:H12	2.00	0.61
1:A:25:MET:N	7:A:905:HOH:O	2.33	0.61
1:A:709:LYS:HG3	1:A:723:CYS:HB3	1.83	0.60
1:A:541:LEU:HD21	1:A:554:THR:HB	1.85	0.59
1:A:94:GLU:N	1:A:554:THR:OG1	2.37	0.58
1:A:680:MET:HB2	1:A:733:ARG:NH1	2.19	0.57
1:A:537:LEU:HD22	1:A:556:MET:HE2	1.86	0.56
1:A:447:LEU:HD13	1:A:455:LEU:HD11	1.88	0.56
1:A:442:ASP:O	1:A:445:ILE:N	2.39	0.55
1:A:695:CYS:HB3	1:A:718:HIS:CE1	2.43	0.53
1:A:621:LEU:HD23	5:A:805:DMS:H12	1.93	0.51
1:A:400:MET:O	1:A:402:PRO:HD3	2.11	0.50
6:A:808:5UJ:N4	6:A:808:5UJ:H9	2.27	0.49
1:A:37:PRO:HD2	1:A:65[B]:ARG:O	2.14	0.48
1:A:90:LEU:HD21	1:A:493:PHE:HB3	1.96	0.47
1:A:89:ARG:HH22	1:A:94:GLU:CD	2.20	0.47
1:A:676:ASP:OD2	1:A:731:LYS:HE2	2.15	0.47
1:A:37:PRO:HD2	1:A:65[A]:ARG:O	2.15	0.46
1:A:37:PRO:HB3	1:A:48:PHE:CE1	2.50	0.46
1:A:74:PHE:O	7:A:901:HOH:O	2.21	0.46
1:A:36:GLU:OE2	1:A:65[A]:ARG:NH2	2.47	0.46
1:A:680:MET:HE2	1:A:682:PHE:CZ	2.51	0.46
1:A:413:LEU:HD12	1:A:413:LEU:HA	1.75	0.45
1:A:61:ILE:HG22	1:A:581:THR:HG22	1.97	0.45
1:A:72:PRO:HA	1:A:73:PRO:HD3	1.89	0.45
1:A:98:ARG:HG3	5:A:807:DMS:H21	1.99	0.45
1:A:398:VAL:HG23	1:A:399:HIS:CD2	2.42	0.45
1:A:400:MET:HB3	1:A:400:MET:HE2	1.65	0.44
1:A:701:MET:HB2	4:A:803:EDO:H21	1.99	0.44
1:A:399:HIS:CD2	1:A:399:HIS:N	2.85	0.44
1:A:69:ASP:OD1	1:A:69:ASP:N	2.47	0.44
1:A:474[A]:HIS:CE1	1:A:636:MET:HE3	2.53	0.44
1:A:93:LEU:HA	1:A:93:LEU:HD23	1.76	0.43
1:A:482:MET:HE2	1:A:482:MET:HB3	1.94	0.43
1:A:39:TRP:HZ3	1:A:69:ASP:CG	2.27	0.42
1:A:95:ALA:HB1	1:A:552:LEU:HD22	2.01	0.42
1:A:67:PRO:HA	1:A:68:PRO:HD3	1.87	0.42
1:A:465:PRO:HG3	1:A:600:VAL:HG21	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:709:LYS:HB3	1:A:712:LEU:HD12	2.01	0.42
1:A:424:GLU:O	1:A:491:MET:HG2	2.20	0.41
1:A:530:LEU:C	1:A:530:LEU:HD23	2.45	0.41
1:A:603:CYS:HB3	1:A:607:TRP:CE3	2.55	0.41
1:A:479:ILE:O	1:A:481:GLY:N	2.43	0.41
1:A:633:ILE:HD13	1:A:633:ILE:HA	1.82	0.40
1:A:680:MET:HB2	1:A:733:ARG:HH11	1.85	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:979:HOH:O	7:A:979:HOH:O[9_765]	1.91	0.29

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	456/481 (95%)	446 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	391/429 (91%)	384 (98%)	7 (2%)	51 70

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	99	VAL
1	A	413	LEU
1	A	455	LEU
1	A	464	MET
1	A	471	VAL
1	A	554	THR
1	A	621	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	399	HIS
1	A	462	ASN
1	A	617	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 9 ligands modelled in this entry, 3 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	DMS	A	807	-	3,3,3	0.65	0	3,3,3	0.47	0
5	DMS	A	805	-	3,3,3	0.74	0	3,3,3	0.79	0
5	DMS	A	806	-	3,3,3	0.59	0	3,3,3	0.40	0
4	EDO	A	803	-	3,3,3	0.68	0	2,2,2	0.13	0
4	EDO	A	804	-	3,3,3	0.48	0	2,2,2	0.23	0
6	5UJ	A	808	3	35,36,36	0.46	1 (2%)	46,51,51	0.68	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	803	-	-	0/1/1/1	-
4	EDO	A	804	-	-	1/1/1/1	-
6	5UJ	A	808	3	-	4/13/23/23	0/5/5/5

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	808	5UJ	C19-N4	2.02	1.33	1.29

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	808	5UJ	C18-C14-N3	2.48	123.70	118.55

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	A	808	5UJ	N3-C14-N1-N2
6	A	808	5UJ	C18-C14-N1-N2
6	A	808	5UJ	N3-C14-N1-C12
6	A	808	5UJ	C18-C14-N1-C12
4	A	804	EDO	O1-C1-C2-O2

There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	807	DMS	1	0
5	A	805	DMS	1	0
5	A	806	DMS	2	0
4	A	803	EDO	1	0
6	A	808	5UJ	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	456/481 (94%)	0.48	60 (13%) 7 8	22, 59, 128, 172	7 (1%)

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	446	LYS	7.5
1	A	447	LEU	6.9
1	A	548	LEU	6.3
1	A	99	VAL	6.2
1	A	543	VAL	5.8
1	A	544	SER	5.5
1	A	434	PHE	5.5
1	A	481	GLY	5.3
1	A	479	ILE	5.0
1	A	450	GLU	4.6
1	A	546	PRO	4.6
1	A	81[A]	LEU	4.5
1	A	76	CYS	4.5
1	A	445	ILE	4.4
1	A	540	GLU	4.3
1	A	754	SER	4.1
1	A	100	LYS	4.1
1	A	75	ALA	4.0
1	A	398	VAL	4.0
1	A	93	LEU	3.9
1	A	468	GLU	3.8
1	A	448	SER	3.8
1	A	480	CYS	3.7
1	A	452	GLU	3.7
1	A	451	GLU	3.7
1	A	545	GLN	3.7
1	A	89	ARG	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	550	HIS	3.6
1	A	549	LEU	3.5
1	A	98	ARG	3.5
1	A	478	ASP	3.4
1	A	82[A]	HIS	3.3
1	A	449	PRO	3.3
1	A	374	ALA	3.3
1	A	94	GLU	3.3
1	A	399	HIS	3.2
1	A	74	PHE	3.2
1	A	727	PRO	3.2
1	A	417	ILE	3.2
1	A	541	LEU	3.1
1	A	551	GLN	3.1
1	A	552	LEU	2.9
1	A	547	ASP	2.9
1	A	553	VAL	2.9
1	A	77	ASP	2.9
1	A	443	GLY	2.8
1	A	78	VAL	2.7
1	A	95	ALA	2.6
1	A	442	ASP	2.5
1	A	418	GLU	2.5
1	A	554	THR	2.4
1	A	537	LEU	2.3
1	A	441	ARG	2.3
1	A	542	PHE	2.2
1	A	708	CYS	2.2
1	A	620	LEU	2.2
1	A	538	ALA	2.2
1	A	724	SER	2.2
1	A	453	GLU	2.1
1	A	440	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	DMS	A	805	4/4	0.82	0.28	77,78,80,101	0
3	MN	A	802	1/1	0.83	0.13	130,130,130,130	1
5	DMS	A	807	4/4	0.84	0.36	75,115,167,170	0
4	EDO	A	804	4/4	0.89	0.20	79,81,83,92	0
4	EDO	A	803	4/4	0.91	0.17	38,43,47,59	0
5	DMS	A	806	4/4	0.92	0.23	64,91,135,148	0
6	5UJ	A	808	32/32	0.95	0.10	38,51,94,170	7
2	ZN	A	801	1/1	1.00	0.02	49,49,49,49	0
3	MN	A	809	1/1	1.00	0.01	39,39,39,39	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 5UJ A 808:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.