



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 02:03 PM UTC

PDB ID : 5LXM / pdb_00005lxm
Title : Crystal structure of Aurora-A bound to a hydrocarbon-stapled proteomimetic of TPX2
Authors : McIntyre, P.J.; Bayliss, R.
Deposited on : 2016-09-22
Resolution : 2.08 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	8172 (2.10-2.06)
Clashscore	190562	8714 (2.10-2.06)
Ramachandran outliers	187476	8641 (2.10-2.06)
Sidechain outliers	187428	8642 (2.10-2.06)
RSRZ outliers	180081	8177 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	283	5% 86% 8% 6%
2	D	39	15% 77% 8% 13%

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 2615 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aurora kinase A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	267	Total	C	N	O	P	S	1	5	0
			2181	1398	379	398	1	5			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	121	MET	-	initiating methionine	UNP O14965
A	290	ALA	CYS	engineered mutation	UNP O14965
A	393	ALA	CYS	engineered mutation	UNP O14965


- Molecule 2 is a protein called Targeting protein for Xklp2.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	D	34	Total	C	N	O	0	0	1
			272	173	41	58			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
D	37	MK8	GLU	conflict	UNP Q9ULW0
D	44	NH2	-	amidation	UNP Q9ULW0

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			7	4	3		
6	A	1	Total	C	O	0	0
			7	4	3		


- Molecule 7 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula:

C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 8 is 1-ETHOXY-2-(2-ETHOXYETHOXY)ETHANE (CCD ID: P4G) (formula: C₈H₁₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			11	8	3		

- Molecule 9 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	1	Total	Cl	0	0
			1	1		

- Molecule 10 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	2	Total	Mg	0	0
			2	2		


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	70	Total	O	0	0
			70	70		
11	D	7	Total	O	0	0
			7	7		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aurora kinase A



- Molecule 2: Targeting protein for Xklp2



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	79.78Å 79.78Å 140.26Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	61.98 – 2.08 61.98 – 2.08	Depositor EDS
% Data completeness (in resolution range)	99.3 (61.98-2.08) 89.4 (61.98-2.08)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.85 (at 2.08Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.200 , 0.243 0.213 , 0.252	Depositor DCC
R_{free} test set	1603 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	44.7	Xtriage
Anisotropy	0.641	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 65.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	0.059 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2615	wwPDB-VP
Average B, all atoms (Å ²)	74.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ADP, MK8, MG, SO4, CL, PEG, NH2, MES, P4G, PG4, TPO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/2231	0.68	2/3025 (0.1%)
2	D	0.33	0/257	0.64	0/344
All	All	0.27	0/2488	0.67	2/3369 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	213	ALA	CA-C-N	5.81	125.67	119.28
1	A	213	ALA	C-N-CA	5.81	125.67	119.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2181	0	2126	17	0
2	D	272	0	234	3	0
3	A	27	0	12	1	0
4	A	13	0	18	2	0
5	A	5	0	0	0	0
6	A	14	0	20	3	0
7	A	12	0	13	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	A	11	0	18	0	0
9	A	1	0	0	0	0
10	A	2	0	0	0	0
11	A	70	0	0	0	0
11	D	7	0	0	1	0
All	All	2615	0	2441	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:211:GLU:OE1	1:A:271:LYS:NZ	2.33	0.62
1:A:295:TYR:OH	1:A:321:GLU:OE1	2.17	0.59
1:A:221[B]:GLU:OE1	1:A:232:ARG:NH2	2.32	0.59
1:A:374:LEU:HB2	6:A:504:PEG:H21	1.87	0.56
2:D:6:SER:N	11:D:201:HOH:O	2.39	0.55
3:A:501:ADP:H8	3:A:501:ADP:H5'1	1.73	0.54
1:A:250:LYS:HE3	2:D:32:ASP:HB3	1.88	0.54
1:A:248:HIS:ND1	6:A:504:PEG:O4	2.23	0.51
1:A:162:LYS:HB3	1:A:208:LEU:HB2	1.95	0.49
1:A:374:LEU:H	6:A:504:PEG:H32	1.78	0.48
1:A:224:LYS:NZ	4:A:502:PG4:H51	2.30	0.47
1:A:224:LYS:HZ3	4:A:502:PG4:H51	1.80	0.47
1:A:180:ARG:NH1	7:A:506:MES:O3S	2.50	0.45
1:A:166:LYS:HZ2	1:A:203:ALA:HA	1.82	0.45
1:A:286:ARG:HB2	1:A:306:HIS:CE1	2.51	0.45
2:D:37:MK8:HB1B	2:D:37:MK8:HGA	1.77	0.44
1:A:233:THR:O	1:A:237:ILE:HG12	2.20	0.42
1:A:230:GLU:OE2	1:A:353:THR:OG1	2.34	0.42
1:A:286:ARG:HB3	1:A:288:TPO:O2P	2.20	0.41
1:A:224:LYS:HB3	1:A:224:LYS:HE2	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	269/283 (95%)	256 (95%)	13 (5%)	0	100	100
2	D	28/39 (72%)	27 (96%)	1 (4%)	0	100	100
All	All	297/322 (92%)	283 (95%)	14 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	227/248 (92%)	225 (99%)	2 (1%)	70	77
2	D	29/33 (88%)	29 (100%)	0	100	100
All	All	256/281 (91%)	254 (99%)	2 (1%)	73	80

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	180	ARG
1	A	336	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	TPO	A	288	1	8,10,11	1.66	1 (12%)	10,14,16	1.83	1 (10%)
2	MK8	D	41	2	5,8,9	1.52	1 (20%)	6,10,12	1.12	0
2	MK8	D	37	2	5,8,9	1.55	1 (20%)	6,10,12	1.76	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	288	1	-	0/9/11/13	-
2	MK8	D	41	2	-	3/6/8/11	-
2	MK8	D	37	2	-	1/6/8/11	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	288	TPO	P-O1P	3.56	1.61	1.50
2	D	37	MK8	CB-CA	-2.81	1.51	1.55
2	D	41	MK8	CB-CA	-2.78	1.52	1.55

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	288	TPO	P-OG1-CB	-5.01	109.72	123.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	37	MK8	CG-CB-CA	-3.95	109.58	115.41

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	37	MK8	CA-CB-CG-CD
2	D	41	MK8	C-CA-CB-CG
2	D	41	MK8	N-CA-CB-CG
2	D	41	MK8	CB1-CA-CB-CG

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	288	TPO	1	0
2	D	37	MK8	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 3 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	SO4	A	503	-	4,4,4	0.24	0	6,6,6	0.09	0
8	P4G	A	507	-	10,10,10	0.61	0	9,9,9	0.19	0
6	PEG	A	505	10	6,6,6	0.58	0	5,5,5	0.26	0
4	PG4	A	502	-	12,12,12	0.69	0	11,11,11	0.20	0
3	ADP	A	501	10	28,29,29	1.43	5 (17%)	43,45,45	1.80	7 (16%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	MES	A	506	-	12,12,12	1.22	2 (16%)	15,16,16	2.52	7 (46%)
6	PEG	A	504	-	6,6,6	0.56	0	5,5,5	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	P4G	A	507	-	-	3/8/8/8	-
6	PEG	A	505	10	-	3/4/4/4	-
4	PG4	A	502	-	-	4/10/10/10	-
7	MES	A	506	-	-	2/6/14/14	0/1/1/1
3	ADP	A	501	10	-	4/16/32/32	0/3/3/3
6	PEG	A	504	-	-	3/4/4/4	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	501	ADP	C5-C4	4.80	1.47	1.39
3	A	501	ADP	C5-C6	2.68	1.48	1.41
3	A	501	ADP	C5-N7	-2.32	1.34	1.39
7	A	506	MES	O2S-S	2.27	1.51	1.45
7	A	506	MES	O1S-S	2.26	1.51	1.45
3	A	501	ADP	C8-N7	2.25	1.36	1.31
3	A	501	ADP	PA-O3A	2.00	1.61	1.59

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	ADP	C5-C4-N3	-6.12	118.28	126.72
3	A	501	ADP	N3-C4-N9	4.79	135.31	127.17
7	A	506	MES	C6-C5-N4	4.26	116.60	110.12
7	A	506	MES	O3S-S-O1S	-4.09	101.16	111.40
7	A	506	MES	C2-C3-N4	4.04	116.25	110.12
7	A	506	MES	C5-N4-C3	4.00	117.46	108.84
3	A	501	ADP	C2-N3-C4	3.75	121.00	111.83
3	A	501	ADP	C4-C5-N7	-3.32	106.79	110.58
7	A	506	MES	O1S-S-C8	3.10	111.41	106.73
3	A	501	ADP	N3-C2-N1	-3.08	123.92	128.58

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	506	MES	O3S-S-C8	2.58	111.05	106.00
7	A	506	MES	O2S-S-C8	2.43	110.41	106.73
3	A	501	ADP	C5-N7-C8	2.37	107.17	103.45
3	A	501	ADP	C4-N9-C8	2.12	107.97	105.74

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	501	ADP	C5'-O5'-PA-O1A
6	A	505	PEG	O2-C3-C4-O4
6	A	505	PEG	O1-C1-C2-O2
4	A	502	PG4	O3-C5-C6-O4
6	A	504	PEG	C1-C2-O2-C3
8	A	507	P4G	O2-C3-C4-O3
3	A	501	ADP	C5'-O5'-PA-O3A
7	A	506	MES	C8-C7-N4-C3
6	A	505	PEG	C4-C3-O2-C2
4	A	502	PG4	O1-C1-C2-O2
8	A	507	P4G	C5-C6-O4-C7
6	A	504	PEG	O1-C1-C2-O2
6	A	504	PEG	C4-C3-O2-C2
4	A	502	PG4	C3-C4-O3-C5
7	A	506	MES	C7-C8-S-O1S
8	A	507	P4G	C6-C5-O3-C4
4	A	502	PG4	C8-C7-O4-C6
3	A	501	ADP	O4'-C4'-C5'-O5'
3	A	501	ADP	PB-O3A-PA-O1A


There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	502	PG4	2	0
3	A	501	ADP	1	0
7	A	506	MES	1	0
6	A	504	PEG	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	266/283 (93%)	0.63	13 (4%) 35 36	35, 68, 102, 132	6 (2%)
2	D	31/39 (79%)	1.43	6 (19%) 3 3	54, 89, 141, 149	0
All	All	297/322 (92%)	0.71	19 (6%) 25 26	35, 69, 107, 149	6 (2%)

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	22	LEU	6.8
1	A	203	ALA	4.7
2	D	28	THR	4.1
1	A	283[A]	SER	3.7
1	A	390	PRO	3.1
2	D	30	ASN	2.9
2	D	33	SER	2.8
1	A	126	ARG	2.8
2	D	32	ASP	2.7
2	D	31	ILE	2.6
1	A	392	ASN	2.5
1	A	169	LEU	2.4
1	A	303	GLY	2.4
1	A	151	ARG	2.3
1	A	334	TYR	2.2
1	A	131	GLU	2.2
1	A	391	SER	2.1
1	A	207	TYR	2.1
1	A	232	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MK8	D	41	9/10	0.95	0.11	51,58,60,67	0
1	TPO	A	288	11/12	0.97	0.07	48,59,62,78	0
2	MK8	D	37	9/10	0.97	0.09	63,65,74,76	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	SO4	A	503	5/5	0.69	0.20	142,143,145,146	0
8	P4G	A	507	11/11	0.85	0.21	69,96,105,107	0
10	MG	A	509	1/1	0.85	0.13	143,143,143,143	0
4	PG4	A	502	13/13	0.86	0.23	93,105,111,111	0
3	ADP	A	501	27/27	0.87	0.13	50,74,163,169	0
6	PEG	A	505	7/7	0.87	0.21	91,95,105,110	0
7	MES	A	506	12/12	0.88	0.15	79,85,112,125	0
6	PEG	A	504	7/7	0.89	0.14	59,66,74,85	0
10	MG	A	510	1/1	0.89	0.15	144,144,144,144	0
9	CL	A	508	1/1	0.95	0.11	116,116,116,116	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.