



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 18, 2026 – 07:36 AM UTC

PDB ID : 4LY8 / pdb_00004ly8
Title : dihydrodipicolinate synthase from *C. jejuni* with pyruvate bound to the active site
Authors : Conly, C.J.T.
Deposited on : 2013-07-30
Resolution : 1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	 86% 10% ..
1	B	306	 88% 8% ..
1	C	306	 2% 87% 10% .
1	D	306	 88% 8% ..

2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 10073 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 4-hydroxy-tetrahydrodipicolinate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	298	2316	1472	384	446	14	0	4	0
1	B	297	2296	1459	383	441	13	0	1	0
1	C	298	2301	1463	381	443	14	0	1	0
1	D	297	2288	1454	380	441	13	2	0	0

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	HIS	-	expression tag	UNP Q9PPB4
A	-6	HIS	-	expression tag	UNP Q9PPB4
A	-5	HIS	-	expression tag	UNP Q9PPB4
A	-4	HIS	-	expression tag	UNP Q9PPB4
A	-3	HIS	-	expression tag	UNP Q9PPB4
A	-2	HIS	-	expression tag	UNP Q9PPB4
A	-1	GLY	-	expression tag	UNP Q9PPB4
A	0	SER	-	expression tag	UNP Q9PPB4
B	-7	HIS	-	expression tag	UNP Q9PPB4
B	-6	HIS	-	expression tag	UNP Q9PPB4
B	-5	HIS	-	expression tag	UNP Q9PPB4
B	-4	HIS	-	expression tag	UNP Q9PPB4
B	-3	HIS	-	expression tag	UNP Q9PPB4
B	-2	HIS	-	expression tag	UNP Q9PPB4
B	-1	GLY	-	expression tag	UNP Q9PPB4
B	0	SER	-	expression tag	UNP Q9PPB4
C	-7	HIS	-	expression tag	UNP Q9PPB4
C	-6	HIS	-	expression tag	UNP Q9PPB4
C	-5	HIS	-	expression tag	UNP Q9PPB4
C	-4	HIS	-	expression tag	UNP Q9PPB4
C	-3	HIS	-	expression tag	UNP Q9PPB4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-2	HIS	-	expression tag	UNP Q9PPB4
C	-1	GLY	-	expression tag	UNP Q9PPB4
C	0	SER	-	expression tag	UNP Q9PPB4
D	-7	HIS	-	expression tag	UNP Q9PPB4
D	-6	HIS	-	expression tag	UNP Q9PPB4
D	-5	HIS	-	expression tag	UNP Q9PPB4
D	-4	HIS	-	expression tag	UNP Q9PPB4
D	-3	HIS	-	expression tag	UNP Q9PPB4
D	-2	HIS	-	expression tag	UNP Q9PPB4
D	-1	GLY	-	expression tag	UNP Q9PPB4
D	0	SER	-	expression tag	UNP Q9PPB4

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: C₆H₁₄O₄).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 10 6 4	0	0
3	A	1	Total C O 10 6 4	0	0
3	A	1	Total C O 10 6 4	0	0
3	A	1	Total C O 10 6 4	0	0
3	B	1	Total C O 10 6 4	0	0
3	B	1	Total C O 10 6 4	0	0
3	B	1	Total C O 10 6 4	0	0
3	B	1	Total C O 10 6 4	0	0
3	C	1	Total C O 10 6 4	0	0
3	C	1	Total C O 10 6 4	0	0
3	D	1	Total C O 10 6 4	0	0

- Molecule 4 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: C₈H₁₈O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			13	8	5		
6	C	1	Total	C	O	0	0
			13	8	5		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	163	Total	O	0	0
			163	163		
7	B	156	Total	O	0	0
			156	156		
7	C	145	Total	O	0	0
			145	145		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	D	146	Total 146	O 146	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 4-hydroxy-tetrahydrodipicolinate synthase




- Molecule 1: 4-hydroxy-tetrahydrodipicolinate synthase



- Molecule 1: 4-hydroxy-tetrahydrodipicolinate synthase



- Molecule 1: 4-hydroxy-tetrahydrodipicolinate synthase





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	81.25Å 101.40Å 147.96Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.01 – 1.70 43.01 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.7 (43.01-1.70) 99.6 (43.01-1.70)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.04 (at 1.60Å)	Xtrriage
Refinement program	PHENIX dev_1327	Depositor
R, R_{free}	0.209 , 0.256 0.212 , 0.256	Depositor DCC
R_{free} test set	7948 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	13.6	Xtrriage
Anisotropy	0.194	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 40.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.41$, $\langle L^2 \rangle = 0.24$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10073	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, GOL, PG4, ACT, PGE, KPI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/2352	0.77	0/3176
1	B	0.43	0/2323	0.76	0/3138
1	C	0.41	0/2328	0.76	0/3146
1	D	0.43	0/2312	0.76	0/3124
All	All	0.43	0/9315	0.76	0/12584

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2316	0	2364	25	0
1	B	2296	0	2336	16	0
1	C	2301	0	2341	19	0
1	D	2288	0	2323	16	0
2	A	24	0	36	0	0
2	B	12	0	18	0	0
2	C	20	0	30	0	0
2	D	28	0	42	0	0
3	A	40	0	56	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	40	0	56	1	0
3	C	20	0	28	3	0
3	D	10	0	14	0	0
4	A	8	0	6	0	0
4	C	4	0	3	0	0
5	A	12	0	16	0	0
5	B	12	0	16	1	0
5	D	6	0	8	0	0
6	A	13	0	18	2	0
6	C	13	0	18	0	0
7	A	163	0	0	1	0
7	B	156	0	0	0	0
7	C	145	0	0	2	0
7	D	146	0	0	0	0
All	All	10073	0	9729	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 72 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:150:ASP:HB3	6:A:313:PG4:H71	1.61	0.82
1:A:280:LYS:HE3	1:A:280:LYS:H	1.49	0.75
1:A:2:ASP:N	1:A:2:ASP:OD1	2.28	0.66
1:A:1:MET:H3	1:A:184:ARG:HD2	1.62	0.65
1:D:45:VAL:HG11	1:D:59:HIS:CD2	2.31	0.64

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	299/306 (98%)	292 (98%)	6 (2%)	1 (0%)	36	22
1	B	295/306 (96%)	289 (98%)	6 (2%)	0	100	100
1	C	296/306 (97%)	289 (98%)	6 (2%)	1 (0%)	36	22
1	D	294/306 (96%)	286 (97%)	7 (2%)	1 (0%)	36	22
All	All	1184/1224 (97%)	1156 (98%)	25 (2%)	3 (0%)	36	22

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	3	LYS
1	C	2	ASP
1	D	3	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	254/257 (99%)	249 (98%)	5 (2%)	48	32
1	B	250/257 (97%)	247 (99%)	3 (1%)	63	51
1	C	251/257 (98%)	247 (98%)	4 (2%)	55	41
1	D	249/257 (97%)	244 (98%)	5 (2%)	48	32
All	All	1004/1028 (98%)	987 (98%)	17 (2%)	53	38

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	160	GLU
1	D	232	GLU
1	B	161	ASN
1	C	1	MET
1	C	45	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	56	HIS
1	B	242	ASN
1	D	56	HIS
1	D	223	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	KPI	D	166	1	11,13,14	2.12	4 (36%)	9,15,17	3.35	2 (22%)
1	KPI	B	166	1	11,13,14	2.11	3 (27%)	9,15,17	3.03	3 (33%)
1	KPI	A	166	1	11,13,14	2.22	3 (27%)	9,15,17	2.75	3 (33%)
1	KPI	C	166	1	11,13,14	1.97	2 (18%)	9,15,17	3.61	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KPI	D	166	1	-	2/13/14/16	-
1	KPI	B	166	1	-	3/13/14/16	-
1	KPI	A	166	1	-	3/13/14/16	-
1	KPI	C	166	1	-	3/13/14/16	-

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	166	KPI	C1-CX1	-5.13	1.39	1.49
1	D	166	KPI	C1-CX1	-5.01	1.40	1.49
1	C	166	KPI	C1-CX1	-4.98	1.40	1.49
1	B	166	KPI	C1-CX1	-4.96	1.40	1.49
1	A	166	KPI	CX2-CX1	3.85	1.54	1.49

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	166	KPI	C1-CX1-CX2	-10.10	108.59	118.11
1	D	166	KPI	C1-CX1-CX2	-9.35	109.29	118.11
1	B	166	KPI	C1-CX1-CX2	-8.29	110.29	118.11
1	A	166	KPI	C1-CX1-CX2	-6.74	111.76	118.11
1	A	166	KPI	CX2-CX1-NZ	3.15	122.29	114.88

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	166	KPI	C1-CX1-NZ-CE
1	A	166	KPI	NZ-CX1-CX2-O1
1	A	166	KPI	NZ-CX1-CX2-O2
1	B	166	KPI	C1-CX1-NZ-CE
1	C	166	KPI	C1-CX1-NZ-CE

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	166	KPI	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

42 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PGE	B	306	-	9,9,9	0.68	0	8,8,8	0.75	0
5	GOL	D	302	-	5,5,5	0.34	0	5,5,5	0.23	0
2	EDO	C	302	-	3,3,3	0.44	0	2,2,2	0.41	0
2	EDO	D	309	-	3,3,3	0.45	0	2,2,2	0.33	0
3	PGE	C	304	-	9,9,9	0.68	0	8,8,8	0.81	0
3	PGE	A	307	-	9,9,9	0.66	0	8,8,8	0.69	0
2	EDO	B	308	-	3,3,3	0.41	0	2,2,2	0.48	0
5	GOL	B	307	-	5,5,5	0.37	0	5,5,5	0.36	0
2	EDO	A	311	-	3,3,3	0.40	0	2,2,2	0.42	0
3	PGE	A	302	-	9,9,9	0.66	0	8,8,8	0.79	0
2	EDO	C	306	-	3,3,3	0.47	0	2,2,2	0.29	0
6	PG4	C	301	-	12,12,12	0.70	0	11,11,11	1.45	0
5	GOL	A	314	-	5,5,5	0.36	0	5,5,5	0.41	0
2	EDO	A	315	-	3,3,3	0.41	0	2,2,2	0.42	0
3	PGE	A	304	-	9,9,9	0.66	0	8,8,8	0.66	0
2	EDO	A	309	-	3,3,3	0.38	0	2,2,2	0.49	0
2	EDO	D	306	-	3,3,3	0.42	0	2,2,2	0.42	0
2	EDO	B	304	-	3,3,3	0.40	0	2,2,2	0.44	0
6	PG4	A	313	-	12,12,12	0.65	0	11,11,11	1.59	2 (18%)
2	EDO	D	308	-	3,3,3	0.40	0	2,2,2	0.41	0
2	EDO	A	303	-	3,3,3	0.42	0	2,2,2	0.28	0
3	PGE	B	303	-	9,9,9	0.67	0	8,8,8	0.70	0
3	PGE	B	301	-	9,9,9	0.62	0	8,8,8	0.75	0
5	GOL	A	310	-	5,5,5	0.37	0	5,5,5	0.15	0
2	EDO	C	305	-	3,3,3	0.44	0	2,2,2	0.35	0
3	PGE	A	305	-	9,9,9	0.64	0	8,8,8	0.78	0
2	EDO	D	307	-	3,3,3	0.47	0	2,2,2	0.29	0
2	EDO	C	307	-	3,3,3	0.47	0	2,2,2	0.37	0
3	PGE	D	301	-	9,9,9	0.69	0	8,8,8	0.71	0
2	EDO	D	304	-	3,3,3	0.42	0	2,2,2	0.44	0
3	PGE	B	305	-	9,9,9	0.66	0	8,8,8	0.79	0
2	EDO	C	303	-	3,3,3	0.44	0	2,2,2	0.41	0
2	EDO	A	301	-	3,3,3	0.42	0	2,2,2	0.29	0
5	GOL	B	302	-	5,5,5	0.34	0	5,5,5	0.38	0
4	ACT	A	308	-	3,3,3	0.83	0	3,3,3	1.27	0
2	EDO	A	306	-	3,3,3	0.49	0	2,2,2	0.19	0
4	ACT	C	309	-	3,3,3	0.92	0	3,3,3	1.45	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	EDO	D	303	-	3,3,3	0.37	0	2,2,2	0.47	0
4	ACT	A	312	-	3,3,3	0.79	0	3,3,3	1.28	0
2	EDO	B	309	-	3,3,3	0.46	0	2,2,2	0.29	0
2	EDO	D	305	-	3,3,3	0.45	0	2,2,2	0.45	0
3	PGE	C	308	-	9,9,9	0.64	0	8,8,8	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PGE	B	306	-	-	5/7/7/7	-
5	GOL	D	302	-	-	4/4/4/4	-
2	EDO	C	302	-	-	0/1/1/1	-
2	EDO	D	309	-	-	0/1/1/1	-
3	PGE	C	304	-	-	4/7/7/7	-
3	PGE	A	307	-	-	2/7/7/7	-
2	EDO	B	308	-	-	1/1/1/1	-
5	GOL	B	307	-	-	4/4/4/4	-
2	EDO	A	311	-	-	0/1/1/1	-
3	PGE	A	302	-	-	4/7/7/7	-
2	EDO	C	306	-	-	1/1/1/1	-
6	PG4	C	301	-	-	6/10/10/10	-
5	GOL	A	314	-	-	4/4/4/4	-
2	EDO	A	315	-	-	0/1/1/1	-
3	PGE	A	304	-	-	4/7/7/7	-
2	EDO	A	309	-	-	1/1/1/1	-
2	EDO	D	306	-	-	1/1/1/1	-
2	EDO	B	304	-	-	0/1/1/1	-
6	PG4	A	313	-	-	5/10/10/10	-
2	EDO	D	308	-	-	1/1/1/1	-
2	EDO	A	303	-	-	0/1/1/1	-
3	PGE	B	303	-	-	5/7/7/7	-
3	PGE	B	301	-	-	1/7/7/7	-
5	GOL	A	310	-	-	2/4/4/4	-
2	EDO	C	305	-	-	0/1/1/1	-
3	PGE	A	305	-	-	2/7/7/7	-
2	EDO	D	307	-	-	0/1/1/1	-
2	EDO	C	307	-	-	1/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PGE	D	301	-	-	2/7/7/7	-
2	EDO	D	304	-	-	0/1/1/1	-
3	PGE	B	305	-	-	3/7/7/7	-
2	EDO	C	303	-	-	0/1/1/1	-
2	EDO	A	301	-	-	1/1/1/1	-
5	GOL	B	302	-	-	4/4/4/4	-
2	EDO	A	306	-	-	0/1/1/1	-
2	EDO	D	303	-	-	1/1/1/1	-
2	EDO	B	309	-	-	0/1/1/1	-
2	EDO	D	305	-	-	1/1/1/1	-
3	PGE	C	308	-	-	4/7/7/7	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	313	PG4	O4-C7-C8	2.22	119.89	110.11
6	A	313	PG4	O4-C6-C5	2.11	119.99	110.35

There are no chirality outliers.

5 of 74 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	310	GOL	C1-C2-C3-O3
5	A	314	GOL	O1-C1-C2-C3
5	A	314	GOL	C1-C2-C3-O3
5	B	302	GOL	O1-C1-C2-C3
5	B	302	GOL	C1-C2-C3-O3

There are no ring outliers.

6 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	304	PGE	1	0
3	A	304	PGE	2	0
6	A	313	PG4	2	0
3	B	303	PGE	1	0
5	B	302	GOL	1	0
3	C	308	PGE	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	297/306 (97%)	-0.36	4 (1%) 75 79	13, 21, 38, 56	7 (2%)
1	B	296/306 (96%)	-0.28	3 (1%) 79 82	12, 23, 42, 60	3 (1%)
1	C	297/306 (97%)	0.14	6 (2%) 65 69	11, 28, 47, 59	4 (1%)
1	D	296/306 (96%)	-0.08	2 (0%) 84 86	19, 28, 48, 91	1 (0%)
All	All	1186/1224 (96%)	-0.14	15 (1%) 75 79	11, 25, 45, 91	15 (1%)

The worst 5 of 15 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	3	LYS	23.1
1	C	1	MET	21.5
1	C	2	ASP	20.6
1	A	2	ASP	4.0
1	A	1	MET	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	KPI	C	166	14/15	0.92	0.08	23,28,41,42	0
1	KPI	B	166	14/15	0.95	0.08	15,21,34,35	0
1	KPI	A	166	14/15	0.95	0.08	13,16,31,31	0
1	KPI	D	166	14/15	0.95	0.08	20,24,31,32	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	A	310	6/6	0.64	0.19	58,59,59,60	0
2	EDO	B	304	4/4	0.69	0.23	63,64,64,65	0
5	GOL	D	302	6/6	0.69	0.16	50,53,54,56	0
4	ACT	C	309	4/4	0.71	0.15	37,42,44,52	0
5	GOL	A	314	6/6	0.73	0.29	46,52,54,57	6
4	ACT	A	312	4/4	0.73	0.16	50,52,53,57	0
2	EDO	C	307	4/4	0.74	0.13	35,35,37,41	4
3	PGE	A	304	10/10	0.75	0.15	36,43,50,52	0
3	PGE	C	308	10/10	0.76	0.18	54,62,67,68	0
2	EDO	B	308	4/4	0.76	0.15	39,40,48,55	0
2	EDO	D	308	4/4	0.77	0.16	56,57,57,58	0
3	PGE	B	305	10/10	0.77	0.15	65,71,73,73	0
3	PGE	A	305	10/10	0.80	0.17	49,52,67,68	1
3	PGE	A	302	10/10	0.80	0.16	59,65,68,71	0
5	GOL	B	307	6/6	0.80	0.20	56,58,59,61	0
3	PGE	C	304	10/10	0.80	0.41	59,68,73,73	10
3	PGE	B	301	10/10	0.81	0.14	42,47,55,55	0
2	EDO	C	302	4/4	0.81	0.12	47,50,50,55	0
2	EDO	A	311	4/4	0.81	0.14	55,57,57,58	0
2	EDO	D	303	4/4	0.81	0.15	50,50,51,54	0
2	EDO	D	305	4/4	0.81	0.12	47,48,50,52	0
2	EDO	A	315	4/4	0.82	0.14	58,58,59,59	0
2	EDO	D	306	4/4	0.82	0.14	47,47,50,51	0
4	ACT	A	308	4/4	0.82	0.15	59,61,62,63	0
2	EDO	D	309	4/4	0.83	0.14	67,69,69,72	0
3	PGE	B	303	10/10	0.83	0.16	60,66,70,70	2
2	EDO	D	307	4/4	0.84	0.14	50,51,52,52	0
2	EDO	A	309	4/4	0.84	0.18	51,52,55,56	0
2	EDO	C	305	4/4	0.84	0.14	59,59,60,60	0
3	PGE	A	307	10/10	0.84	0.14	53,58,65,65	2
5	GOL	B	302	6/6	0.85	0.41	37,53,53,56	6
3	PGE	D	301	10/10	0.85	0.16	41,45,57,61	3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	PGE	B	306	10/10	0.85	0.13	52,60,69,69	0
2	EDO	A	303	4/4	0.86	0.13	54,54,57,57	0
2	EDO	D	304	4/4	0.88	0.13	53,57,60,65	0
6	PG4	C	301	13/13	0.88	0.13	36,43,54,56	3
6	PG4	A	313	13/13	0.89	0.12	30,44,54,55	0
2	EDO	C	303	4/4	0.89	0.11	43,43,44,50	0
2	EDO	C	306	4/4	0.90	0.10	46,46,48,51	0
2	EDO	B	309	4/4	0.92	0.10	48,49,50,51	0
2	EDO	A	306	4/4	0.92	0.12	30,34,35,36	0
2	EDO	A	301	4/4	0.94	0.09	42,43,43,48	0

6.5 Other polymers [i](#)

There are no such residues in this entry.