



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 05:34 AM UTC

PDB ID : 5LYH / pdb_00005lyh
Title : Human PARP14 (ARTD8), catalytic fragment in complex with inhibitor H10
Authors : Karlberg, T.; Thorsell, A.G.; Schuler, H.
Deposited on : 2016-09-28
Resolution : 2.17 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.17 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	8975 (2.20-2.16)
Clashscore	190562	9786 (2.20-2.16)
Ramachandran outliers	187476	9664 (2.20-2.16)
Sidechain outliers	187428	9664 (2.20-2.16)
RSRZ outliers	180081	8979 (2.20-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	193	 87% 8% . .
1	B	193	 3% 84% 7% . . 6%

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Poly [ADP-ribose] polymerase 14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	185	Total 1500	C 948	N 265	O 282	S 5	0	0	0
1	B	182	Total 1493	C 945	N 263	O 281	S 4	0	1	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1609	SER	-	expression tag	UNP Q460N5
A	1610	MET	-	expression tag	UNP Q460N5
B	1609	SER	-	expression tag	UNP Q460N5
B	1610	MET	-	expression tag	UNP Q460N5

- Molecule 2 is 3-[2-[4-[2-[4-[(3-aminocarbonylphenyl)amino]-4-oxidanylidene-butanoyl]amino]ethyl]-1,2,3-triazol-1-yl]ethylsulfamoyl]benzoic acid (CCD ID: 7B8) (formula: $C_{24}H_{27}N_7O_7S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			39	24	7	7	1		
2	B	1	Total	C	N	O	S	0	0
			39	24	7	7	1		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	10	Total	O	0	0
			10	10		
3	B	9	Total	O	0	0
			9	9		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Poly [ADP-ribose] polymerase 14

Chain A:  87% 8% . .



- Molecule 1: Poly [ADP-ribose] polymerase 14

Chain B:  3% 84% 7% . . 6%



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	83.03Å 83.03Å 122.08Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.50 – 2.17 46.50 – 2.17	Depositor EDS
% Data completeness (in resolution range)	99.7 (46.50-2.17) 99.9 (46.50-2.17)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.01 (at 2.16Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.195 , 0.218 0.196 , 0.221	Depositor DCC
R_{free} test set	2004 reflections (7.59%)	wwPDB-VP
Wilson B-factor (Å ²)	46.6	Xtriage
Anisotropy	0.568	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 59.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.40$, $\langle L^2 \rangle = 0.23$	Xtriage
Estimated twinning fraction	0.348 for -h,-k,l	Xtriage
Reported twinning fraction	0.658 for H, K, L 0.342 for -h,-k,l	Depositor
Outliers	0 of 26408 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3090	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 7B8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.93	1/1542 (0.1%)	1.04	2/2095 (0.1%)
1	B	0.98	2/1539 (0.1%)	1.08	4/2091 (0.2%)
All	All	0.96	3/3081 (0.1%)	1.06	6/4186 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1689	VAL	CA-CB	7.19	1.57	1.54
1	A	1689	VAL	CA-CB	6.34	1.57	1.54
1	B	1795	TYR	CA-C	-5.42	1.45	1.52

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1750	HIS	N-CA-C	9.70	122.86	111.71
1	A	1753	HIS	N-CA-C	6.01	118.32	111.11
1	B	1642	HIS	N-CA-C	5.79	119.81	112.87
1	A	1698	ASN	N-CA-C	5.18	121.84	110.80
1	B	1699	ARG	CA-C-N	5.04	130.76	121.70
1	B	1699	ARG	C-N-CA	5.04	130.76	121.70

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1612	MET	Peptide
1	B	1699	ARG	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1500	0	1425	3	0
1	B	1493	0	1414	8	0
2	A	39	0	0	0	0
2	B	39	0	0	0	0
3	A	10	0	0	0	0
3	B	9	0	0	0	0
All	All	3090	0	2839	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1628:GLU:OE1	1:A:1650:ARG:NH1	2.28	0.65
1:B:1699:ARG:CG	1:B:1700:SER:HA	2.29	0.62
1:B:1650:ARG:NH2	1:B:1652:GLN:OE1	2.33	0.60
1:B:1699:ARG:HG3	1:B:1700:SER:HA	1.90	0.54
1:B:1615:GLN:HG3	1:B:1655:ASP:HB2	1.94	0.49
1:B:1753:HIS:CG	1:B:1754:SER:N	2.87	0.43
1:A:1687:GLY:O	1:A:1690:PRO:HD2	2.19	0.42
1:B:1687:GLY:O	1:B:1690:PRO:HD2	2.19	0.42
1:B:1739:TYR:HA	1:B:1795:TYR:O	2.20	0.41
1:A:1679:GLN:HB3	1:A:1740:TYR:CZ	2.55	0.41
1:B:1699:ARG:HB3	1:B:1700:SER:CB	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	181/193 (94%)	176 (97%)	4 (2%)	1 (1%)	21	20
1	B	179/193 (93%)	167 (93%)	11 (6%)	1 (1%)	21	20
All	All	360/386 (93%)	343 (95%)	15 (4%)	2 (1%)	21	20

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1698	ASN
1	B	1753	HIS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	164/170 (96%)	157 (96%)	7 (4%)	26	32
1	B	164/170 (96%)	158 (96%)	6 (4%)	30	38
All	All	328/340 (96%)	315 (96%)	13 (4%)	28	35

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1615	GLN
1	A	1647	LYS
1	A	1663	LYS
1	A	1700	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1707	VAL
1	A	1762	ASN
1	A	1782	LEU
1	B	1615	GLN
1	B	1634	SER
1	B	1647	LYS
1	B	1700	SER
1	B	1720	ASN
1	B	1782	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1642	HIS
1	A	1691	HIS
1	A	1693	ASN
1	A	1695	ASN
1	A	1733	ASN
1	A	1765	ASN
1	A	1779	HIS
1	B	1642	HIS
1	B	1691	HIS
1	B	1693	ASN
1	B	1733	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	7B8	B	1901	-	41,41,41	3.13	12 (29%)	56,56,56	1.96	15 (26%)
2	7B8	A	1901	-	41,41,41	3.85	13 (31%)	56,56,56	2.22	18 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	7B8	B	1901	-	-	13/36/36/36	0/3/3/3
2	7B8	A	1901	-	-	17/36/36/36	0/3/3/3

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1901	7B8	SAY-NAX	14.90	1.83	1.61
2	B	1901	7B8	SAY-NAX	9.52	1.75	1.61
2	A	1901	7B8	OAE-SAY	9.01	1.54	1.43
2	B	1901	7B8	OAE-SAY	8.98	1.53	1.43
2	B	1901	7B8	OAZ-SAY	8.94	1.53	1.43
2	A	1901	7B8	OAZ-SAY	8.74	1.53	1.43
2	A	1901	7B8	CBG-SAY	-6.97	1.65	1.76
2	A	1901	7B8	CBI-CBL	-5.86	1.37	1.49
2	A	1901	7B8	NAU-NAD	-5.84	1.25	1.34
2	A	1901	7B8	CAH-CAG	-5.55	1.42	1.50
2	B	1901	7B8	CAH-CAG	-5.41	1.42	1.50
2	B	1901	7B8	CBI-CBL	-5.16	1.38	1.49
2	B	1901	7B8	NAU-NAD	-4.47	1.27	1.34
2	A	1901	7B8	CAJ-NAK	-3.95	1.33	1.41
2	A	1901	7B8	NAC-NAD	-3.72	1.25	1.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1901	7B8	NAC-NAD	-3.30	1.26	1.32
2	B	1901	7B8	CAN-CAO	3.12	1.57	1.51
2	B	1901	7B8	CAL-NAK	3.05	1.42	1.35
2	B	1901	7B8	CAR-CAS	2.87	1.56	1.50
2	A	1901	7B8	CAM-CAL	2.31	1.56	1.51
2	A	1901	7B8	CAT-CAS	-2.28	1.32	1.36
2	A	1901	7B8	CAR-CAS	2.26	1.54	1.50
2	B	1901	7B8	CAI-CAH	2.24	1.42	1.39
2	B	1901	7B8	CAT-CAS	-2.07	1.33	1.36
2	A	1901	7B8	OBK-CBL	-2.06	1.24	1.30

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1901	7B8	OAE-SAY-OAZ	-8.40	109.32	119.52
2	A	1901	7B8	OAE-SAY-OAZ	-7.51	110.40	119.52
2	A	1901	7B8	NAU-NAD-NAC	4.79	111.33	107.02
2	A	1901	7B8	OAZ-SAY-NAX	4.23	113.62	107.03
2	A	1901	7B8	CBJ-CBG-SAY	4.21	123.37	119.06
2	A	1901	7B8	OAF-CAG-CAH	4.00	124.49	119.60
2	B	1901	7B8	NAU-NAD-NAC	3.97	110.59	107.02
2	B	1901	7B8	OAF-CAG-CAH	3.69	124.11	119.60
2	B	1901	7B8	CBC-CAJ-CAI	-3.68	115.21	119.66
2	A	1901	7B8	CAQ-NAP-CAO	3.21	128.80	122.82
2	A	1901	7B8	OAE-SAY-NAX	3.18	111.97	107.03
2	B	1901	7B8	CAJ-NAK-CAL	-3.17	121.91	127.52
2	B	1901	7B8	CAH-CAI-CAJ	3.15	124.50	120.44
2	A	1901	7B8	OAF-CAG-NAA	-3.13	118.09	122.62
2	B	1901	7B8	OBD-CAL-CAM	-3.08	116.43	122.02
2	A	1901	7B8	OAZ-SAY-CBG	-3.04	104.15	107.98
2	A	1901	7B8	CAH-CAI-CAJ	2.98	124.27	120.44
2	A	1901	7B8	CBF-CBG-SAY	-2.97	116.49	119.76
2	A	1901	7B8	CBC-CAJ-NAK	2.87	130.04	120.41
2	A	1901	7B8	CAI-CAJ-NAK	-2.67	111.44	120.13
2	A	1901	7B8	OAE-SAY-CBG	2.51	111.15	107.98
2	A	1901	7B8	CAT-NAU-NAD	-2.37	108.44	110.75
2	B	1901	7B8	CAT-NAU-NAD	-2.29	108.53	110.75
2	B	1901	7B8	OBK-CBL-CBI	2.25	120.62	114.84
2	A	1901	7B8	CAM-CAL-NAK	-2.22	110.72	114.59
2	A	1901	7B8	CBA-CAH-CAI	-2.21	116.69	119.25
2	B	1901	7B8	CBB-CBC-CAJ	2.17	122.23	119.73
2	B	1901	7B8	OAF-CAG-NAA	-2.16	119.49	122.62

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1901	7B8	CAV-NAU-NAD	2.09	124.97	120.78
2	A	1901	7B8	OBD-CAL-NAK	2.07	127.36	123.64
2	B	1901	7B8	CAW-NAX-SAY	-2.06	114.07	120.27
2	B	1901	7B8	OAZ-SAY-NAX	2.02	110.18	107.03
2	B	1901	7B8	OBD-CAL-NAK	2.01	127.25	123.64

There are no chirality outliers.


All (30) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1901	7B8	NAU-CAV-CAW-NAX
2	A	1901	7B8	NAP-CAQ-CAR-CAS
2	B	1901	7B8	CAQ-CAR-CAS-NAC
2	A	1901	7B8	CAN-CAO-NAP-CAQ
2	B	1901	7B8	CAW-NAX-SAY-OAE
2	B	1901	7B8	CBJ-CBI-CBL-OBM
2	B	1901	7B8	CBH-CBI-CBL-OBK
2	B	1901	7B8	CBH-CBI-CBL-OBM
2	B	1901	7B8	CBJ-CBI-CBL-OBK
2	A	1901	7B8	OAB-CAO-NAP-CAQ
2	B	1901	7B8	CAW-NAX-SAY-CBG
2	A	1901	7B8	OBD-CAL-CAM-CAN
2	A	1901	7B8	CBF-CBG-SAY-OAZ
2	A	1901	7B8	CBJ-CBG-SAY-OAZ
2	A	1901	7B8	NAK-CAL-CAM-CAN
2	A	1901	7B8	CBJ-CBG-SAY-NAX
2	A	1901	7B8	CBF-CBG-SAY-NAX
2	B	1901	7B8	CAQ-CAR-CAS-CAT
2	A	1901	7B8	CBH-CBI-CBL-OBK
2	A	1901	7B8	CBJ-CBI-CBL-OBK
2	A	1901	7B8	CBH-CBI-CBL-OBM
2	A	1901	7B8	CBJ-CBI-CBL-OBM
2	A	1901	7B8	CAW-NAX-SAY-OAZ
2	B	1901	7B8	NAU-CAV-CAW-NAX
2	B	1901	7B8	CAW-CAV-NAU-CAT
2	B	1901	7B8	CBF-CBG-SAY-OAZ
2	B	1901	7B8	CBJ-CBG-SAY-OAZ
2	B	1901	7B8	CAW-CAV-NAU-NAD
2	A	1901	7B8	CAW-NAX-SAY-CBG
2	A	1901	7B8	CAW-NAX-SAY-OAE

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	185/193 (95%)	-0.57	0	100 100	34, 53, 89, 115	0
1	B	182/193 (94%)	-0.27	6 (3%)	49 48	34, 57, 126, 185	1 (0%)
All	All	367/386 (95%)	-0.42	6 (1%)	70 70	34, 54, 114, 185	1 (0%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1756	ILE	3.2
1	B	1701	TYR	3.1
1	B	1763	PRO	2.8
1	B	1765	ASN	2.6
1	B	1758	PRO	2.3
1	B	1766	PRO	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	7B8	A	1901	39/39	0.97	0.09	35,86,116,122	0
2	7B8	B	1901	39/39	0.98	0.08	38,85,126,132	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.