



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 05:19 PM UTC

PDB ID : 4LZK / pdb_00004lzk
Title : Crystal structure of inclusion body protein (PixA pfam12306) from Burkholderia cenocepacia J2315
Authors : Nocek, B.; Li, H.; Clancy, S.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2013-07-31
Resolution : 1.89 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	187	0% 76% 6% • 15%
1	B	187	2% 76% 13% 11%
1	C	187	3% 80% 11% • 7%
1	D	187	2% 75% 10% 16%
1	E	187	2% 72% 12% • 16%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	187	3% 76% 10% 14%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7932 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PixA inclusion body protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	159	Total	C	N	O	Se	0	1	0
			1191	761	199	229	2			
1	B	166	Total	C	N	O	Se	0	3	0
			1253	799	210	241	3			
1	C	174	Total	C	N	O	Se	0	0	0
			1276	812	214	248	2			
1	D	158	Total	C	N	O	Se	0	1	0
			1186	759	201	224	2			
1	E	157	Total	C	N	O	Se	0	0	0
			1181	752	200	227	2			
1	F	161	Total	C	N	O	Se	0	1	0
			1192	761	199	230	2			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP B4EII9
A	-1	ASN	-	expression tag	UNP B4EII9
A	0	ALA	-	expression tag	UNP B4EII9
B	-2	SER	-	expression tag	UNP B4EII9
B	-1	ASN	-	expression tag	UNP B4EII9
B	0	ALA	-	expression tag	UNP B4EII9
C	-2	SER	-	expression tag	UNP B4EII9
C	-1	ASN	-	expression tag	UNP B4EII9
C	0	ALA	-	expression tag	UNP B4EII9
D	-2	SER	-	expression tag	UNP B4EII9
D	-1	ASN	-	expression tag	UNP B4EII9
D	0	ALA	-	expression tag	UNP B4EII9
E	-2	SER	-	expression tag	UNP B4EII9
E	-1	ASN	-	expression tag	UNP B4EII9
E	0	ALA	-	expression tag	UNP B4EII9
F	-2	SER	-	expression tag	UNP B4EII9
F	-1	ASN	-	expression tag	UNP B4EII9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	0	ALA	-	expression tag	UNP B4EII9


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	116	Total 116	O 116	0	0
2	B	102	Total 102	O 102	0	0
2	C	136	Total 136	O 136	0	0
2	D	122	Total 122	O 122	0	0
2	E	98	Total 98	O 98	0	0
2	F	79	Total 79	O 79	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PixA inclusion body protein




- Molecule 1: PixA inclusion body protein




- Molecule 1: PixA inclusion body protein




- Molecule 1: PixA inclusion body protein



● Molecule 1: PixA inclusion body protein



● Molecule 1: PixA inclusion body protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	62.85Å 120.72Å 125.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	32.70 – 1.89 32.70 – 1.89	Depositor EDS
% Data completeness (in resolution range)	89.5 (32.70-1.89) 89.5 (32.70-1.89)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.00 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.176 , 0.212 0.181 , 0.218	Depositor DCC
R_{free} test set	3856 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	18.8	Xtriage
Anisotropy	0.124	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 46.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.025 for -h,l,k	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7932	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.36	1/1218 (0.1%)	1.16	3/1653 (0.2%)
1	B	1.33	4/1286 (0.3%)	1.08	0/1745
1	C	1.38	1/1301 (0.1%)	1.09	1/1769 (0.1%)
1	D	1.34	4/1213 (0.3%)	1.06	0/1647
1	E	1.25	0/1205	1.10	4/1635 (0.2%)
1	F	1.22	2/1220 (0.2%)	1.05	0/1659
All	All	1.32	12/7443 (0.2%)	1.09	8/10108 (0.1%)

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	107	VAL	C-O	-6.46	1.16	1.24
1	B	159	PHE	N-CA	-5.92	1.38	1.45
1	B	71	LEU	N-CA	-5.84	1.39	1.46
1	A	36	SER	N-CA	-5.78	1.39	1.46
1	D	43	THR	N-CA	5.70	1.53	1.46
1	D	169	LYS	C-O	-5.50	1.17	1.24
1	D	53	LEU	N-CA	-5.41	1.39	1.45
1	D	22	VAL	C-O	-5.38	1.18	1.24
1	B	72	LYS	CA-C	-5.31	1.46	1.52
1	F	142	ALA	CA-C	-5.26	1.46	1.52
1	F	25	THR	C-O	-5.15	1.18	1.24
1	B	154	TYR	CA-C	5.15	1.58	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	116	ILE	CB-CA-C	6.72	120.32	110.77
1	E	45	ILE	CA-C-N	6.49	126.41	119.85
1	E	45	ILE	C-N-CA	6.49	126.41	119.85
1	A	7	SER	CA-C-N	-6.23	115.07	123.10
1	A	7	SER	C-N-CA	-6.23	115.07	123.10
1	E	32	HIS	CA-C-N	5.39	125.01	119.56

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	32	HIS	C-N-CA	5.39	125.01	119.56
1	C	28	ILE	CB-CA-C	-5.25	105.25	111.97

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1191	0	1156	12	0
1	B	1253	0	1226	12	0
1	C	1276	0	1231	18	0
1	D	1186	0	1158	9	0
1	E	1181	0	1144	12	0
1	F	1192	0	1144	10	0
2	A	116	0	0	5	0
2	B	102	0	0	5	0
2	C	136	0	0	3	0
2	D	122	0	0	1	0
2	E	98	0	0	3	0
2	F	79	0	0	2	0
All	All	7932	0	7059	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:56:HIS:HE1	1:C:64:GLU:OE2	1.50	0.94
1:C:56:HIS:CE1	1:C:64:GLU:OE2	2.24	0.91
1:E:166:GLN:HG2	1:E:167:ASN:H	1.49	0.77
1:A:47:SER:HB3	2:A:314:HOH:O	1.85	0.76
1:A:47:SER:CB	2:A:314:HOH:O	2.34	0.73
1:E:165:SER:HB3	1:E:168:ARG:HG3	1.67	0.73

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:68:ASN:HB3	2:A:274:HOH:O	1.86	0.73
1:C:8:ILE:HD12	1:F:98:HIS:CD2	2.26	0.70
1:D:9:ASN:HB3	2:D:254:HOH:O	1.92	0.70
1:B:106:GLN:NE2	2:B:286:HOH:O	2.27	0.66
1:C:8:ILE:HB	1:F:100:GLN:HE22	1.61	0.65
1:E:100:GLN:HE21	1:F:10:LEU:HD12	1.63	0.64
1:E:141:GLN:NE2	2:E:258:HOH:O	2.31	0.64
1:B:98:HIS:CD2	1:D:8:ILE:HD12	2.34	0.62
1:E:166:GLN:CG	1:E:167:ASN:H	2.17	0.58
1:C:48:THR:HG21	1:D:27:TYR:OH	2.05	0.56
1:E:112:THR:HG23	2:E:224:HOH:O	2.06	0.56
1:B:38:ASN:OD1	1:B:41:ASN:ND2	2.39	0.55
1:C:165:SER:O	1:C:166:GLN:HG3	2.07	0.54
1:B:53:LEU:HD11	1:F:53:LEU:HD11	1.89	0.54
1:B:18:GLN:NE2	2:B:251:HOH:O	2.26	0.54
1:A:73:LEU:C	1:A:73:LEU:HD12	2.33	0.54
1:C:28:ILE:HD11	1:C:49:ALA:HB1	1.90	0.53
1:C:18:GLN:HG2	2:C:262:HOH:O	2.07	0.53
1:B:28[A]:ILE:HD11	1:B:49:ALA:HB1	1.91	0.52
1:B:112[A]:THR:HG23	2:B:216:HOH:O	2.09	0.52
1:E:155:LEU:HD12	1:E:179:TRP:CZ2	2.45	0.51
1:B:56:HIS:CD2	2:F:279:HOH:O	2.65	0.49
1:E:86:LEU:HG	2:E:292:HOH:O	2.13	0.48
1:C:17:ILE:HD12	1:C:73:LEU:HD12	1.95	0.48
1:C:8:ILE:HD12	1:F:98:HIS:HD2	1.75	0.47
1:D:28[A]:ILE:HD11	1:D:49:ALA:HB1	1.96	0.47
1:A:118:GLN:C	2:A:301:HOH:O	2.58	0.46
1:B:0:ALA:CB	2:B:293:HOH:O	2.63	0.46
1:E:166:GLN:HE22	1:E:168:ARG:HG2	1.81	0.46
1:A:28[A]:ILE:HD11	1:A:49:ALA:HB1	1.97	0.46
1:D:28[A]:ILE:HA	1:D:28[A]:ILE:HD13	1.71	0.45
1:A:73:LEU:C	1:A:73:LEU:CD1	2.90	0.45
1:A:116:ILE:HG12	1:C:90:SER:HB2	1.99	0.44
1:B:92:ASP:OD2	2:B:246:HOH:O	2.21	0.44
1:E:103:SER:HB2	1:E:154:TYR:HB2	1.99	0.44
1:D:117:GLU:O	1:E:164:ARG:NH2	2.44	0.44
1:C:138:GLN:OE1	1:C:140:PHE:CE1	2.71	0.43
1:A:95:LEU:HD13	1:C:133:THR:HG21	2.00	0.43
1:C:17:ILE:CD1	1:C:73:LEU:HD12	2.48	0.43
1:D:28[A]:ILE:HD11	1:D:49:ALA:CB	2.48	0.43
1:F:92:ASP:OD1	2:F:241:HOH:O	2.22	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:88:ASP:OD1	1:C:88:ASP:OD1	2.37	0.43
1:F:164:ARG:HA	1:F:168:ARG:O	2.19	0.43
1:F:38:ASN:ND2	1:F:40:GLN:H	2.18	0.42
1:A:8:ILE:HD11	1:D:158:SER:OG	2.20	0.42
1:B:3:VAL:HG21	1:C:128:PRO:HB3	2.01	0.42
2:C:301:HOH:O	1:F:104:GLY:HA2	2.19	0.42
1:F:28:ILE:HD11	1:F:49:ALA:HB1	2.02	0.42
1:C:68:ASN:ND2	1:C:180:GLN:OE1	2.43	0.41
1:E:166:GLN:NE2	1:E:168:ARG:HG2	2.36	0.41
1:A:112:THR:HG23	2:A:244:HOH:O	2.21	0.41
1:B:73:LEU:N	1:B:73:LEU:HD23	2.36	0.41
1:D:68:ASN:OD1	1:D:180:GLN:HG3	2.20	0.40
1:C:18:GLN:NE2	2:C:313:HOH:O	2.48	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	156/187 (83%)	154 (99%)	2 (1%)	0	100	100
1	B	165/187 (88%)	163 (99%)	2 (1%)	0	100	100
1	C	170/187 (91%)	166 (98%)	3 (2%)	1 (1%)	21	13
1	D	155/187 (83%)	151 (97%)	4 (3%)	0	100	100
1	E	153/187 (82%)	149 (97%)	3 (2%)	1 (1%)	18	10
1	F	158/187 (84%)	152 (96%)	6 (4%)	0	100	100
All	All	957/1122 (85%)	935 (98%)	20 (2%)	2 (0%)	43	36

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	165	SER
1	C	69	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	124/143 (87%)	117 (94%)	7 (6%)	19	11
1	B	132/143 (92%)	125 (95%)	7 (5%)	20	12
1	C	129/143 (90%)	121 (94%)	8 (6%)	16	9
1	D	123/143 (86%)	118 (96%)	5 (4%)	27	19
1	E	123/143 (86%)	115 (94%)	8 (6%)	15	8
1	F	123/143 (86%)	119 (97%)	4 (3%)	33	26
All	All	754/858 (88%)	715 (95%)	39 (5%)	21	13

All (39) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	ILE
1	A	11	GLU
1	A	68	ASN
1	A	73	LEU
1	A	116	ILE
1	A	148	LYS
1	A	165	SER
1	B	48[A]	THR
1	B	48[B]	THR
1	B	50	LEU
1	B	68	ASN
1	B	114	VAL
1	B	116	ILE
1	B	149	SER
1	C	41	ASN
1	C	48	THR
1	C	50	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	73	LEU
1	C	138	GLN
1	C	166	GLN
1	C	169	LYS
1	C	170	SER
1	D	73	LEU
1	D	106	GLN
1	D	114	VAL
1	D	165	SER
1	D	166	GLN
1	E	11	GLU
1	E	48	THR
1	E	50	LEU
1	E	63	SER
1	E	68	ASN
1	E	73	LEU
1	E	114	VAL
1	E	117	GLU
1	F	30	ARG
1	F	63	SER
1	F	68	ASN
1	F	170	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	34	ASN
1	A	74	ASN
1	A	98	HIS
1	A	100	GLN
1	B	74	ASN
1	C	15	GLN
1	C	40	GLN
1	C	56	HIS
1	D	34	ASN
1	D	74	ASN
1	D	98	HIS
1	E	15	GLN
1	E	34	ASN
1	E	40	GLN
1	F	18	GLN
1	F	40	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	F	74	ASN
1	F	100	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	157/187 (83%)	-0.17	2 (1%) 75 78	9, 17, 35, 55	1 (0%)
1	B	163/187 (87%)	-0.04	4 (2%) 58 62	11, 18, 38, 63	3 (1%)
1	C	172/187 (91%)	0.10	6 (3%) 47 50	11, 20, 44, 59	0
1	D	156/187 (83%)	-0.06	4 (2%) 57 61	10, 17, 41, 67	1 (0%)
1	E	155/187 (82%)	0.14	3 (1%) 66 70	12, 22, 41, 83	0
1	F	159/187 (85%)	0.28	6 (3%) 44 47	13, 26, 52, 86	1 (0%)
All	All	962/1122 (85%)	0.04	25 (2%) 57 61	9, 20, 44, 86	6 (0%)

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	116	ILE	4.4
1	C	165	SER	3.5
1	E	8	ILE	3.3
1	A	138	GLN	3.2
1	B	0	ALA	3.1
1	D	138	GLN	2.9
1	F	5	PRO	2.9
1	F	116	ILE	2.8
1	D	165	SER	2.7
1	C	132	ALA	2.6
1	F	168	ARG	2.5
1	B	9	ASN	2.4
1	A	165	SER	2.4
1	D	7	SER	2.4
1	C	166	GLN	2.4
1	F	10	LEU	2.3
1	D	168	ARG	2.3
1	E	138	GLN	2.2
1	C	168	ARG	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	133	THR	2.1
1	F	7	SER	2.1
1	C	64	GLU	2.1
1	B	138	GLN	2.1
1	E	166	GLN	2.1
1	F	27	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.