



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 26, 2026 – 11:35 AM UTC

PDB ID : 5M02 / pdb_00005m02
Title : Crystal structure of murine P14 TCR / H-2Db with PF, modified gp33 peptide from LCMV
Authors : Achour, A.; Sandalova, T.; Sun, R.
Deposited on : 2016-10-03
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	6% 91% 8%
2	B	119	7% 79% 6% 15%
3	G	205	21% 84% 7% 9%
4	H	238	6% 95% 5%
5	P	9	78% 22%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6998 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called H-2 class I histocompatibility antigen, D-B alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	275	Total	C	N	O	S	0	8	0
			2315	1463	411	432	9			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	101	Total	C	N	O	S	0	3	0
			853	545	145	155	8			

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-19	MET	-	initiating methionine	UNP P01887
B	-18	ALA	-	expression tag	UNP P01887
B	-17	ARG	-	expression tag	UNP P01887
B	-16	SER	-	expression tag	UNP P01887
B	-15	VAL	-	expression tag	UNP P01887
B	-14	THR	-	expression tag	UNP P01887
B	-13	LEU	-	expression tag	UNP P01887
B	-12	VAL	-	expression tag	UNP P01887
B	-11	PHE	-	expression tag	UNP P01887
B	-10	LEU	-	expression tag	UNP P01887
B	-9	VAL	-	expression tag	UNP P01887
B	-8	LEU	-	expression tag	UNP P01887
B	-7	VAL	-	expression tag	UNP P01887
B	-6	SER	-	expression tag	UNP P01887
B	-5	LEU	-	expression tag	UNP P01887
B	-4	THR	-	expression tag	UNP P01887
B	-3	GLY	-	expression tag	UNP P01887
B	-2	LEU	-	expression tag	UNP P01887
B	-1	MET	-	expression tag	UNP P01887
B	0	GLY	-	expression tag	UNP P01887

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	85	ASP	ALA	variant	UNP P01887

- Molecule 3 is a protein called Protein Trav14-1,T-cell receptor alpha chain C region.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	187	Total	C	N	O	S	0	0	0
			1467	933	235	291	8			

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	100	LEU	-	linker	UNP A0A0G2JF94
G	101	TYR	-	linker	UNP A0A0G2JF94
G	102	GLY	-	linker	UNP A0A0G2JF94
G	103	ASN	-	linker	UNP A0A0G2JF94
G	104	GLU	-	linker	UNP A0A0G2JF94
G	105	LYS	-	linker	UNP A0A0G2JF94
G	106	ILE	-	linker	UNP A0A0G2JF94
G	107	THR	-	linker	UNP A0A0G2JF94
G	108	PHE	-	linker	UNP A0A0G2JF94
G	109	GLY	-	linker	UNP A0A0G2JF94
G	110	ALA	-	linker	UNP A0A0G2JF94
G	111	GLY	-	linker	UNP A0A0G2JF94
G	112	THR	-	linker	UNP A0A0G2JF94
G	113	LYS	-	linker	UNP A0A0G2JF94
G	114	LEU	-	linker	UNP A0A0G2JF94
G	115	THR	-	linker	UNP A0A0G2JF94
G	116	ILE	-	linker	UNP A0A0G2JF94
G	117	LYS	-	linker	UNP A0A0G2JF94
G	118	PRO	-	linker	UNP A0A0G2JF94
G	119	ASN	-	linker	UNP A0A0G2JF94
G	166	CYS	THR	conflict	UNP P01849

- Molecule 4 is a protein called T-cell receptor beta chain V region C5,T-cell receptor beta-2 chain C region.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	H	237	Total	C	N	O	S	0	1	0
			1881	1184	333	358	6			

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	93	ASP	-	linker	UNP P04213
H	94	ALA	-	linker	UNP P04213
H	95	GLY	-	linker	UNP P04213
H	96	GLY	-	linker	UNP P04213
H	97	ARG	-	linker	UNP P04213
H	98	ASN	-	linker	UNP P04213
H	99	THR	-	linker	UNP P04213
H	100	LEU	-	linker	UNP P04213
H	101	TYR	-	linker	UNP P04213
H	102	PHE	-	linker	UNP P04213
H	103	GLY	-	linker	UNP P04213
H	104	ALA	-	linker	UNP P04213
H	105	GLY	-	linker	UNP P04213
H	106	THR	-	linker	UNP P04213
H	107	ARG	-	linker	UNP P04213
H	108	LEU	-	linker	UNP P04213
H	109	SER	-	linker	UNP P04213
H	110	VAL	-	linker	UNP P04213
H	111	LEU	-	linker	UNP P04213
H	168	CYS	SER	conflict	UNP P01851
H	182	SER	CYS	conflict	UNP P01851

- Molecule 5 is a protein called LCMV-DERIVED GP33 ALTERED PEPTIDE LIGAND PF.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	P	9	Total	C	N	O	S	0	0	0
			72	48	11	12	1			

- Molecule 6 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	H	1	Total	C	O	0	0
			6	3	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	138	Total	O	0	0
			138	138		
7	B	56	Total	O	0	0
			56	56		
7	G	63	Total	O	0	0
			63	63		
7	H	131	Total	O	0	0
			131	131		
7	P	4	Total	O	0	0
			4	4		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: H-2 class I histocompatibility antigen, D-B alpha chain




- Molecule 2: Beta-2-microglobulin



- Molecule 3: Protein Trav14-1,T-cell receptor alpha chain C region



- Molecule 4: T-cell receptor beta chain V region C5,T-cell receptor beta-2 chain C region



- Molecule 5: LCMV-DERIVED GP33 ALTERED PEPTIDE LIGAND PF

Chain P:



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	255.11Å 46.50Å 89.02Å 90.00° 94.60° 90.00°	Depositor
Resolution (Å)	127.14 – 1.75 127.14 – 1.75	Depositor EDS
% Data completeness (in resolution range)	98.1 (127.14-1.75) 98.1 (127.14-1.75)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.50 (at 1.75Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.188 , 0.214 0.197 , 0.221	Depositor DCC
R_{free} test set	5198 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	27.4	Xtriage
Anisotropy	0.326	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 33.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6998	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	0/2405	0.87	4/3259 (0.1%)
2	B	0.69	0/888	0.83	0/1200
3	G	0.88	5/1502 (0.3%)	0.92	8/2037 (0.4%)
4	H	0.71	0/1937	0.81	0/2633
5	P	0.55	0/74	0.82	0/97
All	All	0.76	5/6806 (0.1%)	0.86	12/9226 (0.1%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	G	122	ASN	C-N	5.39	1.40	1.33
3	G	133	PRO	N-CD	5.31	1.55	1.47
3	G	124	GLU	C-N	5.31	1.40	1.33
3	G	125	PRO	N-CD	5.18	1.54	1.47
3	G	123	PRO	N-CD	5.03	1.54	1.47

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	234	ARG	NE-CZ-NH1	10.76	132.26	121.50
1	A	234	ARG	NE-CZ-NH2	-10.04	110.17	119.20
1	A	234	ARG	CD-NE-CZ	6.94	134.11	124.40
1	A	234	ARG	CB-CG-CD	6.15	125.45	111.30
3	G	132	ASP	CA-C-N	-6.11	113.52	119.87
3	G	132	ASP	C-N-CA	-6.11	113.52	119.87
3	G	122	ASN	CA-C-N	-5.86	113.67	119.99
3	G	122	ASN	C-N-CA	-5.86	113.67	119.99
3	G	124	GLU	CA-C-N	-5.62	114.17	119.85
3	G	124	GLU	C-N-CA	-5.62	114.17	119.85
3	G	132	ASP	O-C-N	5.40	125.60	121.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	88	GLN	CB-CA-C	5.19	117.11	109.22

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2315	0	2206	21	0
2	B	853	0	842	8	0
3	G	1467	0	1412	0	0
4	H	1881	0	1792	6	0
5	P	72	0	72	4	0
6	A	12	0	16	0	0
6	H	6	0	8	0	0
7	A	138	0	0	0	0
7	B	56	0	0	0	0
7	G	63	0	0	0	0
7	H	131	0	0	0	0
7	P	4	0	0	0	0
All	All	6998	0	6348	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:234:ARG:HD3	2:B:10:TYR:CE2	2.21	0.75
1:A:111[B]:ARG:HG2	1:A:111[B]:ARG:HH11	1.55	0.71
1:A:97:GLN:HE22	5:P:5:ASN:HD21	1.40	0.67
1:A:96:GLN:OE1	2:B:31:HIS:HE1	1.84	0.61
1:A:234:ARG:HD3	2:B:10:TYR:CZ	2.36	0.60
1:A:263:HIS:CD2	1:A:265:GLY:H	2.20	0.60
2:B:31:HIS:CD2	2:B:32:PRO:HA	2.37	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:H:63:LYS:HG2	4:H:75:ILE:HB	1.87	0.56
1:A:111[B]:ARG:HH11	1:A:111[B]:ARG:CG	2.20	0.54
1:A:137:ASP:OD1	2:B:-1:MET:HE3	2.11	0.51
1:A:70:GLN:NE2	5:P:5:ASN:H	2.10	0.50
1:A:234:ARG:HD2	1:A:242:GLN:OE1	2.12	0.48
1:A:70:GLN:HE22	5:P:5:ASN:H	1.60	0.48
1:A:72:GLN:OE1	4:H:48[A]:TYR:CD1	2.68	0.47
1:A:210:PRO:O	1:A:263:HIS:HE1	1.96	0.47
4:H:23:GLN:HG2	4:H:30:MET:HE3	1.96	0.46
1:A:121:ARG:CD	2:B:-1:MET:HE2	2.46	0.45
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.50	0.45
1:A:263:HIS:HD2	1:A:265:GLY:H	1.64	0.45
1:A:11:ALA:HA	1:A:21:ARG:O	2.16	0.45
1:A:234:ARG:HD2	1:A:242:GLN:HB2	1.98	0.45
4:H:19:LEU:HD22	4:H:106:THR:HG21	1.99	0.44
4:H:143:LEU:HG	4:H:145:ARG:HG3	2.00	0.44
4:H:29:TYR:CZ	4:H:48[B]:TYR:HD1	2.36	0.43
1:A:111[B]:ARG:CG	1:A:111[B]:ARG:NH1	2.80	0.42
1:A:155:HIS:HB3	5:P:6:PHE:CZ	2.54	0.42
2:B:40:LEU:HD11	2:B:81[A]:ARG:HE	1.84	0.42
1:A:260:ARG:NE	1:A:271:THR:HG22	2.35	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	279/276 (101%)	274 (98%)	5 (2%)	0	100	100
2	B	102/119 (86%)	101 (99%)	1 (1%)	0	100	100
3	G	185/205 (90%)	178 (96%)	7 (4%)	0	100	100
4	H	236/238 (99%)	233 (99%)	3 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
5	P	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
All	All	809/847 (96%)	792 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	241/234 (103%)	235 (98%)	6 (2%)	42	22
2	B	98/111 (88%)	98 (100%)	0	100	100
3	G	166/184 (90%)	159 (96%)	7 (4%)	26	8
4	H	204/204 (100%)	203 (100%)	1 (0%)	81	75
5	P	7/7 (100%)	7 (100%)	0	100	100
All	All	716/740 (97%)	702 (98%)	14 (2%)	48	29

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	86	ASN
1	A	114	LEU
1	A	225	THR
1	A	226	GLN
1	A	228	MET
1	A	272	LEU
3	G	14	SER
3	G	75	LYS
3	G	100	LEU
3	G	135	SER
3	G	137	ASP
3	G	157	GLU
3	G	160	THR
4	H	28	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	70	GLN
1	A	127	ASN
1	A	263	HIS
2	B	8	GLN
2	B	31	HIS
2	B	67	HIS
3	G	13	GLN
3	G	83	HIS
3	G	121	GLN
3	G	136	GLN
3	G	149	GLN
4	H	22	HIS
4	H	26	ASN
4	H	39	HIS
4	H	164	HIS
4	H	195	HIS
5	P	5	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	GOL	A	302	-	5,5,5	0.41	0	5,5,5	0.24	0
6	GOL	H	301	-	5,5,5	0.32	0	5,5,5	0.41	0
6	GOL	A	301	-	5,5,5	0.31	0	5,5,5	0.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	302	-	-	0/4/4/4	-
6	GOL	H	301	-	-	3/4/4/4	-
6	GOL	A	301	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	301	GOL	O1-C1-C2-C3
6	H	301	GOL	O2-C2-C3-O3
6	H	301	GOL	C1-C2-C3-O3
6	A	301	GOL	O1-C1-C2-O2
6	H	301	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	275/276 (99%)	0.29	17 (6%) 26 30	16, 28, 47, 70	8 (2%)
2	B	101/119 (84%)	0.39	8 (7%) 18 22	16, 32, 49, 60	3 (2%)
3	G	187/205 (91%)	1.23	43 (22%) 2 2	24, 44, 84, 113	0
4	H	237/238 (99%)	0.34	14 (5%) 28 32	20, 30, 50, 68	1 (0%)
5	P	9/9 (100%)	-0.35	0 100 100	20, 21, 25, 25	0
All	All	809/847 (95%)	0.53	82 (10%) 12 14	16, 32, 62, 113	12 (1%)

All (82) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	G	189	PHE	7.7
3	G	172	ALA	7.3
2	B	-1	MET	6.1
4	H	48[A]	TYR	5.2
3	G	21	TRP	4.9
3	G	30	CYS	4.9
3	G	187	THR	4.7
4	H	174	TYR	4.6
3	G	173	MET	4.4
3	G	150	ILE	4.4
1	A	228	MET	4.3
3	G	175	SER	4.1
3	G	190	THR	4.1
1	A	225	THR	3.9
3	G	20	VAL	3.7
4	H	237	ASP	3.7
3	G	166	CYS	3.7
2	B	0	GLY	3.6
4	H	219	GLY	3.6
3	G	167	VAL	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	2	PRO	3.5
1	A	1	GLY	3.5
3	G	119	ASN	3.4
3	G	188	SER	3.4
3	G	124	GLU	3.2
2	B	48	LYS	3.1
1	A	51	TRP	3.1
4	H	213	GLU	3.1
3	G	194	ILE	3.0
3	G	25	THR	3.0
3	G	156	MET	2.9
2	B	75	THR	2.8
3	G	160	THR	2.8
3	G	195	PHE	2.8
3	G	88	GLN	2.8
3	G	192	GLN	2.8
4	H	236	ALA	2.8
3	G	158	SER	2.8
1	A	181	ARG	2.8
3	G	149	GLN	2.8
1	A	230	LEU	2.7
3	G	86	ASP	2.7
1	A	90	GLY	2.6
3	G	196	LYS	2.6
3	G	168	LEU	2.6
4	H	168	CYS	2.6
4	H	218	GLU	2.5
3	G	141	CYS	2.5
2	B	47	PRO	2.5
3	G	118	PRO	2.5
1	A	227	ASP	2.5
1	A	107	TRP	2.4
1	A	50	PRO	2.4
1	A	89	ALA	2.4
1	A	271	THR	2.4
3	G	120	ILE	2.4
3	G	151	ASN	2.3
1	A	252	GLY	2.3
4	H	203	CYS	2.3
4	H	235	ARG	2.3
3	G	153	PRO	2.2
3	G	186	GLN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	G	155	THR	2.2
4	H	178	ASN	2.2
4	H	212	GLU	2.2
1	A	196	LYS	2.2
4	H	220	SER	2.2
3	G	174	ASP	2.2
1	A	145[A]	ARG	2.2
3	G	171	LYS	2.2
3	G	80	LEU	2.1
3	G	183	TRP	2.1
1	A	194	ARG	2.1
3	G	137	ASP	2.1
2	B	77	THR	2.1
3	G	115	THR	2.1
2	B	20	PRO	2.1
3	G	122	ASN	2.0
3	G	60	VAL	2.0
4	H	89	CYS	2.0
3	G	161	PHE	2.0
2	B	34	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	GOL	H	301	6/6	0.57	0.30	64,67,68,72	0
6	GOL	A	301	6/6	0.70	0.21	61,63,65,66	0
6	GOL	A	302	6/6	0.80	0.17	31,36,43,49	0

6.5 Other polymers ⓘ

There are no such residues in this entry.