



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 26, 2026 – 06:17 AM EDT

PDB ID : 9M00 / pdb_00009m00
Title : Crystal Structure of the Basal pilin EbpB from *Enterococcus faecalis* with a partially disordered N-terminal domain.
Authors : Sharma, V.; Krishnan, V.
Deposited on : 2025-02-23
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	436	
1	B	436	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4988 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Endocarditis and biofilm-associated pilus minor subunit EbpB..

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	312	Total	C	N	O	S	0	0	0
			2340	1481	395	462	2			
1	A	326	Total	C	N	O	S	0	0	0
			2469	1564	419	484	2			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	91	Total	O	0	0
			91	91		
2	A	88	Total	O	0	0
			88	88		

- Molecule 1: Endocarditis and biofilm-associated pilus minor subunit EbpB.



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	39.30Å 48.08Å 127.01Å 96.44° 98.04° 90.12°	Depositor
Resolution (Å)	43.06 – 2.70 43.06 – 2.70	Depositor EDS
% Data completeness (in resolution range)	89.3 (43.06-2.70) 89.3 (43.06-2.70)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.18 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.8.0419	Depositor
R, R_{free}	0.240 , 0.279 0.242 , 0.281	Depositor DCC
R_{free} test set	1183 reflections (4.71%)	wwPDB-VP
Wilson B-factor (Å ²)	21.0	Xtriage
Anisotropy	0.039	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 34.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.042 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.85	EDS
Total number of atoms	4988	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	0/2506	1.00	3/3389 (0.1%)
1	B	0.56	0/2372	1.03	3/3206 (0.1%)
All	All	0.55	0/4878	1.01	6/6595 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	256	THR	CA-CB-OG1	-6.49	99.87	109.60
1	B	267	ASP	CA-CB-CG	5.48	118.08	112.60
1	B	180	ASN	CB-CG-OD1	-5.35	110.11	120.80
1	A	191	GLU	CB-CG-CD	5.15	121.35	112.60
1	A	267	ASP	CA-CB-CG	5.12	117.72	112.60
1	A	331	GLU	CB-CG-CD	5.10	121.27	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2469	0	2363	4	0
1	B	2340	0	2219	7	0
2	A	88	0	0	0	0
2	B	91	0	0	0	0
All	All	4988	0	4582	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:340:ALA:HB2	1:B:397:PRO:HG3	1.87	0.56
1:A:340:ALA:HB2	1:A:397:PRO:HG3	1.88	0.55
1:B:115:THR:HA	1:B:119:GLU:O	2.07	0.53
1:B:40:LEU:HB2	1:B:66:LEU:HD22	1.94	0.50
1:B:345:LYS:HA	1:B:350:GLU:O	2.16	0.46
1:A:201:PHE:CD2	1:A:207:ILE:HG12	2.51	0.46
1:A:345:LYS:HA	1:A:350:GLU:O	2.16	0.46
1:B:81:TYR:O	1:B:82:GLN:CB	2.67	0.42
1:B:374:ASN:OD1	1:B:374:ASN:C	2.62	0.42
1:A:374:ASN:OD1	1:A:374:ASN:C	2.62	0.41
1:B:328:VAL:HB	1:B:333:ALA:HA	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	310/436 (71%)	302 (97%)	8 (3%)	0	100	100
1	B	292/436 (67%)	283 (97%)	9 (3%)	0	100	100
All	All	602/872 (69%)	585 (97%)	17 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	251/377 (67%)	248 (99%)	3 (1%)	63	84
1	B	236/377 (63%)	235 (100%)	1 (0%)	84	93
All	All	487/754 (65%)	483 (99%)	4 (1%)	73	88

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	371	ILE
1	A	38	LYS
1	A	197	LYS
1	A	371	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	156	ASN
1	B	198	GLN
1	B	208	ASN
1	B	260	GLN
1	B	286	GLN
1	B	388	GLN
1	A	198	GLN
1	A	208	ASN
1	A	286	GLN
1	A	388	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	326/436 (74%)	0.59	27 (8%) 17 14	10, 19, 51, 58	0
1	B	312/436 (71%)	0.60	33 (10%) 11 9	8, 17, 54, 69	0
All	All	638/872 (73%)	0.60	60 (9%) 14 12	8, 18, 53, 69	0

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	91	GLN	5.6
1	B	157	LEU	4.6
1	B	94	GLN	4.3
1	A	76	VAL	4.2
1	B	122	VAL	4.1
1	A	159	MET	4.1
1	B	141	PHE	3.9
1	B	37	HIS	3.8
1	B	44	GLY	3.6
1	B	86	GLU	3.6
1	B	97	LEU	3.4
1	B	177	TYR	3.3
1	B	196	ASP	3.3
1	B	120	ASP	3.2
1	B	116	ILE	3.2
1	B	96	GLN	3.0
1	B	63	TYR	3.0
1	B	114	GLN	3.0
1	B	404	GLN	2.9
1	B	36	LEU	2.9
1	A	109	ALA	2.9
1	B	93	ALA	2.8
1	A	177	TYR	2.8
1	B	66	LEU	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	138	ALA	2.7
1	A	355	THR	2.6
1	A	108	ILE	2.6
1	B	38	LYS	2.6
1	A	145	GLU	2.6
1	A	123	VAL	2.6
1	A	139	TYR	2.5
1	A	73	VAL	2.5
1	B	43	ASP	2.5
1	A	118	GLY	2.5
1	B	82	GLN	2.4
1	A	124	SER	2.4
1	A	38	LYS	2.3
1	A	99	GLU	2.3
1	B	73	VAL	2.3
1	A	98	ALA	2.2
1	B	74	TYR	2.2
1	A	37	HIS	2.2
1	B	92	GLU	2.2
1	B	81	TYR	2.2
1	B	185	TYR	2.2
1	A	96	GLN	2.2
1	A	43	ASP	2.2
1	A	119	GLU	2.2
1	A	74	TYR	2.2
1	B	72	GLN	2.1
1	A	404	GLN	2.1
1	B	121	GLY	2.1
1	A	102	ALA	2.1
1	B	418	THR	2.1
1	A	100	THR	2.1
1	B	85	SER	2.1
1	A	110	GLU	2.0
1	B	355	THR	2.0
1	A	144	ALA	2.0
1	A	39	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.