



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:22 PM UTC

PDB ID : 1M1E / pdb_00001m1e
Title : Beta-catenin armadillo repeat domain bound to ICAT
Authors : Daniels, D.L.; Weis, W.I.
Deposited on : 2002-06-18
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	538	
2	B	81	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4708 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-catenin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	512	Total	C	N	O	S	0	0	0
			3902	2451	708	717	26			

- Molecule 2 is a protein called ICAT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	65	Total	C	N	O	S	0	0	0
			496	305	85	102	4			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	286	Total	O	0	0
			286	286		
3	B	24	Total	O	0	0
			24	24		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Beta-catenin



• Molecule 2: ICAT



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	97.75Å 97.32Å 86.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.56 – 2.10 43.56 – 2.10	Depositor EDS
% Data completeness (in resolution range)	97.4 (43.56-2.10) 97.5 (43.56-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.31 (at 2.10Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.202 , 0.228 0.202 , 0.228	Depositor DCC
R_{free} test set	2415 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	26.8	Xtriage
Anisotropy	0.511	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 49.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.017 for k,h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4708	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/3958	0.81	3/5372 (0.1%)
2	B	0.39	0/499	0.81	0/668
All	All	0.39	0/4457	0.81	3/6040 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	389	SER	N-CA-C	7.46	120.35	111.33
1	A	474	ARG	N-CA-C	6.62	120.53	111.39
1	A	265	HIS	N-CA-C	5.78	121.20	114.04

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3902	0	4023	86	0
2	B	496	0	473	13	0
3	A	286	0	0	6	0
3	B	24	0	0	0	0
All	All	4708	0	4496	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:452:LEU:HD23	1:A:493:VAL:HG11	1.56	0.88
1:A:399:GLU:HG3	1:A:437:MET:SD	2.15	0.87
1:A:153:ILE:HB	1:A:154:PRO:HD3	1.58	0.85
1:A:164:ASP:O	1:A:167:VAL:HG12	1.79	0.82
1:A:668:GLN:HG3	1:A:669:ASP:H	1.46	0.80
1:A:160:LEU:HD12	1:A:194:MET:HE1	1.62	0.80
1:A:414:ILE:O	1:A:418:THR:HG23	1.83	0.79
3:A:773:HOH:O	2:B:64:ALA:HB1	1.85	0.74
1:A:354:LYS:HB3	1:A:355:PRO:HD3	1.72	0.72
1:A:515:ARG:C	1:A:515:ARG:HD2	2.17	0.69
1:A:165:GLN:O	1:A:168:VAL:HG22	1.93	0.68
1:A:546:ASP:HA	1:A:549:ARG:NH1	2.11	0.66
2:B:25:MET:HE3	2:B:28:LYS:HB3	1.80	0.64
1:A:418:THR:HG22	1:A:460:ILE:HG12	1.81	0.63
1:A:452:LEU:CD2	1:A:493:VAL:HG11	2.27	0.63
1:A:600:VAL:O	1:A:603:LEU:HB2	1.99	0.62
1:A:433:LYS:O	1:A:437:MET:HG3	2.01	0.61
1:A:194:MET:HE2	1:A:198:ILE:HG13	1.83	0.60
1:A:620:GLU:O	1:A:623:GLN:HG2	2.00	0.60
1:A:432:TYR:O	1:A:436:MET:HG2	2.02	0.60
1:A:475:HIS:CD2	1:A:477:GLU:H	2.20	0.60
1:A:670:TYR:O	1:A:671:LYS:HB2	2.01	0.60
1:A:208:VAL:HG23	1:A:248:VAL:HG21	1.85	0.59
1:A:434:ASN:HA	1:A:437:MET:HE3	1.85	0.59
2:B:34:THR:OG1	2:B:37:GLU:HG3	2.03	0.58
1:A:425:SER:OG	2:B:69:MET:HE2	2.04	0.58
1:A:189:MET:HE1	1:A:224:HIS:CD2	2.39	0.57
1:A:546:ASP:HA	1:A:549:ARG:HH11	1.67	0.57
1:A:151:ARG:HG3	1:A:152:ALA:H	1.70	0.57
1:A:308:ASN:HA	2:B:78:ASP:HA	1.86	0.57
1:A:151:ARG:HG3	1:A:152:ALA:N	2.20	0.56
2:B:69:MET:HE3	2:B:71:PHE:CZ	2.41	0.56
1:A:418:THR:HG22	1:A:460:ILE:CG1	2.36	0.55
1:A:208:VAL:HG13	1:A:209:GLU:N	2.22	0.55
1:A:515:ARG:HD2	1:A:515:ARG:O	2.07	0.53
2:B:69:MET:HE3	2:B:71:PHE:HZ	1.73	0.53
2:B:26:LEU:HA	2:B:29:MET:HE2	1.89	0.53
1:A:243:MET:HE3	1:A:251:VAL:HG11	1.90	0.53
1:A:666:LYS:N	1:A:667:PRO:CD	2.72	0.53
1:A:474:ARG:HH11	1:A:474:ARG:HG3	1.73	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:543:ALA:O	1:A:547:THR:HG23	2.09	0.53
1:A:287:ASN:HD22	1:A:287:ASN:H	1.58	0.52
1:A:196:SER:O	1:A:200:ARG:HG3	2.09	0.52
1:A:471:LEU:O	1:A:475:HIS:HE1	1.92	0.52
1:A:233:LYS:NZ	1:A:233:LYS:HB3	2.25	0.51
1:A:596:ILE:HG22	1:A:634:GLU:HG3	1.92	0.51
1:A:666:LYS:H	1:A:667:PRO:CD	2.23	0.51
1:A:287:ASN:HD22	1:A:287:ASN:N	2.09	0.50
1:A:515:ARG:HD3	1:A:579:ILE:HG13	1.92	0.50
1:A:664:GLU:O	1:A:665:ASP:C	2.54	0.50
1:A:666:LYS:O	1:A:667:PRO:C	2.55	0.49
1:A:168:VAL:HG23	1:A:169:ASN:N	2.27	0.49
2:B:25:MET:O	2:B:29:MET:HG3	2.13	0.49
1:A:309:GLN:HE21	1:A:313:LEU:HG	1.78	0.48
1:A:565:ARG:HB3	1:A:567:GLU:OE1	2.13	0.48
1:A:612:ARG:NH1	3:A:735:HOH:O	2.43	0.48
1:A:499:HIS:HD2	3:A:945:HOH:O	1.97	0.48
1:A:473:SER:OG	1:A:474:ARG:HG3	2.14	0.47
1:A:195:VAL:O	1:A:199:VAL:HG23	2.14	0.47
1:A:260:HIS:CE1	1:A:264:LEU:HD11	2.50	0.47
1:A:194:MET:HE3	1:A:194:MET:HA	1.95	0.47
1:A:600:VAL:HA	1:A:603:LEU:HD22	1.95	0.47
1:A:171:ALA:O	1:A:175:VAL:HG23	2.15	0.46
1:A:546:ASP:OD2	1:A:549:ARG:NH1	2.49	0.46
1:A:186:HIS:HA	1:A:189:MET:HG2	1.98	0.45
1:A:666:LYS:N	1:A:667:PRO:HD2	2.32	0.45
1:A:474:ARG:NH1	2:B:64:ALA:O	2.50	0.45
1:A:194:MET:HE2	1:A:198:ILE:CG1	2.46	0.45
1:A:399:GLU:H	1:A:399:GLU:CD	2.25	0.45
1:A:370:LEU:HB2	1:A:404:THR:HG21	1.99	0.45
1:A:541:VAL:O	1:A:545:GLN:HG3	2.17	0.45
1:A:436:MET:SD	1:A:477:GLU:HG2	2.57	0.45
1:A:431:ASN:O	1:A:435:LYS:HG3	2.17	0.45
1:A:625:LYS:HG2	3:A:794:HOH:O	2.18	0.44
1:A:169:ASN:HD21	1:A:207:ASP:CB	2.30	0.44
1:A:270:LYS:O	1:A:274:ARG:HG3	2.17	0.44
1:A:153:ILE:HB	1:A:154:PRO:CD	2.38	0.44
1:A:525:ALA:O	1:A:529:GLU:HG3	2.17	0.44
1:A:237:ILE:HB	1:A:238:PRO:HD3	2.00	0.43
1:A:160:LEU:CD1	1:A:194:MET:HE1	2.41	0.43
1:A:515:ARG:HG2	1:A:576:ALA:HA	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:463:PRO:HB3	2:B:69:MET:HE1	2.00	0.42
2:B:25:MET:HE2	2:B:29:MET:HG3	2.00	0.42
1:A:428:THR:O	1:A:435:LYS:HE2	2.19	0.42
1:A:379:GLN:NE2	1:A:415:ASN:HB3	2.35	0.41
1:A:567:GLU:H	1:A:567:GLU:CD	2.29	0.41
1:A:239:ALA:O	1:A:243:MET:HG2	2.21	0.41
1:A:310:GLU:N	2:B:78:ASP:OD2	2.51	0.41
1:A:202:MET:HB2	1:A:214:THR:HB	2.02	0.41
1:A:334:GLU:HB3	3:A:772:HOH:O	2.20	0.41
1:A:666:LYS:O	1:A:668:GLN:N	2.53	0.41
1:A:658:VAL:O	1:A:662:MET:HG3	2.21	0.41
1:A:533:ILE:HB	1:A:534:PRO:HD3	2.02	0.40
1:A:433:LYS:HG3	3:A:911:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	508/538 (94%)	498 (98%)	8 (2%)	2 (0%)	30	28
2	B	61/81 (75%)	54 (88%)	5 (8%)	2 (3%)	3	1
All	All	569/619 (92%)	552 (97%)	13 (2%)	4 (1%)	18	15

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	665	ASP
2	B	61	ASP
1	A	667	PRO
2	B	64	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	422/447 (94%)	416 (99%)	6 (1%)	59	67
2	B	53/71 (75%)	52 (98%)	1 (2%)	50	58
All	All	475/518 (92%)	468 (98%)	7 (2%)	57	65

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	287	ASN
1	A	402	LEU
1	A	469	ARG
1	A	598	LEU
1	A	603	LEU
1	A	667	PRO
2	B	65	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	177	GLN
1	A	204	ASN
1	A	265	HIS
1	A	266	GLN
1	A	287	ASN
1	A	302	GLN
1	A	309	GLN
1	A	379	GLN
1	A	407	GLN
1	A	415	ASN
1	A	475	HIS
1	A	530	GLN
1	A	544	HIS
2	B	17	GLN
2	B	49	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	512/538 (95%)	0.38	60 (11%) 9 9	7, 28, 67, 101	2 (0%)
2	B	65/81 (80%)	1.34	16 (24%) 2 2	26, 39, 84, 91	0
All	All	577/619 (93%)	0.49	76 (13%) 7 7	7, 29, 73, 101	2 (0%)

All (76) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	60	ILE	9.1
1	A	169	ASN	8.3
1	A	339	THR	5.9
1	A	670	TYR	5.9
2	B	64	ALA	5.6
1	A	168	VAL	5.3
1	A	665	ASP	5.3
1	A	208	VAL	5.0
1	A	165	GLN	4.9
2	B	74	SER	4.8
1	A	560	PHE	4.8
2	B	54	GLN	4.7
1	A	206	ASN	4.7
1	A	207	ASP	4.6
1	A	166	VAL	4.5
1	A	561	VAL	4.4
1	A	174	MET	4.3
1	A	667	PRO	4.1
1	A	205	THR	3.9
2	B	62	GLN	3.9
2	B	77	GLU	3.8
1	A	151	ARG	3.8
2	B	52	LEU	3.7
1	A	559	GLN	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	167	VAL	3.5
1	A	248	VAL	3.3
2	B	63	GLY	3.3
1	A	432	TYR	3.3
2	B	61	ASP	3.3
2	B	78	ASP	3.2
1	A	173	VAL	3.1
1	A	668	GLN	3.1
1	A	663	SER	3.0
2	B	75	GLU	3.0
1	A	669	ASP	3.0
1	A	437	MET	3.0
1	A	664	GLU	2.9
2	B	50	SER	2.8
1	A	176	HIS	2.8
2	B	73	ARG	2.8
1	A	253	PHE	2.8
1	A	546	ASP	2.8
1	A	211	ALA	2.7
1	A	210	THR	2.6
1	A	178	LEU	2.6
1	A	182	GLU	2.5
1	A	175	VAL	2.5
1	A	245	GLY	2.5
1	A	453	ARG	2.5
2	B	76	THR	2.5
1	A	247	PRO	2.5
1	A	180	LYS	2.5
2	B	69	MET	2.4
1	A	209	GLU	2.4
2	B	65	GLU	2.4
1	A	418	THR	2.4
1	A	203	GLN	2.4
1	A	565	ARG	2.4
1	A	158	LYS	2.3
1	A	171	ALA	2.3
1	A	159	LEU	2.3
1	A	160	LEU	2.3
1	A	204	ASN	2.3
1	A	156	LEU	2.3
1	A	163	GLU	2.2
1	A	549	ARG	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	449	ARG	2.2
1	A	666	LYS	2.1
1	A	480	MET	2.1
1	A	164	ASP	2.1
1	A	474	ARG	2.1
1	A	170	LYS	2.1
1	A	246	SER	2.1
1	A	212	ARG	2.1
1	A	223	HIS	2.0
1	A	493	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.