



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 18, 2026 – 05:16 PM UTC

PDB ID : 4M1H / pdb_00004m1h
Title : X-ray crystal structure of Chlamydia trachomatis apo NrdB
Authors : Boal, A.K.; Rosenzweig, A.C.
Deposited on : 2013-08-02
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	366	
1	B	366	
1	C	366	
1	D	366	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 11834 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribonucleoside-diphosphate reductase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	331	Total	C	N	O	S	0	0	0
			2714	1743	463	496	12			
1	B	336	Total	C	N	O	S	0	0	0
			2759	1771	471	505	12			
1	C	331	Total	C	N	O	S	0	0	0
			2726	1749	464	501	12			
1	D	334	Total	C	N	O	S	0	0	0
			2747	1763	469	503	12			

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP O84835
A	-18	GLY	-	expression tag	UNP O84835
A	-17	SER	-	expression tag	UNP O84835
A	-16	SER	-	expression tag	UNP O84835
A	-15	HIS	-	expression tag	UNP O84835
A	-14	HIS	-	expression tag	UNP O84835
A	-13	HIS	-	expression tag	UNP O84835
A	-12	HIS	-	expression tag	UNP O84835
A	-11	HIS	-	expression tag	UNP O84835
A	-10	HIS	-	expression tag	UNP O84835
A	-9	SER	-	expression tag	UNP O84835
A	-8	SER	-	expression tag	UNP O84835
A	-7	GLY	-	expression tag	UNP O84835
A	-6	LEU	-	expression tag	UNP O84835
A	-5	VAL	-	expression tag	UNP O84835
A	-4	PRO	-	expression tag	UNP O84835
A	-3	ARG	-	expression tag	UNP O84835
A	-2	GLY	-	expression tag	UNP O84835
A	-1	SER	-	expression tag	UNP O84835
A	0	HIS	-	expression tag	UNP O84835
B	-19	MET	-	expression tag	UNP O84835

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-18	GLY	-	expression tag	UNP O84835
B	-17	SER	-	expression tag	UNP O84835
B	-16	SER	-	expression tag	UNP O84835
B	-15	HIS	-	expression tag	UNP O84835
B	-14	HIS	-	expression tag	UNP O84835
B	-13	HIS	-	expression tag	UNP O84835
B	-12	HIS	-	expression tag	UNP O84835
B	-11	HIS	-	expression tag	UNP O84835
B	-10	HIS	-	expression tag	UNP O84835
B	-9	SER	-	expression tag	UNP O84835
B	-8	SER	-	expression tag	UNP O84835
B	-7	GLY	-	expression tag	UNP O84835
B	-6	LEU	-	expression tag	UNP O84835
B	-5	VAL	-	expression tag	UNP O84835
B	-4	PRO	-	expression tag	UNP O84835
B	-3	ARG	-	expression tag	UNP O84835
B	-2	GLY	-	expression tag	UNP O84835
B	-1	SER	-	expression tag	UNP O84835
B	0	HIS	-	expression tag	UNP O84835
C	-19	MET	-	expression tag	UNP O84835
C	-18	GLY	-	expression tag	UNP O84835
C	-17	SER	-	expression tag	UNP O84835
C	-16	SER	-	expression tag	UNP O84835
C	-15	HIS	-	expression tag	UNP O84835
C	-14	HIS	-	expression tag	UNP O84835
C	-13	HIS	-	expression tag	UNP O84835
C	-12	HIS	-	expression tag	UNP O84835
C	-11	HIS	-	expression tag	UNP O84835
C	-10	HIS	-	expression tag	UNP O84835
C	-9	SER	-	expression tag	UNP O84835
C	-8	SER	-	expression tag	UNP O84835
C	-7	GLY	-	expression tag	UNP O84835
C	-6	LEU	-	expression tag	UNP O84835
C	-5	VAL	-	expression tag	UNP O84835
C	-4	PRO	-	expression tag	UNP O84835
C	-3	ARG	-	expression tag	UNP O84835
C	-2	GLY	-	expression tag	UNP O84835
C	-1	SER	-	expression tag	UNP O84835
C	0	HIS	-	expression tag	UNP O84835
D	-19	MET	-	expression tag	UNP O84835
D	-18	GLY	-	expression tag	UNP O84835
D	-17	SER	-	expression tag	UNP O84835

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-16	SER	-	expression tag	UNP O84835
D	-15	HIS	-	expression tag	UNP O84835
D	-14	HIS	-	expression tag	UNP O84835
D	-13	HIS	-	expression tag	UNP O84835
D	-12	HIS	-	expression tag	UNP O84835
D	-11	HIS	-	expression tag	UNP O84835
D	-10	HIS	-	expression tag	UNP O84835
D	-9	SER	-	expression tag	UNP O84835
D	-8	SER	-	expression tag	UNP O84835
D	-7	GLY	-	expression tag	UNP O84835
D	-6	LEU	-	expression tag	UNP O84835
D	-5	VAL	-	expression tag	UNP O84835
D	-4	PRO	-	expression tag	UNP O84835
D	-3	ARG	-	expression tag	UNP O84835
D	-2	GLY	-	expression tag	UNP O84835
D	-1	SER	-	expression tag	UNP O84835
D	0	HIS	-	expression tag	UNP O84835


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	246	Total O 246 246	0	0
2	B	265	Total O 265 265	0	0
2	C	197	Total O 197 197	0	0
2	D	180	Total O 180 180	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ribonucleoside-diphosphate reductase subunit beta




- Molecule 1: Ribonucleoside-diphosphate reductase subunit beta




- Molecule 1: Ribonucleoside-diphosphate reductase subunit beta



- Molecule 1: Ribonucleoside-diphosphate reductase subunit beta





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	74.97Å 97.39Å 99.24Å 90.00° 97.76° 90.00°	Depositor
Resolution (Å)	29.75 – 1.70 29.75 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.3 (29.75-1.70) 99.3 (29.75-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.89 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.179 , 0.210 0.184 , 0.216	Depositor DCC
R_{free} test set	7791 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	25.0	Xtriage
Anisotropy	0.068	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 36.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11834	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/2775	0.73	1/3754 (0.0%)
1	B	0.52	0/2820	0.73	0/3813
1	C	0.49	0/2786	0.74	0/3767
1	D	0.48	0/2808	0.73	0/3797
All	All	0.50	0/11189	0.74	1/15131 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	198	TYR	N-CA-C	5.71	118.71	111.69

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2714	0	2697	13	0
1	B	2759	0	2745	18	0
1	C	2726	0	2705	8	0
1	D	2747	0	2731	9	0
2	A	246	0	0	0	0
2	B	265	0	0	1	0
2	C	197	0	0	0	0
2	D	180	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	11834	0	10878	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:89:GLU:HG2	1:A:123:HIS:HB2	1.62	0.81
1:B:24:ASN:HD22	1:B:24:ASN:H	1.36	0.74
1:B:318:MET:O	1:B:322:ILE:HG12	1.89	0.72
1:D:215:GLY:O	1:D:219:GLN:HG3	1.93	0.68
1:C:97:VAL:HG13	1:D:97:VAL:HG13	1.75	0.68
1:A:97:VAL:HG13	1:B:97:VAL:HG13	1.79	0.64
1:A:89:GLU:HG3	1:A:120:GLU:OE1	1.98	0.62
1:B:52:LEU:HG	2:B:612:HOH:O	2.01	0.60
1:C:199:SER:HB2	1:C:268:GLU:OE2	2.04	0.57
1:B:85:PHE:HB3	1:B:127:PHE:CZ	2.40	0.56
1:C:79:LEU:HD11	1:D:-1:SER:C	2.34	0.52
1:A:72:GLU:HG2	1:B:-2:GLY:HA3	1.92	0.51
1:B:202:VAL:HG22	1:B:318:MET:HG3	1.91	0.51
1:A:24:ASN:HD22	1:A:24:ASN:C	2.18	0.51
1:B:295:ILE:O	1:B:299:ARG:HG2	2.12	0.50
1:A:85:PHE:HB3	1:A:127:PHE:CZ	2.48	0.48
1:B:24:ASN:HD22	1:B:24:ASN:N	2.06	0.47
1:C:85:PHE:HB3	1:C:127:PHE:CZ	2.50	0.46
1:A:295:ILE:O	1:A:299:ARG:HG2	2.16	0.46
1:B:7:ASP:O	1:B:10:GLN:HG2	2.16	0.45
1:C:262:LYS:HE2	1:C:308:ILE:HD12	1.97	0.45
1:D:24:ASN:HD22	1:D:24:ASN:C	2.25	0.45
1:D:280:ILE:HD12	1:D:281:LEU:HD22	1.99	0.45
1:A:179:GLN:HG2	1:A:253:LEU:HD22	1.99	0.44
1:B:280:ILE:HD11	1:B:283:LEU:HD12	1.98	0.44
1:C:201:PHE:HB3	1:C:318:MET:HE1	2.00	0.43
1:A:51:TRP:CE2	1:A:222:TYR:HB3	2.53	0.43
1:C:51:TRP:CE2	1:C:222:TYR:HB3	2.53	0.43
1:D:85:PHE:HB3	1:D:127:PHE:CZ	2.53	0.43
1:B:198:TYR:HA	1:B:201:PHE:CD2	2.53	0.43
1:B:51:TRP:CE2	1:B:222:TYR:HB3	2.54	0.43
1:A:149:ALA:HA	1:B:5:ILE:HD11	2.02	0.42
1:B:245:ASN:O	1:B:248:ILE:HG12	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:56:ILE:HD13	1:D:219:GLN:HG2	2.01	0.42
1:A:126:THR:HG23	1:A:219:GLN:OE1	2.20	0.42
1:B:123:HIS:O	1:B:126:THR:HB	2.19	0.42
1:C:280:ILE:HD11	1:C:283:LEU:HD12	2.02	0.42
1:A:75:ARG:NH2	1:B:-4:PRO:HG2	2.36	0.41
1:D:89:GLU:HG3	1:D:127:PHE:HE2	1.86	0.41
1:A:101:PHE:CG	1:B:97:VAL:HG11	2.56	0.40
1:D:179:GLN:HG2	1:D:253:LEU:HD22	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	329/366 (90%)	327 (99%)	2 (1%)	0	100	100
1	B	334/366 (91%)	332 (99%)	2 (1%)	0	100	100
1	C	329/366 (90%)	324 (98%)	5 (2%)	0	100	100
1	D	332/366 (91%)	330 (99%)	2 (1%)	0	100	100
All	All	1324/1464 (90%)	1313 (99%)	11 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	295/327 (90%)	290 (98%)	5 (2%)	53	38
1	B	300/327 (92%)	297 (99%)	3 (1%)	68	58
1	C	297/327 (91%)	290 (98%)	7 (2%)	43	26
1	D	299/327 (91%)	292 (98%)	7 (2%)	44	27
All	All	1191/1308 (91%)	1169 (98%)	22 (2%)	51	36

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	ASN
1	A	89	GLU
1	A	145	TYR
1	A	164	LYS
1	A	211	ASN
1	B	24	ASN
1	B	145	TYR
1	B	211	ASN
1	C	24	ASN
1	C	145	TYR
1	C	164	LYS
1	C	211	ASN
1	C	281	LEU
1	C	321	THR
1	C	324	LEU
1	D	7	ASP
1	D	11	LYS
1	D	24	ASN
1	D	145	TYR
1	D	211	ASN
1	D	276	LEU
1	D	281	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (41) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	24	ASN
1	A	29	ASN
1	A	30	GLN
1	A	45	ASN
1	A	49	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	94	ASN
1	A	116	GLN
1	A	211	ASN
1	A	232	ASN
1	A	273	GLN
1	A	294	HIS
1	B	2	GLN
1	B	24	ASN
1	B	30	GLN
1	B	49	ASN
1	B	94	ASN
1	B	116	GLN
1	B	158	GLN
1	B	221	GLN
1	B	273	GLN
1	C	14	ASN
1	C	29	ASN
1	C	49	ASN
1	C	103	HIS
1	C	143	ASN
1	C	146	ASN
1	C	158	GLN
1	C	169	ASN
1	C	221	GLN
1	D	14	ASN
1	D	24	ASN
1	D	30	GLN
1	D	49	ASN
1	D	94	ASN
1	D	158	GLN
1	D	169	ASN
1	D	179	GLN
1	D	210	GLN
1	D	273	GLN
1	D	294	HIS
1	D	325	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	331/366 (90%)	0.03	9 (2%) 56 60	14, 23, 36, 52	0
1	B	336/366 (91%)	0.01	14 (4%) 40 45	15, 23, 42, 56	0
1	C	331/366 (90%)	0.23	11 (3%) 49 53	18, 27, 43, 60	0
1	D	334/366 (91%)	0.30	13 (3%) 43 47	18, 30, 41, 50	0
All	All	1332/1464 (90%)	0.14	47 (3%) 47 52	14, 26, 41, 60	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	321	THR	4.4
1	B	-7	GLY	4.3
1	A	161	ILE	4.0
1	C	6	LEU	4.0
1	C	3	ALA	3.7
1	D	6	LEU	3.7
1	B	319	SER	3.6
1	B	324	LEU	3.6
1	D	7	ASP	3.5
1	C	7	ASP	3.4
1	B	322	ILE	3.3
1	A	6	LEU	3.2
1	C	164	LYS	3.1
1	A	252	GLU	3.1
1	C	5	ILE	3.0
1	A	321	THR	2.9
1	A	322	ILE	2.9
1	D	68	ASP	2.9
1	A	159	MET	2.8
1	D	322	ILE	2.8
1	B	25	GLN	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	-5	VAL	2.6
1	B	317	TRP	2.5
1	C	167	ASP	2.5
1	B	320	GLU	2.5
1	D	67	SER	2.5
1	B	-3	ARG	2.4
1	D	2	GLN	2.4
1	D	63	GLU	2.4
1	B	323	ASP	2.3
1	D	169	ASN	2.3
1	D	283	LEU	2.3
1	B	6	LEU	2.3
1	C	319	SER	2.3
1	A	25	GLN	2.2
1	C	166	LEU	2.2
1	B	5	ILE	2.2
1	D	-5	VAL	2.2
1	C	163	GLY	2.2
1	D	5	ILE	2.2
1	B	10	GLN	2.2
1	A	-3	ARG	2.2
1	D	1	MET	2.1
1	C	255	GLN	2.1
1	D	-2	GLY	2.1
1	A	7	ASP	2.0
1	C	-1	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.