



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 01:33 AM UTC

PDB ID : 1M2T / pdb_00001m2t
Title : Mistletoe Lectin I from Viscum album in Complex with Adenine Monophosphate. Crystal Structure at 1.9 Å Resolution
Authors : Krauspenhaar, R.; Rypniewski, W.; Kalkura, N.; Moore, K.; DeLucas, L.; Stoeva, S.; Mikhailov, A.; Voelter, W.; Betzel, C.
Deposited on : 2002-06-25
Resolution : 1.89 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	254	7% 71% 20% 6% ..
2	B	263	5% 75% 17% 6% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	A	502	-	-	-	X
4	FUC	A	501	-	-	-	X
4	FUC	B	601	-	-	-	X
6	GOL	A	600	-	X	-	-
6	GOL	A	601	-	X	-	-
6	GOL	A	702	-	-	X	-
6	GOL	B	701	-	-	X	-
6	GOL	B	705	-	X	-	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 4572 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called mistletoe lectin I A chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	248	Total	C	N	O	S	0	0	0
			1912	1211	323	374	4			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	8	THR	VAL	conflict	UNP P81446
A	9	ASP	THR	conflict	UNP P81446
A	10	GLN	HIS	conflict	UNP P81446
A	15	ALA	GLU	conflict	UNP P81446
A	19	SER	ARG	conflict	UNP P81446
A	23	VAL	LEU	conflict	UNP P81446
A	36	ASN	GLU	conflict	UNP P81446
A	45	VAL	ILE	conflict	UNP P81446
A	49	GLU	ASP	conflict	UNP P81446
A	50	GLY	ALA	conflict	UNP P81446
A	61	ALA	GLN	conflict	UNP P81446
A	63	GLY	GLN	conflict	UNP P81446
A	65	THR	SER	conflict	UNP P81446
A	66	ILE	VAL	conflict	UNP P81446
A	75	LEU	ALA	conflict	UNP P81446
A	81	GLU	GLN	conflict	UNP P81446
A	84	ASN	ASP	conflict	UNP P81446
A	90	SER	ARG	conflict	UNP P81446
A	94	ALA	ARG	conflict	UNP P81446
A	99	GLN	HIS	conflict	UNP P81446
A	100	ASP	LEU	conflict	UNP P81446
A	102	SER	THR	conflict	UNP P81446
A	106	SER	ARG	conflict	UNP P81446
A	109	GLN	LEU	conflict	UNP P81446
A	148	GLN	SER	conflict	UNP P81446
A	150	LYS	ARG	conflict	UNP P81446
A	176	ALA	TYR	conflict	UNP P81446

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	219	ALA	ARG	conflict	UNP P81446
A	223	ALA	PRO	conflict	UNP P81446
A	226	VAL	ASN	conflict	UNP P81446
A	227	ILE	PHE	conflict	UNP P81446
A	233	ILE	VAL	conflict	UNP P81446

- Molecule 2 is a protein called mistletoe lectin I B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	262	Total	C	N	O	S	0	0	0
			1995	1239	352	392	12			

There are 26 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	249	ALA	ASP	conflict	UNP P81830
B	253	THR	SER	conflict	UNP P81830
B	258	ILE	THR	conflict	UNP P81830
B	268	THR	ARG	conflict	UNP P81830
B	301	LYS	ARG	conflict	UNP P81830
B	342	GLY	ASP	conflict	UNP P81830
B	390	THR	VAL	conflict	UNP P81830
B	403	ALA	ASN	conflict	UNP P81830
B	408	TYR	TRP	conflict	UNP P81830
B	413	THR	ASP	conflict	UNP P81830
B	414	ALA	SER	conflict	UNP P81830
B	415	GLY	SER	conflict	UNP P81830
B	417	GLU	LYS	conflict	UNP P81830
B	?	-	GLY	deletion	UNP P81830
B	420	ARG	LYS	conflict	UNP P81830
B	434	LEU	ASN	conflict	UNP P81830
B	436	SER	ASP	conflict	UNP P81830
B	441	ASN	SER	conflict	UNP P81830
B	446	ILE	VAL	conflict	UNP P81830
B	457	ALA	-	insertion	UNP P81830
B	459	SER	ALA	conflict	UNP P81830
B	?	-	SER	deletion	UNP P81830
B	489	SER	LYS	conflict	UNP P81830
B	491	GLN	ARG	conflict	UNP P81830
B	501	ASN	LYS	conflict	UNP P81830
B	510	PRO	PHE	conflict	UNP P81830

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is alpha-L-fucopyranose (CCD ID: FUC) (formula: $C_6H_{12}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			10	6	4		
4	B	1	Total	C	O	0	0
			10	6	4		

- Molecule 5 is ADENINE (CCD ID: ADE) (formula: $C_5H_5N_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	N	0	0
			10	5	5		

- Molecule 6 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	203	Total	O	0	0
			203	203		
7	B	300	Total	O	0	0
			300	300		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: mistletoe lectin I A chain



- Molecule 2: mistletoe lectin I B chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	107.12Å 107.12Å 309.82Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	25.00 – 1.89 25.00 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.8 (25.00-1.89) 99.6 (25.00-1.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.43 (at 1.90Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.210 , 0.240 0.197 , 0.224	Depositor DCC
R_{free} test set	4159 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	38.2	Xtriage
Anisotropy	0.350	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 58.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4572	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NAG, ADE, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.73	0/1950	1.46	15/2659 (0.6%)
2	B	0.83	1/2034 (0.0%)	1.55	23/2774 (0.8%)
All	All	0.78	1/3984 (0.0%)	1.51	38/5433 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
2	B	0	8
All	All	0	10

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	510	PRO	N-CD	5.71	1.55	1.47

The worst 5 of 38 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	383	ASN	CA-CB-CG	-8.70	103.90	112.60
1	A	112	ASN	CA-CB-CG	8.67	121.27	112.60
1	A	215	ASN	OD1-CG-ND2	-7.70	114.90	122.60
1	A	222	ILE	CB-CA-C	7.56	122.78	110.69
2	B	383	ASN	OD1-CG-ND2	-7.39	115.21	122.60

There are no chirality outliers.

5 of 10 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	7	ARG	Mainchain
1	A	86	SER	Mainchain
2	B	255	SER	Mainchain
2	B	288	LYS	Mainchain
2	B	343	ASN	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1912	0	1877	50	0
2	B	1995	0	1937	60	0
3	A	28	0	26	5	0
3	B	56	0	52	12	0
4	A	10	0	10	3	0
4	B	10	0	10	0	0
5	A	10	0	4	1	0
6	A	24	0	23	6	0
6	B	24	0	25	7	0
7	A	203	0	0	3	0
7	B	300	0	0	10	1
All	All	4572	0	3964	115	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 115 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:112:ASN:HD21	3:A:500:NAG:C1	1.59	1.13
2:B:503:ASN:HD21	6:B:701:GOL:H31	1.16	1.08
2:B:416:GLN:HE21	2:B:418:ASN:HD21	1.04	1.00
2:B:288:LYS:H	2:B:295:GLN:HE22	1.11	0.97
2:B:343:ASN:HD21	3:B:602:NAG:C1	1.78	0.96

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:B:768:HOH:O	7:B:768:HOH:O[12_554]	1.76	0.44

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	246/254 (97%)	239 (97%)	6 (2%)	1 (0%)	30	22
2	B	260/263 (99%)	256 (98%)	3 (1%)	1 (0%)	30	22
All	All	506/517 (98%)	495 (98%)	9 (2%)	2 (0%)	30	22

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	254	ALA
1	A	223	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	208/215 (97%)	192 (92%)	16 (8%)	12	5
2	B	218/219 (100%)	208 (95%)	10 (5%)	24	16
All	All	426/434 (98%)	400 (94%)	26 (6%)	17	9

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	230	LEU
2	B	255	SER
2	B	477	ASN
2	B	253	THR
2	B	258	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	308	ASN
2	B	503	ASN
2	B	352	ASN
2	B	485	GLN
2	B	348	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	B	604	-	14,14,15	1.34	1 (7%)	17,19,21	2.04	5 (29%)
4	FUC	A	501	-	10,10,11	1.06	1 (10%)	14,14,16	1.51	4 (28%)
6	GOL	A	601	-	5,5,5	1.61	1 (20%)	5,5,5	2.19	3 (60%)
6	GOL	B	703	-	5,5,5	0.74	0	5,5,5	1.09	0
3	NAG	A	500	-	14,14,15	1.29	1 (7%)	17,19,21	2.09	4 (23%)
6	GOL	B	705	-	5,5,5	1.57	1 (20%)	5,5,5	2.15	2 (40%)
3	NAG	B	600	-	14,14,15	1.28	1 (7%)	17,19,21	1.77	4 (23%)
6	GOL	A	700	-	5,5,5	1.32	0	5,5,5	1.74	1 (20%)
6	GOL	B	701	-	5,5,5	1.52	0	5,5,5	1.59	1 (20%)
5	ADE	A	9550	-	11,11,11	1.07	0	15,15,15	1.53	2 (13%)
6	GOL	A	702	-	5,5,5	1.25	1 (20%)	5,5,5	1.61	1 (20%)
3	NAG	B	603	2	14,14,15	1.57	2 (14%)	17,19,21	2.62	4 (23%)
6	GOL	A	600	-	5,5,5	1.49	1 (20%)	5,5,5	1.90	2 (40%)
3	NAG	B	602	-	14,14,15	1.34	1 (7%)	17,19,21	1.93	3 (17%)
4	FUC	B	601	-	10,10,11	0.88	0	14,14,16	2.00	3 (21%)
6	GOL	B	704	-	5,5,5	1.39	0	5,5,5	1.40	0
3	NAG	A	502	-	14,14,15	1.39	3 (21%)	17,19,21	3.03	7 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. ' - ' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	604	-	-	2/6/23/26	0/1/1/1
6	GOL	A	601	-	-	3/4/4/4	-
6	GOL	B	703	-	-	0/4/4/4	-
4	FUC	A	501	-	-	-	0/1/1/1
3	NAG	A	500	-	-	3/6/23/26	0/1/1/1
6	GOL	B	705	-	-	3/4/4/4	-
3	NAG	B	600	-	-	2/6/23/26	0/1/1/1
6	GOL	A	700	-	-	2/4/4/4	-
6	GOL	B	701	-	-	3/4/4/4	-
5	ADE	A	9550	-	-	-	0/2/2/2
6	GOL	A	702	-	-	2/4/4/4	-
3	NAG	B	603	2	-	0/6/23/26	0/1/1/1
6	GOL	A	600	-	-	3/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	602	-	-	3/6/23/26	0/1/1/1
6	GOL	B	704	-	-	2/4/4/4	-
4	FUC	B	601	-	-	-	0/1/1/1
3	NAG	A	502	-	-	4/6/23/26	0/1/1/1

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	602	NAG	O7-C7	-4.06	1.14	1.23
3	B	604	NAG	O7-C7	-4.03	1.14	1.23
3	B	603	NAG	C1-C2	-3.95	1.47	1.52
3	B	600	NAG	O7-C7	-3.67	1.15	1.23
3	A	500	NAG	O7-C7	-3.55	1.15	1.23

The worst 5 of 46 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	603	NAG	O5-C1-C2	8.16	123.91	111.29
3	B	602	NAG	C1-O5-C5	-5.91	104.27	112.19
3	A	502	NAG	C1-O5-C5	5.87	120.05	112.19
3	A	502	NAG	C1-C2-N2	5.64	119.32	110.43
3	A	500	NAG	C3-C4-C5	5.55	120.29	110.23

There are no chirality outliers.

5 of 32 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	500	NAG	O7-C7-N2-C2
3	A	502	NAG	C8-C7-N2-C2
3	A	502	NAG	O7-C7-N2-C2
3	B	602	NAG	C1-C2-N2-C7
3	B	602	NAG	C8-C7-N2-C2

There are no ring outliers.

13 monomers are involved in 31 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	604	NAG	2	0
4	A	501	FUC	3	0
6	A	601	GOL	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	500	NAG	5	0
6	B	705	GOL	1	0
3	B	600	NAG	5	0
6	B	701	GOL	5	0
5	A	9550	ADE	1	0
6	A	702	GOL	4	0
3	B	603	NAG	2	0
6	A	600	GOL	1	0
3	B	602	NAG	5	0
6	B	704	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	248/254 (97%)	0.51	17 (6%) 23 24	34, 45, 65, 75	0
2	B	262/263 (99%)	0.17	12 (4%) 37 40	30, 39, 56, 75	0
All	All	510/517 (98%)	0.34	29 (5%) 29 31	30, 41, 63, 75	0

The worst 5 of 29 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	249	ALA	7.2
2	B	253	THR	7.0
1	A	248	GLY	6.7
2	B	250	VAL	5.8
1	A	223	ALA	5.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	FUC	B	601	10/11	0.55	0.62	76,77,78,79	10
3	NAG	A	500	14/15	0.56	0.20	91,92,93,93	0
3	NAG	A	502	14/15	0.58	0.48	74,75,76,77	14
3	NAG	B	600	14/15	0.63	0.19	63,76,81,84	0
6	GOL	A	601	6/6	0.69	0.27	88,89,91,92	0
6	GOL	A	600	6/6	0.71	0.24	83,88,90,93	0
4	FUC	A	501	10/11	0.72	0.40	78,79,80,81	10
3	NAG	B	604	14/15	0.72	0.18	62,74,79,79	0
3	NAG	B	602	14/15	0.78	0.17	61,64,67,71	0
6	GOL	B	704	6/6	0.81	0.17	55,58,63,63	0
6	GOL	B	701	6/6	0.85	0.20	55,57,63,68	0
6	GOL	A	702	6/6	0.88	0.13	49,50,53,54	0
5	ADE	A	9550	10/10	0.90	0.10	38,55,61,64	0
3	NAG	B	603	14/15	0.92	0.11	35,44,50,53	0
6	GOL	A	700	6/6	0.93	0.11	42,48,52,55	0
6	GOL	B	703	6/6	0.94	0.10	33,37,43,47	0
6	GOL	B	705	6/6	0.94	0.10	41,57,60,62	0

6.5 Other polymers [i](#)

There are no such residues in this entry.