



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:05 PM UTC

PDB ID : 4M2L / pdb_00004m2l
Title : Gamma subunit of the translation initiation factor 2 from *Sulfolobus solfataricus* in nucleotide-free form
Authors : Nikonov, O.S.; Stolboushkina, E.A.; Arkhipova, V.I.; Gabdulkhakov, A.G.; Nikulin, A.D.; Garber, M.B.; Nikonov, S.V.
Deposited on : 2013-08-05
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3689 (2.16-2.12)
Clashscore	190562	3812 (2.16-2.12)
Ramachandran outliers	187476	3773 (2.16-2.12)
Sidechain outliers	187428	3772 (2.16-2.12)
RSRZ outliers	180081	3691 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	410	13% 75% 18% ...

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3483 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Translation initiation factor 2 subunit gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	398	3080	1977	522	570	11	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	LEU	deletion	UNP Q980A5
A	?	-	LYS	deletion	UNP Q980A5
A	?	-	ARG	deletion	UNP Q980A5
A	?	-	GLY	deletion	UNP Q980A5
A	?	-	MET	deletion	UNP Q980A5

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	255	Total O 255 255	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Translation initiation factor 2 subunit gamma



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	95.81Å 95.81Å 165.24Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.91 – 2.15 47.91 – 2.15	Depositor EDS
% Data completeness (in resolution range)	98.2 (47.91-2.15) 99.9 (47.91-2.15)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.26 (at 2.14Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.188 , 0.228 0.195 , 0.235	Depositor DCC
R_{free} test set	2446 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	40.1	Xtriage
Anisotropy	0.595	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 51.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.029 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3483	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, EDO, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/3133	0.82	4/4242 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	17	HIS	N-CA-C	-9.77	97.07	110.35
1	A	263	LYS	N-CA-C	-6.04	102.40	110.55
1	A	48	LYS	CB-CA-C	-6.00	109.67	116.63
1	A	104	MET	N-CA-CB	5.32	117.94	110.12

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3080	0	3206	73	0
2	A	20	0	0	0	0
3	A	20	0	0	0	0
4	A	108	0	162	17	0
5	A	255	0	0	3	0
All	All	3483	0	3368	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:101:MET:HG2	1:A:358:MET:HE1	1.46	0.97
1:A:149:ASN:HD21	4:A:534:EDO:H21	1.30	0.94
1:A:207:ARG:HH12	4:A:513:EDO:H11	1.46	0.79
1:A:340:VAL:HG22	1:A:345:MET:HE1	1.65	0.78
1:A:150:LYS:HG2	4:A:534:EDO:H12	1.65	0.78
1:A:356:THR:H	4:A:519:EDO:H21	1.48	0.75
1:A:17:HIS:CD2	1:A:128:GLN:HB2	2.27	0.70
1:A:108:ALA:HB1	1:A:139:ILE:HD12	1.77	0.67
1:A:344:GLU:HA	1:A:344:GLU:OE1	1.99	0.62
1:A:138:ILE:HD11	1:A:336:LEU:HD11	1.82	0.61
1:A:94:ALA:HB3	1:A:95:PRO:HD3	1.84	0.60
1:A:22:LYS:CE	1:A:95:PRO:HG3	2.33	0.59
1:A:64:LYS:H	4:A:527:EDO:H12	1.68	0.59
1:A:23:THR:HG22	1:A:93:ASP:OD1	2.03	0.59
1:A:337:GLU:HG2	1:A:338:ARG:HG2	1.87	0.56
1:A:347:LYS:HZ2	1:A:347:LYS:HB3	1.71	0.56
1:A:229:GLN:HB2	1:A:232:GLU:HG3	1.88	0.56
1:A:263:LYS:CG	1:A:266:LYS:HE2	2.34	0.56
1:A:22:LYS:HE2	1:A:95:PRO:CD	2.36	0.56
1:A:110:LEU:HD22	1:A:241:SER:HB3	1.88	0.56
1:A:94:ALA:HB3	1:A:95:PRO:CD	2.36	0.55
1:A:22:LYS:NZ	1:A:95:PRO:HG3	2.22	0.55
1:A:263:LYS:HG2	1:A:266:LYS:HE2	1.87	0.55
1:A:355:GLU:HB2	4:A:519:EDO:H12	1.87	0.55
1:A:20:HIS:HA	5:A:847:HOH:O	2.06	0.54
1:A:22:LYS:HE2	1:A:95:PRO:HD3	1.88	0.54
1:A:263:LYS:HG3	1:A:264:GLN:N	2.22	0.54
1:A:149:ASN:ND2	4:A:534:EDO:H21	2.13	0.53
1:A:50:GLY:HA3	1:A:93:ASP:HB2	1.90	0.53
1:A:263:LYS:HG2	1:A:266:LYS:HD3	1.91	0.52
1:A:212:LYS:HE3	4:A:531:EDO:H12	1.92	0.52
1:A:338:ARG:HA	1:A:348:VAL:HG23	1.94	0.49
1:A:101:MET:HG3	1:A:105:LEU:CD1	2.42	0.49
1:A:207:ARG:NH1	4:A:513:EDO:H11	2.22	0.48
1:A:130:ARG:HD3	1:A:345:MET:HE1	1.94	0.48
1:A:287:LYS:HD3	1:A:287:LYS:H	1.79	0.48
1:A:105:LEU:HD13	1:A:360:SER:OG	2.15	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:22:LYS:HE2	1:A:95:PRO:HG3	1.97	0.47
1:A:287:LYS:H	1:A:287:LYS:CD	2.27	0.47
1:A:279:ILE:HG21	1:A:289:ALA:HB2	1.97	0.47
1:A:391:ASN:HD22	4:A:525:EDO:H22	1.80	0.47
1:A:349:ASP:O	1:A:399:ARG:NH2	2.35	0.46
1:A:151:VAL:HG11	1:A:182:PRO:CB	2.46	0.46
1:A:74:CYS:SG	1:A:83:PRO:HB3	2.56	0.46
1:A:175:ALA:O	1:A:178:VAL:HG22	2.15	0.46
1:A:252:GLU:OE1	1:A:275:LYS:HE3	2.16	0.45
1:A:313:GLY:O	4:A:510:EDO:H11	2.17	0.45
1:A:372:SER:HB3	1:A:379:GLU:HB2	1.97	0.45
1:A:94:ALA:N	1:A:95:PRO:HD2	2.32	0.45
1:A:139:ILE:HG13	1:A:141:VAL:HG23	1.99	0.45
1:A:337:GLU:H	1:A:337:GLU:CD	2.25	0.44
1:A:101:MET:HG2	1:A:358:MET:CE	2.33	0.44
1:A:359:LEU:HD12	1:A:382:LEU:HD21	1.99	0.44
1:A:48:LYS:HB3	1:A:49:LEU:H	1.49	0.43
1:A:358:MET:HG3	1:A:367:LEU:HD13	2.01	0.43
1:A:401:ILE:HB	1:A:406:ARG:HD2	2.00	0.43
1:A:255:VAL:HB	1:A:272:ILE:HB	2.01	0.42
1:A:315:ILE:HG23	4:A:510:EDO:H12	2.00	0.42
1:A:263:LYS:HB3	1:A:266:LYS:HE2	2.02	0.42
1:A:100:LEU:HD22	1:A:100:LEU:HA	1.91	0.42
1:A:280:ARG:HE	1:A:280:ARG:HB2	1.54	0.42
1:A:216:LEU:HD12	1:A:315:ILE:HG22	2.01	0.42
1:A:101:MET:HG3	1:A:105:LEU:HD12	2.02	0.42
1:A:324:PRO:HD2	1:A:388:VAL:O	2.19	0.42
1:A:14:VAL:HG12	1:A:22:LYS:HB2	2.02	0.41
1:A:263:LYS:CB	1:A:266:LYS:HE2	2.50	0.41
1:A:211:GLN:HG2	4:A:515:EDO:H11	2.03	0.41
1:A:17:HIS:HB3	4:A:530:EDO:H21	2.03	0.41
1:A:251:GLN:HG2	4:A:522:EDO:H11	2.01	0.41
1:A:22:LYS:NZ	5:A:848:HOH:O	2.54	0.41
4:A:510:EDO:H22	5:A:758:HOH:O	2.21	0.41
1:A:20:HIS:CE1	1:A:150:LYS:HE2	2.56	0.40
1:A:150:LYS:O	1:A:153:VAL:HG22	2.21	0.40
1:A:365:THR:HB	4:A:528:EDO:H12	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	392/410 (96%)	385 (98%)	6 (2%)	1 (0%)	36	33

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	21	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	342/353 (97%)	315 (92%)	27 (8%)	11	7

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	HIS
1	A	20	HIS
1	A	22	LYS
1	A	25	LEU
1	A	27	GLN
1	A	48	LYS
1	A	49	LEU
1	A	60	GLU
1	A	100	LEU
1	A	101	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	104	MET
1	A	131	GLU
1	A	212	LYS
1	A	216	LEU
1	A	217	VAL
1	A	218	ILE
1	A	232	GLU
1	A	246	LEU
1	A	263	LYS
1	A	264	GLN
1	A	287	LYS
1	A	304	SER
1	A	339	VAL
1	A	343	LYS
1	A	347	LYS
1	A	367	LEU
1	A	384	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	244	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

35 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	502	-	4,4,4	0.25	0	6,6,6	0.10	0
4	EDO	A	516	-	3,3,3	0.43	0	2,2,2	0.38	0
4	EDO	A	514	-	3,3,3	0.46	0	2,2,2	0.40	0
4	EDO	A	527	-	3,3,3	0.39	0	2,2,2	0.41	0
4	EDO	A	512	-	3,3,3	0.41	0	2,2,2	0.45	0
4	EDO	A	520	-	3,3,3	0.43	0	2,2,2	0.31	0
4	EDO	A	535	-	3,3,3	0.44	0	2,2,2	0.36	0
3	PO4	A	505	-	4,4,4	0.95	0	6,6,6	0.54	0
4	EDO	A	530	-	3,3,3	0.49	0	2,2,2	0.27	0
4	EDO	A	517	-	3,3,3	0.48	0	2,2,2	0.25	0
4	EDO	A	518	-	3,3,3	0.48	0	2,2,2	0.26	0
3	PO4	A	507	-	4,4,4	1.56	0	6,6,6	0.75	0
2	SO4	A	501	-	4,4,4	0.22	0	6,6,6	0.21	0
4	EDO	A	526	-	3,3,3	0.44	0	2,2,2	0.41	0
4	EDO	A	513	-	3,3,3	0.47	0	2,2,2	0.29	0
4	EDO	A	515	-	3,3,3	0.47	0	2,2,2	0.38	0
4	EDO	A	534	-	3,3,3	0.47	0	2,2,2	0.31	0
4	EDO	A	525	-	3,3,3	0.43	0	2,2,2	0.40	0
4	EDO	A	509	-	3,3,3	0.49	0	2,2,2	0.26	0
4	EDO	A	522	-	3,3,3	0.45	0	2,2,2	0.30	0
4	EDO	A	523	-	3,3,3	0.43	0	2,2,2	0.37	0
3	PO4	A	508	-	4,4,4	0.85	0	6,6,6	0.65	0
2	SO4	A	504	-	4,4,4	0.27	0	6,6,6	0.38	0
4	EDO	A	532	-	3,3,3	0.45	0	2,2,2	0.34	0
4	EDO	A	533	-	3,3,3	0.47	0	2,2,2	0.31	0
4	EDO	A	528	-	3,3,3	0.50	0	2,2,2	0.23	0
3	PO4	A	506	-	4,4,4	0.92	0	6,6,6	0.45	0
2	SO4	A	503	-	4,4,4	0.42	0	6,6,6	0.07	0
4	EDO	A	529	-	3,3,3	0.41	0	2,2,2	0.49	0
4	EDO	A	521	-	3,3,3	0.44	0	2,2,2	0.42	0
4	EDO	A	524	-	3,3,3	0.44	0	2,2,2	0.36	0
4	EDO	A	531	-	3,3,3	0.39	0	2,2,2	0.47	0
4	EDO	A	510	-	3,3,3	0.40	0	2,2,2	0.46	0
4	EDO	A	511	-	3,3,3	0.43	0	2,2,2	0.51	0
4	EDO	A	519	-	3,3,3	0.43	0	2,2,2	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	516	-	-	1/1/1/1	-
4	EDO	A	514	-	-	0/1/1/1	-
4	EDO	A	527	-	-	1/1/1/1	-
4	EDO	A	512	-	-	0/1/1/1	-
4	EDO	A	520	-	-	0/1/1/1	-
4	EDO	A	535	-	-	0/1/1/1	-
4	EDO	A	530	-	-	0/1/1/1	-
4	EDO	A	517	-	-	0/1/1/1	-
4	EDO	A	518	-	-	1/1/1/1	-
4	EDO	A	526	-	-	0/1/1/1	-
4	EDO	A	513	-	-	0/1/1/1	-
4	EDO	A	515	-	-	0/1/1/1	-
4	EDO	A	534	-	-	0/1/1/1	-
4	EDO	A	525	-	-	0/1/1/1	-
4	EDO	A	509	-	-	0/1/1/1	-
4	EDO	A	522	-	-	1/1/1/1	-
4	EDO	A	523	-	-	0/1/1/1	-
4	EDO	A	532	-	-	1/1/1/1	-
4	EDO	A	533	-	-	0/1/1/1	-
4	EDO	A	528	-	-	0/1/1/1	-
4	EDO	A	529	-	-	0/1/1/1	-
4	EDO	A	521	-	-	1/1/1/1	-
4	EDO	A	524	-	-	0/1/1/1	-
4	EDO	A	531	-	-	0/1/1/1	-
4	EDO	A	510	-	-	0/1/1/1	-
4	EDO	A	511	-	-	0/1/1/1	-
4	EDO	A	519	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	516	EDO	O1-C1-C2-O2
4	A	527	EDO	O1-C1-C2-O2
4	A	532	EDO	O1-C1-C2-O2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	A	521	EDO	O1-C1-C2-O2
4	A	522	EDO	O1-C1-C2-O2
4	A	518	EDO	O1-C1-C2-O2

There are no ring outliers.

11 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	527	EDO	1	0
4	A	530	EDO	1	0
4	A	513	EDO	2	0
4	A	515	EDO	1	0
4	A	534	EDO	3	0
4	A	525	EDO	1	0
4	A	522	EDO	1	0
4	A	528	EDO	1	0
4	A	531	EDO	1	0
4	A	510	EDO	3	0
4	A	519	EDO	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	398/410 (97%)	0.45	52 (13%) 7 9	16, 34, 75, 115	96 (24%)

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	49	LEU	17.2
1	A	48	LYS	9.4
1	A	18	VAL	7.9
1	A	99	VAL	6.5
1	A	94	ALA	6.3
1	A	342	ALA	6.1
1	A	101	MET	6.0
1	A	100	LEU	5.8
1	A	22	LYS	5.8
1	A	340	VAL	5.7
1	A	265	GLY	5.5
1	A	95	PRO	5.4
1	A	20	HIS	4.9
1	A	104	MET	4.8
1	A	23	THR	4.8
1	A	102	ALA	4.8
1	A	341	GLY	4.4
1	A	264	GLN	4.3
1	A	46	THR	4.1
1	A	17	HIS	4.1
1	A	263	LYS	3.9
1	A	16	GLY	3.8
1	A	347	LYS	3.8
1	A	21	GLY	3.6
1	A	105	LEU	3.5
1	A	19	ASP	3.4
1	A	109	ALA	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	32	ILE	3.2
1	A	103	THR	3.2
1	A	403	GLY	3.2
1	A	266	LYS	3.1
1	A	343	LYS	3.1
1	A	262	GLU	3.1
1	A	142	LYS	3.0
1	A	106	SER	3.0
1	A	209	LEU	2.9
1	A	267	VAL	2.9
1	A	110	LEU	2.9
1	A	47	ILE	2.8
1	A	345	MET	2.8
1	A	107	GLY	2.8
1	A	283	ASP	2.8
1	A	108	ALA	2.7
1	A	344	GLU	2.6
1	A	404	ARG	2.6
1	A	346	LEU	2.4
1	A	50	GLY	2.4
1	A	339	VAL	2.2
1	A	338	ARG	2.2
1	A	261	VAL	2.1
1	A	168	GLN	2.0
1	A	384	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	EDO	A	535	4/4	0.65	0.25	81,82,85,89	0
4	EDO	A	533	4/4	0.69	0.35	63,68,72,75	2
4	EDO	A	524	4/4	0.72	0.21	68,71,78,81	0
4	EDO	A	529	4/4	0.72	0.32	47,49,54,54	3
4	EDO	A	520	4/4	0.73	0.29	47,57,60,63	2
4	EDO	A	530	4/4	0.75	0.31	48,56,65,68	3
4	EDO	A	518	4/4	0.75	0.25	49,50,56,59	1
4	EDO	A	515	4/4	0.75	0.20	41,44,48,58	1
4	EDO	A	525	4/4	0.78	0.23	69,71,75,79	3
4	EDO	A	514	4/4	0.79	0.23	46,48,51,62	3
4	EDO	A	511	4/4	0.79	0.23	55,67,71,79	1
4	EDO	A	516	4/4	0.80	0.14	66,75,79,81	0
4	EDO	A	522	4/4	0.81	0.17	44,52,54,66	1
4	EDO	A	531	4/4	0.82	0.18	42,43,46,56	2
4	EDO	A	532	4/4	0.82	0.24	33,33,35,55	4
4	EDO	A	517	4/4	0.83	0.21	40,42,54,65	3
4	EDO	A	528	4/4	0.83	0.22	43,48,50,52	2
3	PO4	A	506	5/5	0.85	0.12	61,64,68,77	2
4	EDO	A	526	4/4	0.85	0.25	57,57,65,67	3
4	EDO	A	512	4/4	0.85	0.20	50,65,65,67	1
4	EDO	A	527	4/4	0.87	0.17	43,44,55,70	1
3	PO4	A	508	5/5	0.87	0.11	35,53,58,63	4
4	EDO	A	523	4/4	0.88	0.16	46,56,59,60	2
4	EDO	A	534	4/4	0.88	0.19	33,45,47,50	2
4	EDO	A	519	4/4	0.88	0.21	36,50,51,66	2
4	EDO	A	513	4/4	0.89	0.18	36,41,55,56	2
2	SO4	A	503	5/5	0.91	0.14	51,53,56,74	4
4	EDO	A	521	4/4	0.92	0.12	44,45,48,49	1
4	EDO	A	510	4/4	0.93	0.14	26,35,45,47	3
2	SO4	A	504	5/5	0.94	0.09	56,58,62,72	4
3	PO4	A	505	5/5	0.94	0.10	58,62,65,73	4
3	PO4	A	507	5/5	0.96	0.09	38,50,53,62	4
4	EDO	A	509	4/4	0.96	0.08	29,30,36,41	3
2	SO4	A	502	5/5	0.98	0.07	41,50,52,54	5
2	SO4	A	501	5/5	0.98	0.05	37,39,40,40	3

6.5 Other polymers ⓘ

There are no such residues in this entry.