



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 01:29 PM UTC

PDB ID : 5M2T / pdb_00005m2t
Title : X-ray structure of uridine phosphorylase from *Vibrio cholerae* in complex with uridine at 1.03 Å resolution
Authors : Prokofev, I.I.; Lashkov, A.A.; Gabdulkhakov, A.G.; Betzel, C.; Mikhailov, A.M.
Deposited on : 2016-10-13
Resolution : 1.03 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1036 (1.06-1.02)
Clashscore	190562	1063 (1.06-1.02)
Ramachandran outliers	187476	1034 (1.06-1.02)
Sidechain outliers	187428	1035 (1.06-1.02)
RSRZ outliers	180081	1035 (1.06-1.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	253	2% 89% 11%
1	B	253	2% 87% 13%
1	C	253	8% 88% 11%
1	D	253	3% 88% 11%
1	E	253	6% 87% 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	253	% 89% 10%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	CL	D	303	-	-	X	-
6	EDO	E	303	-	-	X	-

2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 28257 atoms, of which 13402 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uridine phosphorylase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	251	Total	C	H	N	O	S	0	49	0
			4312	1328	2186	376	405	17			
1	B	252	Total	C	H	N	O	S	0	51	0
			4322	1336	2190	377	401	18			
1	C	251	Total	C	H	N	O	S	0	62	0
			4415	1359	2228	389	420	19			
1	D	252	Total	C	H	N	O	S	0	55	0
			4377	1348	2214	384	413	18			
1	E	252	Total	C	H	N	O	S	0	59	0
			4491	1381	2279	398	415	18			
1	F	251	Total	C	H	N	O	S	0	50	0
			4384	1346	2227	388	406	17			

- Molecule 2 is URIDINE (CCD ID: URI) (formula: $C_9H_{12}N_2O_6$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	0	0
			29	9	12	2	6		
2	B	1	Total	C	H	N	O	0	0
			29	9	12	2	6		
2	C	1	Total	C	H	N	O	0	0
			29	9	12	2	6		
2	D	1	Total	C	H	N	O	0	0
			29	9	12	2	6		
2	E	1	Total	C	H	N	O	0	0
			29	9	12	2	6		
2	F	1	Total	C	H	N	O	0	0
			29	9	12	2	6		

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		
3	C	1	Total	Na	0	0
			1	1		
3	E	1	Total	Na	0	0
			1	1		


- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Cl	0	0
			1	1		
4	C	2	Total	Cl	0	0
			2	2		
4	D	2	Total	Cl	0	0
			2	2		
4	F	1	Total	Cl	0	0
			1	1		

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	1	Total	Mg	0	0
			1	1		

- Molecule 6 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	E	1	Total	C	H	O	0	0
			10	2	6	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	327	Total	O	0	3
			330	330		
7	B	301	Total	O	0	0
			301	301		
7	C	235	Total	O	0	3
			238	238		
7	D	290	Total	O	0	3
			293	293		
7	E	268	Total	O	0	2
			270	270		
7	F	328	Total	O	0	2
			330	330		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Uridine phosphorylase




- Molecule 1: Uridine phosphorylase



- Molecule 1: Uridine phosphorylase





- Molecule 1: Uridine phosphorylase







● Molecule 1: Uridine phosphorylase



● Molecule 1: Uridine phosphorylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	64.32Å 72.04Å 89.19Å 110.56° 107.53° 85.83°	Depositor
Resolution (Å)	44.46 – 1.03 44.46 – 1.03	Depositor EDS
% Data completeness (in resolution range)	93.2 (44.46-1.03) 93.2 (44.46-1.03)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.53 (at 1.03Å)	Xtriage
Refinement program	PHENIX 1.10_2155	Depositor
R, R_{free}	0.118 , 0.137 0.121 , 0.141	Depositor DCC
R_{free} test set	6560 reflections (0.93%)	wwPDB-VP
Wilson B-factor (Å ²)	8.5	Xtriage
Anisotropy	0.158	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 52.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	28257	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: URI, MG, EDO, NA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.62	0/2309	0.75	0/3121
1	B	0.62	0/2343	0.80	2/3172 (0.1%)
1	C	0.56	0/2436	0.74	2/3297 (0.1%)
1	D	0.61	0/2374	0.74	0/3208
1	E	0.57	0/2432	0.70	0/3283
1	F	0.60	1/2371 (0.0%)	0.74	0/3204
All	All	0.60	1/14265 (0.0%)	0.75	4/19285 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	90	ARG	CG-CD	-5.78	1.35	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	226	GLU	CA-C-N	-7.80	114.91	123.02
1	C	226	GLU	C-N-CA	-7.80	114.91	123.02
1	B	90[A]	ARG	NE-CZ-NH1	-7.58	113.92	121.50
1	B	90[B]	ARG	NE-CZ-NH1	-7.58	113.92	121.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2126	2186	2025	27	0
1	B	2132	2190	2020	26	0
1	C	2187	2228	2003	21	0
1	D	2163	2214	2024	26	0
1	E	2212	2279	2101	28	0
1	F	2157	2227	2063	25	0
2	A	17	12	12	0	0
2	B	17	12	12	1	0
2	C	17	12	12	0	0
2	D	17	12	12	0	0
2	E	17	12	12	2	0
2	F	17	12	12	0	0
3	A	1	0	0	0	0
3	C	1	0	0	0	0
3	E	1	0	0	0	0
4	B	1	0	0	1	0
4	C	2	0	0	1	0
4	D	2	0	0	2	0
4	F	1	0	0	0	0
5	D	1	0	0	0	0
6	E	4	6	6	6	0
7	A	330	0	0	10	0
7	B	301	0	0	9	0
7	C	238	0	0	9	0
7	D	293	0	0	14	0
7	E	270	0	0	11	0
7	F	330	0	0	15	0
All	All	14855	13402	12314	154	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 154 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:177[A]:ARG:NH2	7:C:402:HOH:O	1.77	1.16
1:F:122:PHE:O	7:F:401:HOH:O	1.77	1.01
1:E:177[A]:ARG:NH2	7:E:403:HOH:O	1.97	0.98
1:A:53[A]:ARG:NH1	7:A:403:HOH:O	1.98	0.94
1:B:97[B]:ILE:HD11	1:B:194:PHE:CD2	2.03	0.94

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	299/253 (118%)	296 (99%)	2 (1%)	1 (0%)	36	10
1	B	303/253 (120%)	300 (99%)	2 (1%)	1 (0%)	36	10
1	C	315/253 (124%)	309 (98%)	5 (2%)	1 (0%)	36	10
1	D	307/253 (121%)	303 (99%)	3 (1%)	1 (0%)	36	10
1	E	312/253 (123%)	309 (99%)	2 (1%)	1 (0%)	36	10
1	F	304/253 (120%)	301 (99%)	2 (1%)	1 (0%)	36	10
All	All	1840/1518 (121%)	1818 (99%)	16 (1%)	6 (0%)	30	10

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	162	TYR
1	A	162	TYR
1	B	162	TYR
1	C	162	TYR
1	D	162	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	245/203 (121%)	244 (100%)	1 (0%)	84	63

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	247/203 (122%)	244 (99%)	3 (1%)	63	26
1	C	258/203 (127%)	251 (97%)	7 (3%)	39	5
1	D	250/203 (123%)	249 (100%)	1 (0%)	84	63
1	E	259/203 (128%)	253 (98%)	6 (2%)	44	8
1	F	253/203 (125%)	253 (100%)	0	100	100
All	All	1512/1218 (124%)	1494 (99%)	18 (1%)	68	26

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	232[B]	THR
1	E	253	LYS
1	E	235[B]	GLU
1	C	242[A]	LYS
1	E	232[A]	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	102	ASN
1	C	180	GLN
1	F	144	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 17 ligands modelled in this entry, 10 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	URI	A	301	-	18,18,18	0.99	1 (5%)	26,26,26	1.35	4 (15%)
2	URI	B	301	-	18,18,18	0.78	0	26,26,26	1.74	6 (23%)
2	URI	E	301	-	18,18,18	0.84	0	26,26,26	1.62	5 (19%)
2	URI	F	301	-	18,18,18	0.74	1 (5%)	26,26,26	1.51	6 (23%)
2	URI	C	301	-	18,18,18	0.90	1 (5%)	26,26,26	1.38	3 (11%)
6	EDO	E	303	-	3,3,3	0.27	0	2,2,2	0.17	0
2	URI	D	301	-	18,18,18	1.31	1 (5%)	26,26,26	1.43	4 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	URI	A	301	-	-	4/6/22/22	0/2/2/2
2	URI	B	301	-	-	2/6/22/22	0/2/2/2
2	URI	E	301	-	-	4/6/22/22	0/2/2/2
2	URI	F	301	-	-	4/6/22/22	0/2/2/2
2	URI	C	301	-	-	4/6/22/22	0/2/2/2
6	EDO	E	303	-	-	1/1/1/1	-
2	URI	D	301	-	-	4/6/22/22	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	301	URI	O2-C2	-5.00	1.14	1.23
2	A	301	URI	O2-C2	-3.33	1.17	1.23
2	C	301	URI	O2-C2	-3.10	1.17	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	301	URI	O2-C2	-2.27	1.19	1.23

The worst 5 of 28 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	URI	C1'-N1-C2	4.63	125.92	117.59
2	E	301	URI	C1'-N1-C2	4.15	125.05	117.59
2	B	301	URI	O4'-C1'-N1	-3.94	99.44	108.36
2	F	301	URI	C1'-N1-C2	3.90	124.59	117.59
2	E	301	URI	C2'-C1'-N1	3.46	122.88	113.25

There are no chirality outliers.

5 of 23 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	301	URI	C2'-C1'-N1-C6
2	B	301	URI	C2'-C1'-N1-C2
2	E	301	URI	C2'-C1'-N1-C2
2	E	301	URI	C2'-C1'-N1-C6
2	F	301	URI	C2'-C1'-N1-C6

There are no ring outliers.

3 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	301	URI	1	0
2	E	301	URI	2	0
6	E	303	EDO	6	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	251/253 (99%)	-0.27	6 (2%) 59 71	4, 11, 24, 48	33 (13%)
1	B	252/253 (99%)	-0.26	6 (2%) 59 71	4, 10, 20, 45	34 (13%)
1	C	251/253 (99%)	0.18	19 (7%) 20 22	4, 14, 38, 75	39 (15%)
1	D	252/253 (99%)	-0.06	8 (3%) 50 63	4, 12, 31, 54	36 (14%)
1	E	252/253 (99%)	-0.01	15 (5%) 27 32	4, 11, 26, 51	42 (16%)
1	F	251/253 (99%)	-0.23	3 (1%) 76 86	4, 11, 25, 49	33 (13%)
All	All	1509/1518 (99%)	-0.11	57 (3%) 44 55	4, 11, 28, 75	217 (14%)

The worst 5 of 57 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	2	THR	7.3
1	B	2	THR	5.9
1	E	232[A]	THR	5.1
1	C	227	ILE	4.7
1	E	231[A]	ALA	4.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	EDO	E	303	4/4	0.89	0.14	31,37,38,39	10
4	CL	B	302	1/1	0.96	0.14	35,35,35,35	0
2	URI	D	301	17/17	0.97	0.07	9,13,16,18	29
2	URI	B	301	17/17	0.97	0.06	7,10,18,21	2
4	CL	D	302	1/1	0.97	0.07	19,19,19,19	1
5	MG	D	304	1/1	0.97	0.09	16,16,16,16	1
2	URI	C	301	17/17	0.97	0.06	11,14,17,20	29
2	URI	F	301	17/17	0.98	0.05	9,13,18,19	4
4	CL	D	303	1/1	0.98	0.06	32,32,32,32	1
2	URI	A	301	17/17	0.98	0.05	8,11,15,17	29
4	CL	C	302	1/1	0.98	0.06	28,28,28,28	1
4	CL	C	303	1/1	0.99	0.06	19,19,19,19	1
3	NA	C	304	1/1	0.99	0.04	15,15,15,15	1
3	NA	E	302	1/1	0.99	0.05	13,13,13,13	1
4	CL	F	302	1/1	0.99	0.06	21,21,21,21	1
2	URI	E	301	17/17	0.99	0.04	7,10,15,16	7
3	NA	A	302	1/1	0.99	0.04	10,10,10,10	1

6.5 Other polymers [i](#)

There are no such residues in this entry.