



Full wwPDB NMR Structure Validation Report ⓘ

Mar 26, 2026 – 04:14 AM UTC

PDB ID : 2M3L / pdb_00002m3l
BMRB ID : 18967
Title : Solution structure of the C-terminal zinc-binding domain of HPV51 oncoprotein E6
Authors : Mischo, A.; Ohlenschlager, O.; Gorlach, M.
Deposited on : 2013-01-21

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 90%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	76	71% 9% 20%

2 Ensemble composition and analysis

This entry contains 20 models. Model 18 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:80-A:140 (61)	0.74	18

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 3 single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 4, 5, 6, 8, 12, 14, 16, 18, 20
2	7, 9, 13, 15, 19
Single-model clusters	10; 11; 17

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1246 atoms, of which 620 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Protein E6.

Mol	Chain	Residues	Atoms						Trace
1	A	76	Total	C	H	N	O	S	0
			1245	381	620	125	114	5	

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	76	GLY	-	expression tag	UNP P26554
A	77	SER	-	expression tag	UNP P26554
A	78	HIS	-	expression tag	UNP P26554
A	79	MET	-	expression tag	UNP P26554

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	
2	A	1	Total	Zn
			1	1

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Protein E6




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Protein E6




4.2.2 Score per residue for model 2

- Molecule 1: Protein E6




4.2.3 Score per residue for model 3

- Molecule 1: Protein E6




4.2.4 Score per residue for model 4

- Molecule 1: Protein E6




4.2.5 Score per residue for model 5

- Molecule 1: Protein E6




4.2.6 Score per residue for model 6

- Molecule 1: Protein E6




4.2.7 Score per residue for model 7

- Molecule 1: Protein E6




4.2.8 Score per residue for model 8

- Molecule 1: Protein E6




4.2.9 Score per residue for model 9

- Molecule 1: Protein E6




4.2.10 Score per residue for model 10

- Molecule 1: Protein E6




4.2.11 Score per residue for model 11

- Molecule 1: Protein E6




4.2.12 Score per residue for model 12

- Molecule 1: Protein E6




4.2.13 Score per residue for model 13

- Molecule 1: Protein E6




4.2.14 Score per residue for model 14

- Molecule 1: Protein E6




4.2.15 Score per residue for model 15

- Molecule 1: Protein E6




4.2.16 Score per residue for model 16

- Molecule 1: Protein E6




4.2.17 Score per residue for model 17

- Molecule 1: Protein E6




4.2.18 Score per residue for model 18 (medoid)

- Molecule 1: Protein E6




4.2.19 Score per residue for model 19

- Molecule 1: Protein E6



4.2.20 Score per residue for model 20

- Molecule 1: Protein E6



5 Refinement protocol and experimental data overview

The models were refined using the following method: *DGSA-distance geometry simulated annealing, molecular dynamics*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	3
CNS	refinement	
CYANA	refinement	3

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	938
Number of shifts mapped to atoms	938
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	90%

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.13±0.04	0±1/509 (0.0± 0.1%)	1.05±0.04	0±0/684 (0.0± 0.0%)
All	All	1.13	5/10180 (0.0%)	1.06	1/13680 (0.0%)

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	94	LYS	CA-C	-5.33	1.46	1.52	12	1
1	A	132	TRP	C-N	-5.26	1.26	1.33	6	1
1	A	108	ARG	NE-CZ	5.24	1.38	1.33	16	2
1	A	116	GLN	C-O	-5.16	1.18	1.24	1	1

All unique angle outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	120	ASP	CA-CB-CG	5.71	118.31	112.60	4	1

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	498	496	496	6±2
All	All	9980	9920	9920	123

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:89:GLU:O	1:A:93:LYS:HA	0.64	1.93	8	6
1:A:114:GLU:O	1:A:118:LEU:HG	0.61	1.95	10	9
1:A:89:GLU:OE2	1:A:95:SER:HA	0.60	1.96	20	3
1:A:131:ARG:HA	1:A:131:ARG:NE	0.60	2.11	19	2
1:A:80:SER:HA	1:A:127:GLU:O	0.59	1.98	10	13
1:A:114:GLU:OE2	1:A:136:CYS:HB2	0.58	1.99	6	5
1:A:81:ARG:HB3	1:A:127:GLU:HB3	0.56	1.77	13	3
1:A:136:CYS:SG	1:A:138:ASN:HB2	0.56	2.40	9	1
1:A:89:GLU:HA	1:A:94:LYS:O	0.56	2.01	2	8
1:A:114:GLU:CG	1:A:136:CYS:HA	0.56	2.31	15	2
1:A:110:LEU:HA	1:A:114:GLU:OE2	0.56	2.01	15	1
1:A:105:ARG:HD3	1:A:139:CYS:SG	0.56	2.41	20	1
1:A:110:LEU:O	1:A:115:LYS:HE3	0.55	2.01	13	3
1:A:93:LYS:HB3	1:A:93:LYS:NZ	0.54	2.17	14	2
1:A:93:LYS:NZ	1:A:93:LYS:HB2	0.54	2.18	1	1
1:A:105:ARG:HD3	1:A:105:ARG:O	0.52	2.04	13	3
1:A:103:CYS:HA	1:A:134:GLY:O	0.51	2.06	4	6
1:A:136:CYS:SG	1:A:138:ASN:HB3	0.50	2.47	6	1
1:A:99:LEU:HD21	1:A:132:TRP:CD1	0.49	2.43	16	1
1:A:127:GLU:HB2	1:A:132:TRP:CZ3	0.49	2.42	4	1
1:A:114:GLU:OE1	1:A:136:CYS:HB2	0.49	2.08	2	2
1:A:114:GLU:HG3	1:A:136:CYS:HA	0.49	1.84	12	1
1:A:114:GLU:HG2	1:A:136:CYS:HA	0.48	1.85	19	1
1:A:102:ARG:HD3	1:A:107:GLN:O	0.47	2.09	8	3
1:A:110:LEU:O	1:A:115:LYS:HE2	0.47	2.08	3	1
1:A:82:SER:HA	1:A:125:PHE:O	0.47	2.10	1	7
1:A:135:GLN:HG2	1:A:140:TRP:HB2	0.46	1.86	9	1
1:A:122:LYS:N	1:A:122:LYS:HD2	0.45	2.26	11	3
1:A:113:GLU:H	1:A:113:GLU:CD	0.45	2.18	7	3
1:A:85:GLY:O	1:A:89:GLU:HG3	0.44	2.12	9	3
1:A:121:GLU:HB3	1:A:123:LYS:NZ	0.44	2.28	20	5
1:A:80:SER:HB3	1:A:126:HIS:NE2	0.44	2.27	9	1
1:A:101:ILE:O	1:A:110:LEU:HG	0.44	2.13	2	1
1:A:117:LYS:HA	1:A:120:ASP:HB3	0.43	1.89	4	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:80:SER:HB2	1:A:126:HIS:NE2	0.43	2.28	20	1
1:A:137:ALA:HA	1:A:140:TRP:HB3	0.43	1.91	18	2
1:A:114:GLU:HG3	1:A:136:CYS:CA	0.42	2.44	12	1
1:A:85:GLY:CA	1:A:96:LEU:HD21	0.42	2.44	2	1
1:A:114:GLU:HG2	1:A:136:CYS:CB	0.42	2.45	4	1
1:A:94:LYS:HD2	1:A:98:ASP:OD1	0.42	2.14	11	1
1:A:111:GLY:O	1:A:115:LYS:HG2	0.42	2.15	6	1
1:A:107:GLN:HA	1:A:107:GLN:OE1	0.42	2.15	18	1
1:A:102:ARG:O	1:A:133:THR:HA	0.41	2.15	5	1
1:A:101:ILE:HA	1:A:132:TRP:O	0.41	2.14	16	1
1:A:117:LYS:HA	1:A:120:ASP:CB	0.41	2.45	4	1
1:A:81:ARG:HD3	1:A:127:GLU:OE1	0.41	2.14	1	1
1:A:93:LYS:HE2	1:A:93:LYS:HA	0.41	1.92	9	1
1:A:81:ARG:HD2	1:A:127:GLU:OE1	0.41	2.15	10	1
1:A:103:CYS:SG	1:A:105:ARG:HB3	0.40	2.56	12	1
1:A:117:LYS:CE	1:A:117:LYS:HA	0.40	2.46	17	1
1:A:114:GLU:CD	1:A:136:CYS:HB2	0.40	2.42	1	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	61/76 (80%)	59±1 (97±2%)	2±1 (3±2%)	0±0 (0±0%)	49	83
All	All	1220/1520 (80%)	1182 (97%)	37 (3%)	1 (0%)	49	83

All 1 unique Ramachandran outliers are listed below.

Mol	Chain	Res	Type	Models (Total)
1	A	112	PRO	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR

entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	54/68 (79%)	52±1 (95±2%)	2±1 (5±2%)	25	76
All	All	1080/1360 (79%)	1030 (95%)	50 (5%)	25	76

All 13 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	93	LYS	11
1	A	123	LYS	6
1	A	117	LYS	5
1	A	122	LYS	4
1	A	131	ARG	4
1	A	116	GLN	4
1	A	128	ILE	4
1	A	99	LEU	3
1	A	127	GLU	3
1	A	113	GLU	2
1	A	105	ARG	2
1	A	88	LEU	1
1	A	126	HIS	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 90% for the well-defined parts and 87% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	938
Number of shifts mapped to atoms	938
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	11

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	74	0.09 ± 0.26	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	70	0.07 ± 0.37	None needed (< 0.5 ppm)
$^{13}\text{C}'$	74	0.06 ± 0.25	None needed (< 0.5 ppm)
^{15}N	70	-0.22 ± 0.45	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 90%, i.e. 772 atoms were assigned a chemical shift out of a possible 860. 0 out of 7 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	303/305 (99%)	123/124 (99%)	122/122 (100%)	58/59 (98%)
Sidechain	419/489 (86%)	283/313 (90%)	132/148 (89%)	4/28 (14%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	50/66 (76%)	23/33 (70%)	25/29 (86%)	2/4 (50%)
Overall	772/860 (90%)	429/470 (91%)	279/299 (93%)	64/91 (70%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 87%, i.e. 936 atoms were assigned a chemical shift out of a possible 1079. 0 out of 8 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	366/381 (96%)	148/155 (95%)	148/152 (97%)	70/74 (95%)
Sidechain	516/625 (83%)	348/398 (87%)	160/186 (86%)	8/41 (20%)
Aromatic	54/73 (74%)	25/37 (68%)	27/31 (87%)	2/5 (40%)
Overall	936/1079 (87%)	521/590 (88%)	335/369 (91%)	80/120 (67%)

7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	127	GLU	HB3	-0.29	0.95 – 3.05	-10.9
1	A	127	GLU	HG3	0.27	1.20 – 3.30	-9.4
1	A	88	LEU	HD21	-1.30	-0.65 – 2.13	-7.3
1	A	88	LEU	HD22	-1.30	-0.65 – 2.13	-7.3
1	A	88	LEU	HD23	-1.30	-0.65 – 2.13	-7.3
1	A	132	TRP	CE2	175.42	105.29 – 170.61	5.7
1	A	127	GLU	HG2	1.10	1.24 – 3.30	-5.7
1	A	140	TRP	CE2	174.33	105.29 – 170.61	5.6
1	A	88	LEU	HD11	-0.63	-0.61 – 2.12	-5.1
1	A	88	LEU	HD12	-0.63	-0.61 – 2.12	-5.1
1	A	88	LEU	HD13	-0.63	-0.61 – 2.12	-5.1

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:

