



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 04:20 AM UTC

PDB ID : 3M3I / pdb_00003m3i
Title : Hypothetical protein from Leishmania major
Authors : Merritt, E.A.; Structural Genomics of Pathogenic Protozoa Consortium (SGPP)
Deposited on : 2010-03-09
Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1596 (2.36-2.36)
Clashscore	190562	1663 (2.36-2.36)
Ramachandran outliers	187476	1646 (2.36-2.36)
Sidechain outliers	187428	1646 (2.36-2.36)
RSRZ outliers	180081	1598 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	225	32% 67% 11% 21%
1	B	225	35% 64% 14% 21%
1	C	225	22% 64% 14% 20%
1	D	225	20% 65% 13% 20%
1	E	225	24% 67% 12% 21%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	225	
1	G	225	
1	H	225	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 11442 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative uncharacterized protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	177	Total	C	N	O	S	0	0	0
			1404	893	245	260	6			
1	B	177	Total	C	N	O	S	0	1	0
			1424	904	245	269	6			
1	C	179	Total	C	N	O	S	0	1	0
			1431	909	250	266	6			
1	D	179	Total	C	N	O	S	0	0	0
			1425	906	248	265	6			
1	E	177	Total	C	N	O	S	0	1	0
			1413	898	247	262	6			
1	F	176	Total	C	N	O	S	0	1	0
			1401	893	241	261	6			
1	G	176	Total	C	N	O	S	0	1	0
			1407	896	244	261	6			
1	H	177	Total	C	N	O	S	0	1	0
			1416	898	247	265	6			

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	expression tag	UNP Q4FX13
A	-6	ALA	-	expression tag	UNP Q4FX13
A	-5	HIS	-	expression tag	UNP Q4FX13
A	-4	HIS	-	expression tag	UNP Q4FX13
A	-3	HIS	-	expression tag	UNP Q4FX13
A	-2	HIS	-	expression tag	UNP Q4FX13
A	-1	HIS	-	expression tag	UNP Q4FX13
A	0	HIS	-	expression tag	UNP Q4FX13
B	-7	MET	-	expression tag	UNP Q4FX13
B	-6	ALA	-	expression tag	UNP Q4FX13
B	-5	HIS	-	expression tag	UNP Q4FX13
B	-4	HIS	-	expression tag	UNP Q4FX13
B	-3	HIS	-	expression tag	UNP Q4FX13

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	HIS	-	expression tag	UNP Q4FX13
B	-1	HIS	-	expression tag	UNP Q4FX13
B	0	HIS	-	expression tag	UNP Q4FX13
C	-7	MET	-	expression tag	UNP Q4FX13
C	-6	ALA	-	expression tag	UNP Q4FX13
C	-5	HIS	-	expression tag	UNP Q4FX13
C	-4	HIS	-	expression tag	UNP Q4FX13
C	-3	HIS	-	expression tag	UNP Q4FX13
C	-2	HIS	-	expression tag	UNP Q4FX13
C	-1	HIS	-	expression tag	UNP Q4FX13
C	0	HIS	-	expression tag	UNP Q4FX13
D	-7	MET	-	expression tag	UNP Q4FX13
D	-6	ALA	-	expression tag	UNP Q4FX13
D	-5	HIS	-	expression tag	UNP Q4FX13
D	-4	HIS	-	expression tag	UNP Q4FX13
D	-3	HIS	-	expression tag	UNP Q4FX13
D	-2	HIS	-	expression tag	UNP Q4FX13
D	-1	HIS	-	expression tag	UNP Q4FX13
D	0	HIS	-	expression tag	UNP Q4FX13
E	-7	MET	-	expression tag	UNP Q4FX13
E	-6	ALA	-	expression tag	UNP Q4FX13
E	-5	HIS	-	expression tag	UNP Q4FX13
E	-4	HIS	-	expression tag	UNP Q4FX13
E	-3	HIS	-	expression tag	UNP Q4FX13
E	-2	HIS	-	expression tag	UNP Q4FX13
E	-1	HIS	-	expression tag	UNP Q4FX13
E	0	HIS	-	expression tag	UNP Q4FX13
F	-7	MET	-	expression tag	UNP Q4FX13
F	-6	ALA	-	expression tag	UNP Q4FX13
F	-5	HIS	-	expression tag	UNP Q4FX13
F	-4	HIS	-	expression tag	UNP Q4FX13
F	-3	HIS	-	expression tag	UNP Q4FX13
F	-2	HIS	-	expression tag	UNP Q4FX13
F	-1	HIS	-	expression tag	UNP Q4FX13
F	0	HIS	-	expression tag	UNP Q4FX13
G	-7	MET	-	expression tag	UNP Q4FX13
G	-6	ALA	-	expression tag	UNP Q4FX13
G	-5	HIS	-	expression tag	UNP Q4FX13
G	-4	HIS	-	expression tag	UNP Q4FX13
G	-3	HIS	-	expression tag	UNP Q4FX13
G	-2	HIS	-	expression tag	UNP Q4FX13
G	-1	HIS	-	expression tag	UNP Q4FX13

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	0	HIS	-	expression tag	UNP Q4FX13
H	-7	MET	-	expression tag	UNP Q4FX13
H	-6	ALA	-	expression tag	UNP Q4FX13
H	-5	HIS	-	expression tag	UNP Q4FX13
H	-4	HIS	-	expression tag	UNP Q4FX13
H	-3	HIS	-	expression tag	UNP Q4FX13
H	-2	HIS	-	expression tag	UNP Q4FX13
H	-1	HIS	-	expression tag	UNP Q4FX13
H	0	HIS	-	expression tag	UNP Q4FX13


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	15	Total O 15 15	0	0
2	B	18	Total O 18 18	0	0
2	C	19	Total O 19 19	0	0
2	D	22	Total O 22 22	0	0
2	E	17	Total O 17 17	0	0
2	F	10	Total O 10 10	0	0
2	G	6	Total O 6 6	0	0
2	H	14	Total O 14 14	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Putative uncharacterized protein




• Molecule 1: Putative uncharacterized protein




• Molecule 1: Putative uncharacterized protein







● Molecule 1: Putative uncharacterized protein




● Molecule 1: Putative uncharacterized protein




● Molecule 1: Putative uncharacterized protein



● Molecule 1: Putative uncharacterized protein



● Molecule 1: Putative uncharacterized protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	52.98Å 98.43Å 100.76Å 71.87° 80.19° 89.57°	Depositor
Resolution (Å)	46.72 – 2.35 46.72 – 2.35	Depositor EDS
% Data completeness (in resolution range)	92.8 (46.72-2.35) 92.8 (46.72-2.35)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.72 (at 2.34Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.239 , 0.290 0.249 , 0.294	Depositor DCC
R_{free} test set	3684 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	35.1	Xtriage
Anisotropy	0.280	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 52.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	11442	wwPDB-VP
Average B, all atoms (Å ²)	74.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.66	0/1442	0.81	2/1959 (0.1%)
1	B	0.61	0/1463	0.77	1/1989 (0.1%)
1	C	0.70	0/1471	0.81	0/2000
1	D	0.68	0/1465	0.82	0/1991
1	E	0.67	0/1451	0.80	1/1971 (0.1%)
1	F	0.63	0/1440	0.81	1/1959 (0.1%)
1	G	0.61	0/1446	0.78	0/1966
1	H	0.61	0/1454	0.77	0/1976
All	All	0.65	0/11632	0.80	5/15811 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	173	GLY	N-CA-C	7.73	120.82	111.93
1	B	173	GLY	N-CA-C	6.29	119.25	112.33
1	A	31	TYR	N-CA-C	5.57	116.73	108.60
1	E	173	GLY	N-CA-C	5.16	118.01	112.33
1	A	28	GLY	N-CA-C	-5.15	105.71	111.63

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1404	0	1328	19	0
1	B	1424	0	1338	25	0
1	C	1431	0	1351	27	0
1	D	1425	0	1351	29	0
1	E	1413	0	1335	20	0
1	F	1401	0	1312	20	0
1	G	1407	0	1323	28	0
1	H	1416	0	1332	20	0
2	A	15	0	0	2	0
2	B	18	0	0	1	0
2	C	19	0	0	3	0
2	D	22	0	0	0	0
2	E	17	0	0	0	0
2	F	10	0	0	1	0
2	G	6	0	0	0	0
2	H	14	0	0	1	0
All	All	11442	0	10670	156	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 156 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:169:ILE:CD1	1:H:169:ILE:HD12	1.85	1.06
1:A:169:ILE:HD12	1:H:169:ILE:HD12	1.06	1.06
1:E:195:GLU:OE1	1:F:19:ARG:NH1	1.95	0.99
1:A:169:ILE:HD12	1:H:169:ILE:CD1	1.95	0.96
1:D:77:MET:HE3	1:G:77:MET:HE3	1.48	0.93

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	173/225 (77%)	167 (96%)	5 (3%)	1 (1%)	21	24
1	B	174/225 (77%)	169 (97%)	4 (2%)	1 (1%)	21	24
1	C	176/225 (78%)	172 (98%)	3 (2%)	1 (1%)	21	24
1	D	175/225 (78%)	168 (96%)	6 (3%)	1 (1%)	21	24
1	E	174/225 (77%)	167 (96%)	6 (3%)	1 (1%)	21	24
1	F	173/225 (77%)	167 (96%)	5 (3%)	1 (1%)	21	24
1	G	173/225 (77%)	168 (97%)	4 (2%)	1 (1%)	21	24
1	H	174/225 (77%)	169 (97%)	4 (2%)	1 (1%)	21	24
All	All	1392/1800 (77%)	1347 (97%)	37 (3%)	8 (1%)	21	24

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	93	PRO
1	A	93	PRO
1	E	93	PRO
1	H	93	PRO
1	D	93	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	145/188 (77%)	142 (98%)	3 (2%)	47	61
1	B	149/188 (79%)	144 (97%)	5 (3%)	32	43
1	C	149/188 (79%)	146 (98%)	3 (2%)	48	63
1	D	149/188 (79%)	145 (97%)	4 (3%)	39	52
1	E	146/188 (78%)	141 (97%)	5 (3%)	32	43
1	F	144/188 (77%)	141 (98%)	3 (2%)	47	61
1	G	145/188 (77%)	141 (97%)	4 (3%)	38	51
1	H	147/188 (78%)	146 (99%)	1 (1%)	76	86

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1174/1504 (78%)	1146 (98%)	28 (2%)	43 57

5 of 28 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	169	ILE
1	H	77	MET
1	E	71	CYS
1	G	71	CYS
1	E	32	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	25	HIS
1	G	43	ASN
1	H	121	GLN
1	G	68	HIS
1	C	10	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	177/225 (78%)	2.04	73 (41%) 0 0	57, 69, 99, 135	0
1	B	177/225 (78%)	2.09	78 (44%) 0 0	37, 73, 101, 162	1 (0%)
1	C	179/225 (79%)	1.65	50 (27%) 1 1	32, 65, 83, 101	1 (0%)
1	D	179/225 (79%)	1.59	45 (25%) 1 1	57, 65, 84, 113	0
1	E	177/225 (78%)	1.70	53 (29%) 1 1	32, 66, 84, 124	1 (0%)
1	F	176/225 (78%)	2.23	85 (48%) 0 0	37, 74, 113, 205	1 (0%)
1	G	176/225 (78%)	2.44	103 (58%) 0 0	38, 77, 121, 180	1 (0%)
1	H	177/225 (78%)	2.16	88 (49%) 0 0	35, 74, 107, 142	1 (0%)
All	All	1418/1800 (78%)	1.99	575 (40%) 0 0	32, 70, 103, 205	6 (0%)

The worst 5 of 575 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	98	ARG	9.1
1	G	98	ARG	7.8
1	B	176[A]	TYR	7.8
1	G	176[A]	TYR	6.9
1	G	97	ASP	6.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.