



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 1, 2026 – 02:25 PM UTC

PDB ID : 6M36 / pdb_00006m36
Title : The crystal structure of B. subtilis RsbV/RsbW complex in the monoclinic crystal form
Authors : Pathak, D.; Kwon, E.; Kim, D.Y.
Deposited on : 2020-03-02
Resolution : 3.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1001 (3.44-3.36)
Clashscore	190562	1022 (3.44-3.36)
Ramachandran outliers	187476	1012 (3.44-3.36)
Sidechain outliers	187428	1012 (3.44-3.36)
RSRZ outliers	180081	1001 (3.44-3.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	141	
1	C	141	
1	E	141	
1	G	141	
1	I	141	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	K	141	
1	M	141	
1	O	141	
2	B	103	
2	D	103	
2	F	103	
2	H	103	
2	J	103	
2	L	103	
2	N	103	
2	P	103	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 12580 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine-protein kinase RsbW.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	103	Total	C	N	O	S	0	0	0
			802	508	131	157	6			
1	C	105	Total	C	N	O	S	0	0	0
			824	524	133	161	6			
1	E	100	Total	C	N	O	S	0	0	0
			777	493	124	154	6			
1	G	104	Total	C	N	O	S	0	0	0
			814	517	132	159	6			
1	I	104	Total	C	N	O	S	0	0	0
			808	511	132	159	6			
1	K	103	Total	C	N	O	S	0	0	0
			802	508	131	157	6			
1	M	105	Total	C	N	O	S	0	0	0
			824	524	133	161	6			
1	O	98	Total	C	N	O	S	0	0	0
			760	483	123	148	6			

- Molecule 2 is a protein called Anti-sigma-B factor antagonist.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	100	Total	C	N	O	S	0	0	0
			774	490	130	151	3			
2	D	99	Total	C	N	O	S	0	0	0
			768	487	129	149	3			
2	F	95	Total	C	N	O	S	0	0	0
			742	471	124	144	3			
2	H	101	Total	C	N	O	S	0	0	0
			779	493	131	152	3			
2	J	100	Total	C	N	O	S	0	0	0
			774	490	130	151	3			
2	L	101	Total	C	N	O	S	0	0	0
			779	493	131	152	3			


Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	N	101	Total	C	N	O	S	0	0	0
			779	493	131	152	3			
2	P	100	Total	C	N	O	S	0	0	0
			774	490	130	151	3			





- Molecule 1: Serine-protein kinase RsbW





- Molecule 1: Serine-protein kinase RsbW



- Molecule 1: Serine-protein kinase RsbW



- Molecule 1: Serine-protein kinase RsbW



- Molecule 2: Anti-sigma-B factor antagonist

Chain B:  70% 27% .



- Molecule 2: Anti-sigma-B factor antagonist

Chain D:  63% 33% .



- Molecule 2: Anti-sigma-B factor antagonist

Chain F:  66% 26% 8% .




- Molecule 2: Anti-sigma-B factor antagonist

Chain H:  64% 34% .



- Molecule 2: Anti-sigma-B factor antagonist


Chain J:  50% 47% .





- Molecule 2: Anti-sigma-B factor antagonist

Chain L:  54% 44% .







● Molecule 2: Anti-sigma-B factor antagonist



● Molecule 2: Anti-sigma-B factor antagonist



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	117.23Å 70.94Å 137.68Å 90.00° 105.35° 90.00°	Depositor
Resolution (Å)	29.79 – 3.40 29.79 – 3.40	Depositor EDS
% Data completeness (in resolution range)	96.1 (29.79-3.40) 95.9 (29.79-3.40)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.15 (at 3.39Å)	Xtriage
Refinement program	PHENIX dev_3051	Depositor
R, R_{free}	0.207 , 0.290 0.210 , 0.288	Depositor DCC
R_{free} test set	1476 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	71.2	Xtriage
Anisotropy	0.328	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 58.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	12580	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.56	0/810	0.87	0/1092
1	C	0.55	0/833	0.78	0/1122
1	E	0.47	0/783	0.78	0/1055
1	G	0.48	0/823	0.81	1/1110 (0.1%)
1	I	0.57	0/817	0.90	0/1103
1	K	0.52	0/810	0.78	0/1092
1	M	0.44	0/833	0.73	0/1122
1	O	0.48	0/766	0.76	0/1032
2	B	0.53	0/780	0.82	0/1050
2	D	0.48	0/774	0.76	0/1042
2	F	0.43	0/747	0.73	0/1004
2	H	0.53	0/785	0.86	0/1057
2	J	0.53	0/780	0.86	0/1050
2	L	0.42	0/785	0.73	0/1057
2	N	0.56	0/785	0.85	0/1057
2	P	0.51	0/780	0.83	0/1050
All	All	0.51	0/12691	0.80	1/17095 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
2	P	0	1
All	All	0	2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	124	MET	CA-CB-CG	-5.75	102.61	114.10

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	33	MET	Peptide
2	P	9	GLN	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	802	0	795	27	0
1	C	824	0	817	48	0
1	E	777	0	776	41	0
1	G	814	0	804	41	0
1	I	808	0	801	42	0
1	K	802	0	795	32	0
1	M	824	0	817	45	0
1	O	760	0	766	58	0
2	B	774	0	804	19	0
2	D	768	0	799	27	0
2	F	742	0	771	21	0
2	H	779	0	809	31	0
2	J	774	0	804	41	0
2	L	779	0	809	38	0
2	N	779	0	809	36	0
2	P	774	0	804	36	0
All	All	12580	0	12780	507	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 507 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:31:ARG:HG3	2:F:61:VAL:HG23	1.22	1.11
1:M:33:MET:HE1	1:M:78:LEU:HD13	1.49	0.94
1:G:124:MET:HE3	1:G:141:LYS:HB2	1.47	0.94

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:8:ILE:HG12	1:K:12:VAL:HG12	1.48	0.94
1:M:12:VAL:HG11	1:M:18:TYR:HB3	1.53	0.90

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	95/141 (67%)	94 (99%)	1 (1%)	0	100	100
1	C	97/141 (69%)	93 (96%)	4 (4%)	0	100	100
1	E	92/141 (65%)	90 (98%)	2 (2%)	0	100	100
1	G	96/141 (68%)	92 (96%)	3 (3%)	1 (1%)	12	40
1	I	98/141 (70%)	93 (95%)	4 (4%)	1 (1%)	12	40
1	K	95/141 (67%)	94 (99%)	1 (1%)	0	100	100
1	M	97/141 (69%)	95 (98%)	2 (2%)	0	100	100
1	O	90/141 (64%)	88 (98%)	2 (2%)	0	100	100
2	B	98/103 (95%)	93 (95%)	4 (4%)	1 (1%)	12	40
2	D	97/103 (94%)	93 (96%)	4 (4%)	0	100	100
2	F	91/103 (88%)	86 (94%)	4 (4%)	1 (1%)	11	38
2	H	99/103 (96%)	95 (96%)	4 (4%)	0	100	100
2	J	98/103 (95%)	93 (95%)	5 (5%)	0	100	100
2	L	99/103 (96%)	98 (99%)	1 (1%)	0	100	100
2	N	99/103 (96%)	94 (95%)	5 (5%)	0	100	100
2	P	98/103 (95%)	88 (90%)	10 (10%)	0	100	100
All	All	1539/1952 (79%)	1479 (96%)	56 (4%)	4 (0%)	36	65

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	77	ARG
2	B	41	GLY
2	F	74	GLY
1	I	133	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	87/120 (72%)	87 (100%)	0	100	100
1	C	89/120 (74%)	89 (100%)	0	100	100
1	E	85/120 (71%)	85 (100%)	0	100	100
1	G	88/120 (73%)	88 (100%)	0	100	100
1	I	88/120 (73%)	88 (100%)	0	100	100
1	K	87/120 (72%)	87 (100%)	0	100	100
1	M	89/120 (74%)	89 (100%)	0	100	100
1	O	83/120 (69%)	83 (100%)	0	100	100
2	B	88/90 (98%)	88 (100%)	0	100	100
2	D	87/90 (97%)	87 (100%)	0	100	100
2	F	85/90 (94%)	85 (100%)	0	100	100
2	H	88/90 (98%)	88 (100%)	0	100	100
2	J	88/90 (98%)	88 (100%)	0	100	100
2	L	88/90 (98%)	88 (100%)	0	100	100
2	N	88/90 (98%)	88 (100%)	0	100	100
2	P	88/90 (98%)	88 (100%)	0	100	100
All	All	1396/1680 (83%)	1396 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
2	L	72	GLN
2	L	80	ASN
2	P	80	ASN
2	N	4	ASN
2	H	72	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	103/141 (73%)	-0.29	0 100 100	30, 50, 79, 87	0
1	C	105/141 (74%)	-0.34	0 100 100	34, 48, 76, 100	0
1	E	100/141 (70%)	0.35	2 (2%) 65 49	58, 86, 105, 110	0
1	G	104/141 (73%)	-0.02	0 100 100	54, 70, 89, 104	0
1	I	104/141 (73%)	-0.30	0 100 100	33, 49, 74, 84	0
1	K	103/141 (73%)	-0.08	0 100 100	45, 66, 84, 91	0
1	M	105/141 (74%)	0.13	2 (1%) 66 51	48, 75, 101, 125	0
1	O	98/141 (69%)	0.30	1 (1%) 79 66	50, 75, 98, 110	0
2	B	100/103 (97%)	-0.15	0 100 100	40, 65, 93, 116	0
2	D	99/103 (96%)	-0.02	0 100 100	42, 72, 104, 141	0
2	F	95/103 (92%)	0.22	0 100 100	63, 91, 118, 131	0
2	H	101/103 (98%)	-0.33	0 100 100	42, 57, 82, 118	0
2	J	100/103 (97%)	-0.19	1 (1%) 79 66	39, 58, 89, 116	0
2	L	101/103 (98%)	-0.12	0 100 100	47, 71, 94, 113	0
2	N	101/103 (98%)	-0.15	0 100 100	41, 58, 83, 118	0
2	P	100/103 (97%)	-0.16	0 100 100	40, 60, 94, 110	0
All	All	1619/1952 (82%)	-0.08	6 (0%) 88 81	30, 66, 100, 141	0

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	M	76	ASP	2.6
1	E	31	SER	2.6
1	M	143	LEU	2.2
2	J	74	GLY	2.2
1	E	120	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.