



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 03:50 PM UTC

PDB ID : 6M3Y / pdb_00006m3y
Title : Crystal structure of pilus adhesin, SpaC from *Lactobacillus rhamnosus* GG - open conformation
Authors : Kant, A.; Palva, A.; von Ossowski, I.; Krishnan, V.
Deposited on : 2020-03-04
Resolution : 1.93 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1452 (1.94-1.94)
Clashscore	190562	1494 (1.94-1.94)
Ramachandran outliers	187476	1479 (1.94-1.94)
Sidechain outliers	187428	1479 (1.94-1.94)
RSRZ outliers	180081	1453 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	822	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6279 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pilus assembly protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	811	Total	C	N	O	S	0	0	0
			6179	3847	1053	1266	13			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	857	LEU	-	expression tag	UNP A0A1Y0DVK9

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	99	Total	O	0	0
			99	99		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pilus assembly protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	81.05Å 110.02Å 128.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	83.53 – 1.93 83.53 – 1.93	Depositor EDS
% Data completeness (in resolution range)	59.8 (83.53-1.93) 60.1 (83.53-1.93)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.31 (at 1.92Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.207 , 0.248 0.217 , 0.255	Depositor DCC
R_{free} test set	2597 reflections (2.97%)	wwPDB-VP
Wilson B-factor (Å ²)	29.4	Xtriage
Anisotropy	0.080	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 32.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6279	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.66	0/6297	0.89	2/8569 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	708	ASP	N-CA-C	6.15	118.78	111.71
1	A	39	ILE	N-CA-C	5.97	118.20	111.88

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6179	0	5920	25	0
2	A	1	0	0	0	0
3	A	99	0	0	0	0
All	All	6279	0	5920	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:608:ASP:OD1	1:A:610:THR:HG22	2.07	0.54
1:A:49:GLY:HA2	1:A:465:PHE:CZ	2.46	0.51
1:A:753:ALA:HB3	1:A:793:LYS:HB3	1.93	0.51
1:A:342:MET:HG2	1:A:346:LYS:HB2	1.94	0.49
1:A:783:ALA:HB2	1:A:813:GLY:HA3	1.95	0.48
1:A:433:LEU:C	1:A:433:LEU:HD23	2.39	0.47
1:A:138:ILE:HD13	1:A:252:MET:HB3	1.96	0.47
1:A:563:TYR:HD2	1:A:584:MET:HE3	1.79	0.47
1:A:36:THR:HG22	1:A:72:GLN:NE2	2.30	0.47
1:A:783:ALA:HB2	1:A:813:GLY:CA	2.46	0.45
1:A:136:VAL:HG23	1:A:250:LYS:HB3	1.98	0.45
1:A:820:ILE:HG21	1:A:831:LYS:HB3	2.01	0.43
1:A:45:THR:HG22	1:A:50:THR:HG23	2.00	0.43
1:A:331:LEU:HD11	1:A:364:ALA:HB2	2.02	0.42
1:A:684:TYR:O	1:A:694:GLY:HA2	2.19	0.42
1:A:736:THR:HG23	1:A:780:LYS:NZ	2.35	0.42
1:A:67:GLN:HG2	1:A:487:GLY:HA3	2.02	0.42
1:A:827:PRO:HG3	1:A:838:ILE:HD12	2.02	0.41
1:A:230:TYR:O	1:A:233:ILE:HG22	2.21	0.41
1:A:44:GLN:HB2	1:A:53:THR:HG22	2.02	0.41
1:A:759:SER:HB2	1:A:786:THR:HB	2.03	0.41
1:A:805:ILE:N	1:A:805:ILE:HD12	2.36	0.41
1:A:132:ASN:O	1:A:132:ASN:CG	2.64	0.40
1:A:735:MET:HE3	1:A:841:SER:HA	2.04	0.40
1:A:855:LEU:HD23	1:A:855:LEU:HA	1.93	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	805/822 (98%)	791 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	665/680 (98%)	659 (99%)	6 (1%)	70	67

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	62	GLN
1	A	111	LYS
1	A	349	GLN
1	A	663	THR
1	A	712	ASP
1	A	773	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	38	ASN
1	A	54	ASN
1	A	349	GLN
1	A	377	GLN
1	A	517	ASN
1	A	521	GLN
1	A	532	ASN
1	A	626	GLN
1	A	639	ASN
1	A	681	GLN
1	A	690	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	811/822 (98%)	0.51	49 (6%)	27 30	18, 37, 65, 90	0

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	246	THR	5.1
1	A	365	ASP	4.8
1	A	245	ASP	4.6
1	A	508	SER	4.4
1	A	570	ALA	4.2
1	A	760	ALA	4.0
1	A	761	GLY	3.9
1	A	569	ASP	3.8
1	A	509	ASN	3.8
1	A	571	SER	3.7
1	A	707	GLY	3.3
1	A	340	HIS	3.3
1	A	36	THR	3.1
1	A	529	LEU	3.1
1	A	573	ILE	3.0
1	A	782	VAL	2.9
1	A	812	THR	2.9
1	A	156	ASN	2.9
1	A	274	ASN	2.8
1	A	855	LEU	2.8
1	A	718	ASP	2.8
1	A	224	ARG	2.8
1	A	406	ASN	2.7
1	A	753	ALA	2.6
1	A	507	LEU	2.5
1	A	786	THR	2.5
1	A	619	ALA	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	612	LEU	2.4
1	A	617	LEU	2.4
1	A	511	SER	2.4
1	A	359	ASP	2.4
1	A	781	LEU	2.3
1	A	810	ALA	2.3
1	A	794	ALA	2.3
1	A	802	PRO	2.3
1	A	752	ALA	2.2
1	A	572	GLY	2.2
1	A	133	VAL	2.2
1	A	416	GLN	2.2
1	A	766	ILE	2.2
1	A	533	TRP	2.1
1	A	608	ASP	2.1
1	A	705	PRO	2.1
1	A	814	LYS	2.1
1	A	618	LYS	2.1
1	A	740	ILE	2.0
1	A	532	ASN	2.0
1	A	614	ALA	2.0
1	A	840	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)


There are no oligosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MG	A	901	1/1	0.98	0.09	28,28,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.