



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 04:45 AM UTC

PDB ID : 7M3I / pdb_00007m3i
Title : Structure of SARS-CoV-2 spike protein receptor binding domain in complex with a neutralizing antibody, CV2-75 Fab
Authors : Hurlburt, N.K.; Pancera, M.
Deposited on : 2021-03-18
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	227	
1	H	227	
2	B	214	
2	L	214	
3	C	273	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	R	273	3% 67% 11% • 21%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9872 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CV2-75 Fab Heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	221	Total	C	N	O	S	0	0	0
			1658	1054	266	334	4			
1	H	205	Total	C	N	O	S	0	0	0
			1561	996	249	312	4			


- Molecule 2 is a protein called CV2-75 Fab Light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	211	Total	C	N	O	S	0	0	0
			1585	990	267	324	4			
2	L	211	Total	C	N	O	S	0	0	0
			1585	990	267	324	4			

- Molecule 3 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	217	Total	C	N	O	S	0	0	0
			1718	1099	290	321	8			
3	R	217	Total	C	N	O	S	0	0	0
			1721	1102	290	321	8			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	R	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	O	0	0
			1	1		
5	C	7	Total	O	0	0
			7	7		
5	H	2	Total	O	0	0
			2	2		
5	L	2	Total	O	0	0
			2	2		
5	R	4	Total	O	0	0
			4	4		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CV2-75 Fab Heavy chain




- Molecule 1: CV2-75 Fab Heavy chain




- Molecule 2: CV2-75 Fab Light chain




- Molecule 2: CV2-75 Fab Light chain






● Molecule 3: Spike protein S1



● Molecule 3: Spike protein S1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	86.37Å 127.52Å 155.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.32 – 2.80 49.32 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.5 (49.32-2.80) 99.4 (49.32-2.80)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.58 (at 2.81Å)	Xtriage
Refinement program	PHENIX 1.17.1 _3660	Depositor
R, R_{free}	0.245 , 0.281 0.249 , 0.285	Depositor DCC
R_{free} test set	2092 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	79.6	Xtriage
Anisotropy	0.433	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 66.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	9872	wwPDB-VP
Average B, all atoms (Å ²)	101.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, PCA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.14	0/1696	0.38	0/2323
1	H	0.14	0/1597	0.39	1/2187 (0.0%)
2	B	0.13	0/1626	0.34	0/2227
2	L	0.14	0/1626	0.37	0/2227
3	C	0.16	0/1765	0.34	0/2401
3	R	0.15	0/1768	0.36	0/2405
All	All	0.15	0/10078	0.36	1/13770 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	H	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	1	PCA	O-C-N	-7.51	110.97	123.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	H	1	PCA	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1658	0	1614	19	0
1	H	1561	0	1513	17	0
2	B	1585	0	1523	17	0
2	L	1585	0	1523	18	0
3	C	1718	0	1645	14	0
3	R	1721	0	1654	18	0
4	C	14	0	13	1	0
4	R	14	0	13	0	0
5	B	1	0	0	0	0
5	C	7	0	0	0	0
5	H	2	0	0	0	0
5	L	2	0	0	0	0
5	R	4	0	0	0	0
All	All	9872	0	9498	99	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (99) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:177:VAL:HG12	2:B:164:THR:HG23	1.63	0.80
3:R:457:ARG:NH1	3:R:467:ASP:OD2	2.23	0.72
2:L:54:PRO:HG2	2:L:57:ILE:HD13	1.75	0.69
1:A:129:VAL:HG22	1:A:150:VAL:HG12	1.75	0.67
1:A:203:ILE:HD13	1:A:218:LYS:HB3	1.77	0.66
1:A:150:VAL:HG11	1:A:206:VAL:HG21	1.79	0.64
2:B:82:GLU:HG3	2:B:106:THR:HA	1.83	0.60
1:H:203:ILE:HG22	1:H:218:LYS:HB3	1.81	0.60
4:C:600:NAG:H83	4:C:600:NAG:H3	1.85	0.59
3:C:321:GLN:HG3	3:C:324:GLU:HG3	1.85	0.59
1:H:151:LYS:HE3	2:L:133:THR:HG21	1.84	0.59
3:C:439:ASN:O	3:C:443:SER:OG	2.20	0.58
1:H:103:GLY:O	1:H:105:LEU:N	2.36	0.58
2:B:151:LYS:HA	2:B:156:PRO:HA	1.85	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:4:LEU:HD22	1:H:24:VAL:HG22	1.86	0.57
1:A:103:GLY:O	1:A:105:LEU:N	2.37	0.57
3:C:358:ILE:HB	3:C:395:VAL:HG23	1.86	0.56
3:C:468:ILE:HD12	3:R:518:LEU:HD21	1.86	0.56
2:L:151:LYS:HA	2:L:156:PRO:HA	1.88	0.56
2:L:169:GLN:NE2	2:L:173:LYS:HB3	2.20	0.55
3:R:352:ALA:HB2	3:R:468:ILE:HG22	1.89	0.55
2:B:168:LYS:HE2	2:B:172:ASN:HA	1.89	0.55
3:R:342:PHE:CE2	3:R:511:VAL:HG11	2.43	0.54
3:R:417:LYS:O	3:R:422:ASN:ND2	2.39	0.54
2:B:12:VAL:HG21	2:B:18:ALA:HB2	1.89	0.53
3:C:523:THR:HG23	3:C:524:VAL:HG23	1.90	0.53
3:R:321:GLN:HG3	3:R:324:GLU:HG3	1.89	0.53
1:A:134:PRO:HG3	1:A:197:LEU:HD11	1.89	0.53
2:L:173:LYS:HG3	2:L:174:TYR:H	1.73	0.53
2:B:34:TRP:HB2	2:B:47:VAL:HB	1.90	0.52
1:A:86:THR:O	1:A:119:VAL:HG21	2.09	0.52
2:L:112:LYS:HG3	2:L:143:PRO:HG2	1.92	0.52
1:H:2:VAL:HA	1:H:26:GLY:HA3	1.93	0.51
1:A:12:VAL:HG21	1:A:18:LEU:HD13	1.92	0.51
1:H:39:GLN:HB2	1:H:45:LEU:HD23	1.91	0.51
3:C:335:LEU:HD23	3:C:362:VAL:HG13	1.93	0.51
3:R:360:ASN:H	3:R:523:THR:HB	1.75	0.51
1:H:18:LEU:HD12	1:H:117:VAL:HG11	1.92	0.50
3:R:461:LEU:HD22	3:R:465:GLU:HB3	1.92	0.50
2:L:173:LYS:HG3	2:L:174:TYR:N	2.27	0.49
3:C:431:GLY:HA2	3:C:515:PHE:CD2	2.46	0.49
1:H:66:ARG:HH21	1:H:82:LEU:HD11	1.78	0.49
1:A:39:GLN:HB2	1:A:45:LEU:HD23	1.93	0.49
1:A:129:VAL:HG21	1:A:206:VAL:HG11	1.93	0.49
1:H:18:LEU:HD21	1:H:20:LEU:HD13	1.95	0.47
1:A:4:LEU:HD22	1:A:24:VAL:HG12	1.97	0.47
2:B:4:LEU:HD11	2:B:89:VAL:HG12	1.97	0.46
2:B:46:VAL:HA	2:B:57:ILE:HG12	1.98	0.46
3:C:431:GLY:HA2	3:C:515:PHE:HD2	1.81	0.46
2:L:39:PRO:HD3	2:L:82:GLU:OE2	2.16	0.46
1:H:96:ALA:HB1	1:H:108:PHE:HB3	1.98	0.46
1:A:127:PRO:HB3	1:A:153:TYR:HB3	1.97	0.46
1:A:146:LEU:HD13	1:A:219:VAL:HG21	1.97	0.45
1:H:171:VAL:HA	1:H:190:VAL:HG13	1.99	0.45
3:R:537:LYS:HA	3:R:537:LYS:HD2	1.63	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:146:VAL:HG12	2:L:199:HIS:HB2	1.99	0.45
2:L:39:PRO:HD3	2:L:82:GLU:OE1	2.16	0.45
3:R:379:CYS:HA	3:R:432:CYS:HA	1.98	0.45
1:A:127:PRO:HD2	1:A:213:THR:HG21	1.99	0.45
3:C:350:VAL:O	3:C:353:TRP:HD1	1.99	0.44
1:H:100:ARG:NH1	1:H:105:LEU:HB3	2.33	0.44
3:C:326:ILE:HD12	3:C:326:ILE:HA	1.88	0.44
1:H:87:ALA:O	1:H:90:THR:HG22	2.16	0.44
2:L:39:PRO:HD3	2:L:82:GLU:CD	2.42	0.43
1:H:18:LEU:HB3	1:H:82:LEU:HB3	1.99	0.43
2:B:35:TYR:HE1	2:B:88:GLN:HB3	1.83	0.43
2:B:20:ILE:HD11	2:B:105:VAL:HG11	2.01	0.42
2:B:169:GLN:HG2	2:B:173:LYS:O	2.19	0.42
3:R:535:LYS:HB2	3:R:535:LYS:HE2	1.72	0.42
1:H:12:VAL:O	1:H:119:VAL:HA	2.20	0.42
2:L:10:VAL:O	2:L:105:VAL:HA	2.19	0.42
2:L:62:SER:O	2:L:72:LEU:HD12	2.19	0.42
3:R:415:THR:HG22	3:R:416:GLY:H	1.85	0.42
3:C:457:ARG:NH1	3:C:459:SER:O	2.51	0.42
2:L:15:GLY:N	2:L:77:VAL:O	2.52	0.42
2:B:38:LYS:HG2	2:B:83:ALA:HB2	2.02	0.42
1:A:68:THR:HB	1:A:81:LYS:HB2	2.00	0.42
2:B:2:TYR:HB2	2:B:91:ASP:OD2	2.20	0.42
2:L:92:SER:HB2	3:R:380:TYR:CD2	2.55	0.42
3:R:350:VAL:HA	3:R:400:PHE:HB2	2.01	0.42
3:R:434:ILE:HB	3:R:511:VAL:HG13	2.00	0.42
1:H:37:ILE:O	1:H:94:TYR:N	2.42	0.41
3:R:533:LEU:HD23	3:R:533:LEU:HA	1.95	0.41
2:L:6:GLN:NE2	2:L:87:CYS:SG	2.93	0.41
1:H:62:SER:HB2	1:H:63:LEU:HD12	2.02	0.41
2:B:31:THR:O	2:B:89:VAL:HG23	2.20	0.41
3:C:455:LEU:HD22	3:C:493:GLN:HG3	2.02	0.41
1:A:18:LEU:O	1:A:81:LYS:HA	2.21	0.41
2:B:6:GLN:NE2	2:B:87:CYS:SG	2.94	0.41
2:L:83:ALA:HB2	2:L:105:VAL:O	2.21	0.41
3:R:326:ILE:HD12	3:R:326:ILE:HA	1.85	0.41
1:A:35:SER:HA	1:A:50:TYR:HA	2.03	0.41
2:B:12:VAL:HG11	2:B:18:ALA:HA	2.02	0.41
2:B:134:LEU:HB2	2:B:180:LEU:HB3	2.02	0.41
1:A:177:VAL:HG22	1:A:185:SER:O	2.20	0.41
3:C:342:PHE:CE2	3:C:511:VAL:HG11	2.55	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:101:PRO:HB3	3:C:374:PHE:O	2.21	0.40
2:L:164:THR:HG22	2:L:177:SER:H	1.86	0.40
3:R:501:ASN:HD22	3:R:501:ASN:HA	1.76	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	219/227 (96%)	204 (93%)	12 (6%)	3 (1%)	9	30
1	H	201/227 (88%)	187 (93%)	11 (6%)	3 (2%)	8	28
2	B	209/214 (98%)	198 (95%)	11 (5%)	0	100	100
2	L	209/214 (98%)	189 (90%)	17 (8%)	3 (1%)	9	30
3	C	215/273 (79%)	198 (92%)	16 (7%)	1 (0%)	24	55
3	R	215/273 (79%)	202 (94%)	13 (6%)	0	100	100
All	All	1268/1428 (89%)	1178 (93%)	80 (6%)	10 (1%)	16	44

All (10) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	155	PRO
1	H	200	GLN
2	L	142	TYR
2	L	191	ARG
1	A	104	ASN
1	A	194	SER
1	H	104	ASN
3	C	533	LEU
2	L	84	ASP
1	A	14	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/197 (97%)	186 (97%)	5 (3%)	40	75
1	H	181/197 (92%)	175 (97%)	6 (3%)	33	69
2	B	179/182 (98%)	178 (99%)	1 (1%)	78	92
2	L	179/182 (98%)	177 (99%)	2 (1%)	65	87
3	C	189/240 (79%)	188 (100%)	1 (0%)	81	93
3	R	190/240 (79%)	183 (96%)	7 (4%)	30	65
All	All	1109/1238 (90%)	1087 (98%)	22 (2%)	48	80

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	GLN
1	A	24	VAL
1	A	132	LEU
1	A	190	VAL
1	A	203	ILE
2	B	164	THR
3	C	395	VAL
1	H	190	VAL
1	H	192	VAL
1	H	197	LEU
1	H	199	THR
1	H	209	LYS
1	H	215	VAL
2	L	74	ILE
2	L	105	VAL
3	R	328	ARG
3	R	368	LEU
3	R	369	TYR
3	R	531	THR
3	R	534	VAL
3	R	535	LYS
3	R	537	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	GLN
2	B	25	ASN
2	B	36	GLN
1	H	5	GLN
2	L	41	GLN
2	L	88	GLN
2	L	199	HIS
3	R	532	ASN
3	R	536	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	PCA	A	1	1	7,8,9	1.92	1 (14%)	9,10,12	2.12	5 (55%)
1	PCA	H	1	1	7,8,9	1.94	1 (14%)	9,10,12	2.20	5 (55%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	A	1	1	-	0/0/11/13	0/1/1/1
1	PCA	H	1	1	-	0/0/11/13	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	1	PCA	CD-N	5.02	1.47	1.34
1	A	1	PCA	CD-N	4.96	1.46	1.34

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	1	PCA	CA-N-CD	-3.12	102.91	113.58
1	A	1	PCA	CA-N-CD	-3.08	103.02	113.58
1	H	1	PCA	OE-CD-CG	-2.96	121.44	126.72
1	A	1	PCA	OE-CD-CG	-2.82	121.68	126.72
1	H	1	PCA	CB-CA-N	2.62	110.46	103.24
1	A	1	PCA	CB-CA-N	2.62	110.45	103.24
1	H	1	PCA	CG-CD-N	2.54	114.61	108.39
1	A	1	PCA	CG-CD-N	2.51	114.52	108.39
1	A	1	PCA	CB-CA-C	-2.36	109.42	112.66
1	H	1	PCA	CB-CA-C	-2.28	109.52	112.66

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	R	600	3	14,14,15	0.36	0	17,19,21	0.54	0
4	NAG	C	600	3	14,14,15	0.54	0	17,19,21	1.36	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	R	600	3	-	2/6/23/26	0/1/1/1
4	NAG	C	600	3	-	6/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	600	NAG	C2-N2-C7	4.53	128.97	122.90
4	C	600	NAG	C1-C2-N2	2.54	114.44	110.43

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	R	600	NAG	O5-C5-C6-O6
4	C	600	NAG	C8-C7-N2-C2
4	C	600	NAG	O7-C7-N2-C2
4	C	600	NAG	C4-C5-C6-O6
4	C	600	NAG	O5-C5-C6-O6
4	R	600	NAG	C4-C5-C6-O6
4	C	600	NAG	C3-C2-N2-C7
4	C	600	NAG	C1-C2-N2-C7

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	600	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	220/227 (96%)	0.35	3 (1%) 73 64	59, 94, 157, 229	0
1	H	204/227 (89%)	0.62	18 (8%) 15 11	58, 103, 178, 188	0
2	B	211/214 (98%)	0.31	3 (1%) 73 64	62, 100, 138, 157	0
2	L	211/214 (98%)	0.74	21 (9%) 12 9	66, 136, 178, 196	0
3	C	217/273 (79%)	0.24	8 (3%) 45 36	47, 77, 134, 160	0
3	R	217/273 (79%)	0.17	7 (3%) 50 40	53, 79, 150, 199	0
All	All	1280/1428 (89%)	0.40	60 (4%) 36 29	47, 94, 168, 229	0

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	L	176	ALA	4.0
3	R	326	ILE	3.9
3	C	326	ILE	3.8
2	L	79	ALA	3.5
2	L	155	SER	3.4
3	R	369	TYR	3.3
1	H	190	VAL	3.3
1	H	122	ALA	3.2
1	H	167	LEU	3.2
1	A	221	PRO	3.1
2	L	109	GLY	3.1
2	L	82	GLU	3.1
2	L	129	ALA	3.0
3	C	534	VAL	3.0
1	A	133	ALA	2.9
2	L	114	ALA	2.8
1	H	158	VAL	2.7
2	L	117	VAL	2.7
1	H	184	TYR	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	L	166	PRO	2.7
3	C	530	SER	2.7
3	R	371	SER	2.7
1	H	189	VAL	2.6
2	L	115	PRO	2.5
2	L	156	PRO	2.5
2	L	142	TYR	2.5
1	H	149	LEU	2.4
3	C	322	PRO	2.4
1	H	160	VAL	2.4
3	R	368	LEU	2.4
1	A	215	VAL	2.4
2	L	174	TYR	2.4
2	L	137	LEU	2.4
2	L	145	ALA	2.3
2	L	118	THR	2.3
2	L	204	VAL	2.3
1	H	192	VAL	2.3
2	B	159	ALA	2.3
1	H	169	SER	2.3
2	L	141	PHE	2.2
3	C	324	GLU	2.2
1	H	148	CYS	2.2
2	L	197	VAL	2.2
2	L	144	GLY	2.2
3	R	372	ALA	2.2
1	H	131	PRO	2.1
2	L	111	PRO	2.1
3	R	322	PRO	2.1
2	B	211	THR	2.1
3	C	323	THR	2.1
3	C	475	ALA	2.1
1	H	31	SER	2.1
1	H	218	LYS	2.1
1	H	165	GLY	2.1
3	C	486	PHE	2.1
1	H	162	TRP	2.1
1	H	219	VAL	2.0
2	B	154	SER	2.0
3	R	342	PHE	2.0
1	H	214	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	PCA	A	1	8/9	0.50	0.14	128,145,147,148	0
1	PCA	H	1	8/9	0.77	0.12	123,127,132,141	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	C	600	14/15	0.72	0.15	68,94,108,111	0
4	NAG	R	600	14/15	0.85	0.12	90,108,118,118	0

6.5 Other polymers [i](#)

There are no such residues in this entry.