wwPDB EM Validation Summary Report

Mar 8, 2026 – 02:15 AM UTC

PDB ID	:	9M4W / pdb_00009m4w
EMDB ID	:	EMD-63631
Title	:	Structure of human TRPC5 bound with (-)-englerin A, class 2.
Authors	:	Chen, Y.K.; Wei, M.; Chen, L.
Deposited on	:	2025-03-05
Resolution	:	2.62 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev132
Mogul	:	2022.3.0, CSD as543be (2022)
MolProbity	:	4-5-2 with Phenix2.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics	:	202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	8810 (2.12 - 3.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	767	26% 69% 17% 14%
1	B	767	26% 68% 18% 14%
1	C	767	26% 68% 18% 14%
1	D	767	26% 68% 18% 14%

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 22180 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Short transient receptor potential channel 5.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	662	Total 5399	C 3523	N 893	O 953	S 30	0	0
1	B	662	Total 5399	C 3523	N 893	O 953	S 30	0	0
1	C	662	Total 5399	C 3523	N 893	O 953	S 30	0	0
1	D	662	Total 5399	C 3523	N 893	O 953	S 30	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	PRO	-	expression tag	UNP Q9UL62
A	-1	GLU	-	expression tag	UNP Q9UL62
A	0	PHE	-	expression tag	UNP Q9UL62
B	-2	PRO	-	expression tag	UNP Q9UL62
B	-1	GLU	-	expression tag	UNP Q9UL62
B	0	PHE	-	expression tag	UNP Q9UL62
C	-2	PRO	-	expression tag	UNP Q9UL62
C	-1	GLU	-	expression tag	UNP Q9UL62
C	0	PHE	-	expression tag	UNP Q9UL62
D	-2	PRO	-	expression tag	UNP Q9UL62
D	-1	GLU	-	expression tag	UNP Q9UL62
D	0	PHE	-	expression tag	UNP Q9UL62

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
2	A	1	Total 1	Zn 1	0
2	B	1	Total 1	Zn 1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	AltConf
2	C	1	Total Zn 1 1	0
2	D	1	Total Zn 1 1	0

- Molecule 3 is PHOSPHATIDYLETHANOLAMINE (CCD ID: PTY) (formula: $C_{40}H_{80}NO_8P$).



Mol	Chain	Residues	Atoms	AltConf
3	A	1	Total C O P 40 31 8 1	0
3	B	1	Total C O P 40 31 8 1	0
3	C	1	Total C O P 40 31 8 1	0
3	D	1	Total C O P 40 31 8 1	0

- Molecule 4 is (2S)-3-(hexadecanoyloxy)-2-[(9Z)-octadec-9-enoyloxy]propyl 2-(trimethylammonio)ethyl phosphate (CCD ID: POV) (formula: $C_{42}H_{82}NO_8P$).



Mol	Chain	Residues	Atoms				AltConf
4	A	1	Total	C	O	P	0
			38	29	8	1	
4	B	1	Total	C	O	P	0
			38	29	8	1	
4	C	1	Total	C	O	P	0
			38	29	8	1	
4	D	1	Total	C	O	P	0
			38	29	8	1	

- Molecule 5 is CHOLESTEROL HEMISUCCINATE (CCD ID: Y01) (formula: $C_{31}H_{50}O_4$).



Mol	Chain	Residues	Atoms	AltConf
5	A	1	Total C O 35 31 4	0
5	B	1	Total C O 35 31 4	0
5	C	1	Total C O 35 31 4	0
5	D	1	Total C O 35 31 4	0

- Molecule 6 is (-)-englerin A (CCD ID: A1L55) (formula: $C_{26}H_{34}O_6$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	AltConf
6	A	1	Total C O 32 26 6	0
6	B	1	Total C O 32 26 6	0
6	C	1	Total C O 32 26 6	0
6	D	1	Total C O 32 26 6	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Short transient receptor potential channel 5

Chain A:



• Molecule 1: Short transient receptor potential channel 5

Chain B:



• Molecule 1: Short transient receptor potential channel 5

Chain C:





• Molecule 1: Short transient receptor potential channel 5

Chain D:




M648
D652
E653
G654
G655
T656
I663
I664
PRO
SER
PRO
LYS
SER
PHE
LEU
TYR
LEU
GLY
ASN
TRP
PHE
ASN
ASN
THR
PHE
CYS
PRO
LYS
ARG
ASP
PRO
ASP
GLY
ARG
ARG
ARG
ARG
ASN
LEU
ARG
SER
PHE
THR
GLU
ARG
ASN
ALA
ASP
S706
L707
I708
H712
L720
V721
M728
I729
R730
N731
S732
K733
THR
HIS
GLU
GLY
LEU
THR
GLU
GLU
ASN
PHE
LYS
GLU
LEU
LYS
GLN
ASP
ILE
SER
SER
PHE
ARG
TYR
GLU
VAL
LEU
ASP
LEU
LEU
LEU
GLY
ASN
ARG

4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	109774	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/Å^2$)	50	Depositor
Minimum defocus (nm)	1500	Depositor
Maximum defocus (nm)	1800	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	6.168	Depositor
Minimum map value	-3.776	Depositor
Average map value	-0.002	Depositor
Map value standard deviation	0.131	Depositor
Recommended contour level	1.06	Depositor
Map size (Å)	333.216, 333.216, 333.216	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.0413, 1.0413, 1.0413	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PTY, POV, ZN, Y01, A1L55

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.26	0/5529	0.40	0/7497
1	B	0.26	0/5529	0.40	0/7497
1	C	0.26	0/5529	0.40	0/7497
1	D	0.26	0/5529	0.40	0/7497
All	All	0.26	0/22116	0.40	0/29988

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5399	0	5453	104	0
1	B	5399	0	5453	107	0
1	C	5399	0	5453	107	0
1	D	5399	0	5453	108	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	40	0	56	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	40	0	56	2	0
3	C	40	0	56	2	0
3	D	40	0	56	2	0
4	A	38	0	47	3	0
4	B	38	0	47	3	0
4	C	38	0	47	4	0
4	D	38	0	47	3	0
5	A	35	0	49	4	0
5	B	35	0	49	6	0
5	C	35	0	49	5	0
5	D	35	0	49	3	0
6	A	32	0	0	0	0
6	B	32	0	0	0	0
6	C	32	0	0	0	0
6	D	32	0	0	0	0
All	All	22180	0	22420	380	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 380 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:519:LYS:HE3	1:D:506:LEU:HD13	1.66	0.76
1:A:506:LEU:HD13	1:D:519:LYS:HE3	1.66	0.76
1:B:519:LYS:HG2	1:C:506:LEU:HD22	1.68	0.75
1:A:519:LYS:HG2	1:B:506:LEU:HD22	1.68	0.75
1:B:519:LYS:HE3	1:C:506:LEU:HD13	1.66	0.75

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	652/767 (85%)	639 (98%)	13 (2%)	0	100	100
1	B	652/767 (85%)	639 (98%)	13 (2%)	0	100	100
1	C	652/767 (85%)	639 (98%)	13 (2%)	0	100	100
1	D	652/767 (85%)	639 (98%)	13 (2%)	0	100	100
All	All	2608/3068 (85%)	2556 (98%)	52 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	590/689 (86%)	590 (100%)	0	100	100
1	B	590/689 (86%)	590 (100%)	0	100	100
1	C	590/689 (86%)	590 (100%)	0	100	100
1	D	590/689 (86%)	590 (100%)	0	100	100
All	All	2360/2756 (86%)	2360 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 26 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	709	GLN
1	C	386	GLN
1	D	625	ASN
1	C	298	GLN
1	C	536	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 20 ligands modelled in this entry, 4 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	PTY	C	802	-	39,39,49	1.07	2 (5%)	42,44,54	1.23	5 (11%)
6	A1L55	B	805	-	34,35,35	1.62	5 (14%)	39,53,53	2.18	13 (33%)
5	Y01	B	804	-	38,38,38	1.22	4 (10%)	57,57,57	1.73	11 (19%)
3	PTY	B	802	-	39,39,49	1.07	2 (5%)	42,44,54	1.23	5 (11%)
5	Y01	D	804	-	38,38,38	1.22	4 (10%)	57,57,57	1.73	11 (19%)
5	Y01	C	804	-	38,38,38	1.22	4 (10%)	57,57,57	1.73	11 (19%)
6	A1L55	A	805	-	34,35,35	1.62	5 (14%)	39,53,53	2.18	13 (33%)
4	POV	B	803	-	37,37,51	1.09	2 (5%)	40,42,59	1.19	4 (10%)
6	A1L55	D	805	-	34,35,35	1.62	5 (14%)	39,53,53	2.18	13 (33%)
3	PTY	D	802	-	39,39,49	1.07	2 (5%)	42,44,54	1.23	5 (11%)
4	POV	A	803	-	37,37,51	1.09	2 (5%)	40,42,59	1.19	4 (10%)
4	POV	D	803	-	37,37,51	1.09	2 (5%)	40,42,59	1.19	4 (10%)
6	A1L55	C	805	-	34,35,35	1.62	5 (14%)	39,53,53	2.18	13 (33%)
3	PTY	A	802	-	39,39,49	1.07	2 (5%)	42,44,54	1.23	5 (11%)
4	POV	C	803	-	37,37,51	1.09	2 (5%)	40,42,59	1.19	4 (10%)
5	Y01	A	804	-	38,38,38	1.22	4 (10%)	57,57,57	1.73	11 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PTY	C	802	-	-	13/41/41/53	-
6	A1L55	B	805	-	-	3/21/67/67	0/5/4/4
5	Y01	B	804	-	-	14/19/77/77	0/4/4/4
3	PTY	B	802	-	-	13/41/41/53	-
5	Y01	D	804	-	-	14/19/77/77	0/4/4/4
5	Y01	C	804	-	-	14/19/77/77	0/4/4/4
6	A1L55	A	805	-	-	3/21/67/67	0/5/4/4
4	POV	B	803	-	-	5/39/39/55	-
6	A1L55	D	805	-	-	3/21/67/67	0/5/4/4
3	PTY	D	802	-	-	13/41/41/53	-
4	POV	A	803	-	-	5/39/39/55	-
4	POV	D	803	-	-	5/39/39/55	-
6	A1L55	C	805	-	-	3/21/67/67	0/5/4/4
3	PTY	A	802	-	-	13/41/41/53	-
4	POV	C	803	-	-	5/39/39/55	-
5	Y01	A	804	-	-	14/19/77/77	0/4/4/4

The worst 5 of 52 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	805	A1L55	C05-C04	4.30	1.66	1.52
6	B	805	A1L55	C05-C04	4.30	1.66	1.52
6	C	805	A1L55	C05-C04	4.30	1.66	1.52
6	D	805	A1L55	C05-C04	4.30	1.66	1.52
4	A	803	POV	O31-C31	4.25	1.45	1.33

The worst 5 of 132 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
6	A	805	A1L55	O03-C02-C25	6.03	124.51	111.39
6	B	805	A1L55	O03-C02-C25	6.03	124.51	111.39
6	C	805	A1L55	O03-C02-C25	6.03	124.51	111.39
6	D	805	A1L55	O03-C02-C25	6.03	124.51	111.39
6	A	805	A1L55	O03-C02-O01	-5.81	114.04	123.34

There are no chirality outliers.

5 of 140 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	802	PTY	C5-O14-P1-O13
3	B	802	PTY	C5-O14-P1-O13
3	C	802	PTY	C5-O14-P1-O13
3	D	802	PTY	C5-O14-P1-O13
5	A	804	Y01	CAV-CBC-OAW-CAY

There are no ring outliers.

12 monomers are involved in 39 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	802	PTY	2	0
5	B	804	Y01	6	0
3	B	802	PTY	2	0
5	D	804	Y01	3	0
5	C	804	Y01	5	0
4	B	803	POV	3	0
3	D	802	PTY	2	0
4	A	803	POV	3	0
4	D	803	POV	3	0
3	A	802	PTY	2	0
4	C	803	POV	4	0
5	A	804	Y01	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand PTY C 802



Bond lengths



Bond angles



Torsions



Rings

Ligand A1L55 B 805



Bond lengths



Bond angles



Torsions



Rings

Ligand Y01 B 804



Bond lengths



Bond angles



Torsions



Rings

Ligand PTY B 802



Bond lengths



Bond angles



Torsions



Rings

Ligand Y01 D 804



Bond lengths



Bond angles



Torsions



Rings

Ligand Y01 C 804



Bond lengths



Bond angles



Torsions



Rings

Ligand A1L55 A 805



Ligand POV B 803



Ligand A1L55 D 805



Ligand PTY D 802



Ligand POV A 803



Bond lengths



Bond angles



Torsions



Rings

Ligand POV D 803



Bond lengths



Bond angles



Torsions



Rings

Ligand A1L55 C 805



Bond lengths



Bond angles



Torsions



Rings

Ligand PTY A 802



Bond lengths



Bond angles



Torsions



Rings

Ligand POV C 803



Bond lengths



Bond angles



Torsions



Rings

Ligand Y01 A 804



Bond lengths



Bond angles



Torsions



Rings

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Map visualisation

This section contains visualisations of the EMDB entry EMD-63631. These allow visual inspection of the internal detail of the map and identification of artifacts.

Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections

6.1.1 Primary map



X



Y



Z

6.1.2 Raw map



X



Y



Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map



X Index: 160



Y Index: 160



Z Index: 160

6.2.2 Raw map



X Index: 160



Y Index: 160



Z Index: 160

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map



X Index: 178



Y Index: 178



Z Index: 185

6.3.2 Raw map



X Index: 143



Y Index: 143



Z Index: 186

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map



X Y Z

6.4.2 Raw map



X Y Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views (i)

This section was not generated.

6.6 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

7.2 Volume estimate (i)



The volume at the recommended contour level is 64 nm^3; this corresponds to an approximate mass of 58 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.382 $Å^{-1}$

8 Fourier-Shell correlation

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC



*Reported resolution corresponds to spatial frequency of 0.382 Å^{-1}

8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.62	-	-
Author-provided FSC curve	2.62	2.99	2.70
Unmasked-calculated*	3.13	3.55	3.16

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.13 differs from the reported value 2.62 by more than 10 %

9 Map-model fit

This section contains information regarding the fit between EMDB map EMD-63631 and PDB model 9M4W. Per-residue inclusion information can be found in section 3 on page 7.

9.1 Map-model overlay



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 1.06 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model



X Y Z

The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model



X Y Z

The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (1.06).

9.4 Atom inclusion (i)



At the recommended contour level, 65% of all backbone atoms, 59% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (1.06) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.5860	0.5800
A	0.5850	0.5800
B	0.5860	0.5810
C	0.5860	0.5800
D	0.5860	0.5800