



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 04:41 AM UTC

PDB ID : 4M5Y / pdb_00004m5y
Title : Crystal structure of broadly neutralizing Fab 5J8
Authors : Lee, P.S.; Wilson, I.A.
Deposited on : 2013-08-08
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2145 (1.56-1.56)
Clashscore	190562	2189 (1.56-1.56)
Ramachandran outliers	187476	2153 (1.56-1.56)
Sidechain outliers	187428	2150 (1.56-1.56)
RSRZ outliers	180081	2146 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	233	12% 83% 10% 7%
1	I	233	4% 88% 8% .
2	L	214	7% 92% 7% .
2	M	214	3% 94% . .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PEG	I	302	-	-	X	-
3	PEG	I	303	-	-	X	-
4	PG4	H	305	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7570 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Fab 5J8 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	217	Total	C	N	O	S	0	9	0
			1683	1079	270	330	4			
1	I	224	Total	C	N	O	S	0	10	0
			1727	1110	276	337	4			

- Molecule 2 is a protein called Fab 5J8 light chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	211	Total	C	N	O	S	0	4	0
			1599	1004	267	324	4			
2	M	211	Total	C	N	O	S	0	5	0
			1598	1004	264	326	4			

- Molecule 3 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	H	1	Total C O 7 4 3	0	0
3	H	1	Total C O 7 4 3	0	0
3	H	1	Total C O 7 4 3	0	0
3	H	1	Total C O 7 4 3	0	0
3	L	1	Total C O 7 4 3	0	0
3	I	1	Total C O 7 4 3	0	0
3	I	1	Total C O 7 4 3	0	0
3	I	1	Total C O 7 4 3	0	0
3	I	1	Total C O 7 4 3	0	0
3	I	1	Total C O 7 4 3	0	0
3	M	1	Total C O 7 4 3	0	0
3	M	1	Total C O 7 4 3	0	0

- Molecule 4 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	H	1	Total	C	O	0	0
			13	8	5		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	H	209	Total	O	0	0
			209	209		
5	L	194	Total	O	0	0
			194	194		
5	I	240	Total	O	0	0
			240	240		
5	M	223	Total	O	0	0
			223	223		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Fab 5J8 heavy chain




• Molecule 1: Fab 5J8 heavy chain



• Molecule 2: Fab 5J8 light chain



• Molecule 2: Fab 5J8 light chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	90.57Å 99.98Å 144.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.15 – 1.55 49.15 – 1.55	Depositor EDS
% Data completeness (in resolution range)	96.8 (49.15-1.55) 96.8 (49.15-1.55)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.94 (at 1.55Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, R_{free}	0.172 , 0.186 0.174 , 0.188	Depositor DCC
R_{free} test set	9216 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	20.1	Xtriage
Anisotropy	0.270	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 44.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7570	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 41.33 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.3987e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PG4, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	H	0.48	0/1751	0.74	0/2394
1	I	0.49	0/1805	0.74	0/2468
2	L	0.45	0/1650	0.73	0/2261
2	M	0.44	0/1652	0.74	1/2264 (0.0%)
All	All	0.47	0/6858	0.74	1/9387 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	M	155	VAL	CB-CA-C	-5.09	103.10	110.63

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1683	0	1661	24	0
1	I	1727	0	1726	19	0
2	L	1599	0	1564	9	0
2	M	1598	0	1563	9	0
3	H	28	0	40	9	0
3	I	35	0	50	15	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	L	7	0	10	0	0
3	M	14	0	20	2	0
4	H	13	0	18	8	0
5	H	209	0	0	6	0
5	I	240	0	0	4	0
5	L	194	0	0	0	0
5	M	223	0	0	0	0
All	All	7570	0	6652	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:81:LYS:HB3	4:H:305:PG4:H11	1.59	0.84
1:H:84:ALA:HB1	3:H:303:PEG:H32	1.63	0.80
1:I:97:ARG:HB2	3:I:302:PEG:H22	1.73	0.70
1:H:46:GLU:OE1	5:H:536:HOH:O	2.14	0.66
1:H:108:LEU:HB2	3:H:304:PEG:H32	1.78	0.66
3:I:302:PEG:O4	5:I:575:HOH:O	2.14	0.65
1:H:99:GLY:HA3	3:I:303:PEG:H31	1.80	0.63
2:M:132[A]:LEU:HD12	2:M:178:LEU:HD23	1.79	0.62
1:H:21[B]:THR:HG22	1:H:79[B]:SER:OG	2.00	0.62
1:I:70[B]:SER:OG	3:I:304:PEG:H32	2.00	0.62
1:I:97:ARG:HH11	3:I:302:PEG:H32	1.66	0.61
1:H:70[B]:SER:OG	4:H:305:PG4:O4	2.12	0.61
1:I:21[B]:THR:HG22	1:I:79[B]:SER:OG	2.01	0.61
2:L:14:ALA:HB3	2:L:17:GLU:HG3	1.87	0.57
1:I:21[B]:THR:HG23	5:I:596:HOH:O	2.06	0.56
4:H:305:PG4:O5	5:H:472:HOH:O	2.18	0.56
2:M:120:PRO:HD3	2:M:132[A]:LEU:CD2	2.38	0.54
1:H:70[A]:SER:OG	4:H:305:PG4:H82	2.07	0.54
1:H:81:LYS:CB	4:H:305:PG4:H11	2.35	0.53
2:L:167:GLN:O	2:L:168:SER:OG	2.24	0.53
1:I:97:ARG:NH1	3:I:302:PEG:H32	2.24	0.52
1:H:21[B]:THR:HG23	5:H:575:HOH:O	2.09	0.52
1:H:100(B):ASP:HB3	2:L:93[A]:ILE:HD11	1.93	0.51
1:H:43:LYS:NZ	3:H:302:PEG:H12	2.26	0.51
1:I:102:LYS:HB3	3:I:301:PEG:H11	1.92	0.50
1:I:171:GLN:NE2	5:I:592:HOH:O	2.40	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:27:TYR:HA	3:I:303:PEG:H32	1.94	0.49
1:H:99:GLY:C	3:I:303:PEG:H22	2.37	0.49
3:H:303:PEG:H42	5:H:481:HOH:O	2.13	0.49
1:H:15:SER:OG	3:H:301:PEG:H22	2.14	0.48
2:L:120:PRO:HD3	2:L:132:LEU:CD2	2.43	0.48
3:I:302:PEG:H41	5:I:582:HOH:O	2.14	0.48
1:I:26:GLY:O	3:I:303:PEG:H32	2.13	0.47
1:H:100(B):ASP:HB3	2:L:93[A]:ILE:CD1	2.44	0.47
1:I:102:LYS:HD2	3:I:301:PEG:H11	1.97	0.46
2:L:54[B]:ARG:HD3	2:L:62:PHE:O	2.16	0.45
2:M:33:LEU:HD12	2:M:90[A]:VAL:HG13	1.98	0.45
3:H:301:PEG:H42	5:H:573:HOH:O	2.17	0.45
1:I:100(B):ASP:HB3	2:M:93:ILE:HD11	1.98	0.45
2:M:33:LEU:HD22	2:M:71:ALA:CB	2.46	0.45
2:L:54[A]:ARG:HD3	2:L:62:PHE:O	2.17	0.45
1:H:40:PRO:HB3	3:H:302:PEG:H21	1.99	0.45
2:M:107:GLY:H	3:M:302:PEG:H12	1.82	0.45
1:I:36:TRP:CG	1:I:80[A]:LEU:HD12	2.53	0.44
1:H:100:TYR:CD2	3:I:303:PEG:H21	2.52	0.44
1:H:68:GLY:HA3	4:H:305:PG4:H62	2.00	0.43
1:I:59:TYR:HD2	1:I:64:GLU:HG3	1.81	0.43
3:H:302:PEG:H41	5:H:579:HOH:O	2.17	0.43
1:I:178:LEU:C	1:I:178:LEU:HD12	2.44	0.43
1:I:100(B):ASP:HB3	2:M:93:ILE:CD1	2.48	0.43
1:I:34:TYR:HE2	3:I:302:PEG:H31	1.84	0.43
1:I:103:TRP:HB2	3:I:301:PEG:H42	2.00	0.42
2:M:107:GLY:N	3:M:302:PEG:H12	2.35	0.42
1:H:68:GLY:C	4:H:305:PG4:H62	2.44	0.42
1:H:106:GLY:O	3:H:304:PEG:H11	2.19	0.41
2:L:33:LEU:HD12	2:L:90[A]:VAL:HG13	2.01	0.41
1:H:81:LYS:HB3	4:H:305:PG4:C1	2.41	0.41
2:M:120:PRO:HD3	2:M:132[A]:LEU:HD23	2.03	0.40
1:H:189:LEU:HD23	1:H:189:LEU:HA	1.91	0.40
2:L:13:VAL:HG11	2:L:19:ALA:HB2	2.03	0.40
1:H:178:LEU:C	1:H:178:LEU:HD12	2.47	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	222/233 (95%)	217 (98%)	5 (2%)	0	100	100
1	I	232/233 (100%)	228 (98%)	4 (2%)	0	100	100
2	L	213/214 (100%)	207 (97%)	6 (3%)	0	100	100
2	M	214/214 (100%)	210 (98%)	4 (2%)	0	100	100
All	All	881/894 (98%)	862 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	196/201 (98%)	191 (97%)	5 (3%)	40	13
1	I	202/201 (100%)	201 (100%)	1 (0%)	81	68
2	L	182/181 (101%)	181 (100%)	1 (0%)	81	68
2	M	183/181 (101%)	183 (100%)	0	100	100
All	All	763/764 (100%)	756 (99%)	7 (1%)	68	51

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	150	VAL
1	H	187	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	H	204	ASN
1	H	209	LYS
1	H	210	ARG
2	L	155	VAL
1	I	210	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
2	L	37	GLN
2	L	95(B)	GLN
2	L	126	GLN
1	I	53	HIS
1	I	171	GLN
2	M	126	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PEG	M	302	-	6,6,6	0.36	0	5,5,5	0.39	0
4	PG4	H	305	-	12,12,12	0.38	0	11,11,11	0.87	0
3	PEG	I	302	-	6,6,6	0.44	0	5,5,5	0.31	0
3	PEG	M	301	-	6,6,6	0.46	0	5,5,5	0.29	0
3	PEG	I	305	-	6,6,6	0.49	0	5,5,5	0.18	0
3	PEG	H	302	-	6,6,6	0.55	0	5,5,5	0.53	0
3	PEG	I	304	-	6,6,6	0.44	0	5,5,5	0.42	0
3	PEG	H	304	-	6,6,6	0.44	0	5,5,5	0.37	0
3	PEG	I	301	-	6,6,6	0.49	0	5,5,5	0.35	0
3	PEG	H	301	-	6,6,6	0.41	0	5,5,5	0.75	0
3	PEG	I	303	-	6,6,6	0.40	0	5,5,5	0.59	0
3	PEG	H	303	-	6,6,6	0.41	0	5,5,5	0.74	0
3	PEG	L	301	-	6,6,6	0.44	0	5,5,5	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEG	M	302	-	-	2/4/4/4	-
4	PG4	H	305	-	-	6/10/10/10	-
3	PEG	I	302	-	-	2/4/4/4	-
3	PEG	M	301	-	-	0/4/4/4	-
3	PEG	I	305	-	-	1/4/4/4	-
3	PEG	H	302	-	-	2/4/4/4	-
3	PEG	I	304	-	-	2/4/4/4	-
3	PEG	H	304	-	-	1/4/4/4	-
3	PEG	I	301	-	-	2/4/4/4	-
3	PEG	H	301	-	-	1/4/4/4	-
3	PEG	I	303	-	-	2/4/4/4	-
3	PEG	H	303	-	-	4/4/4/4	-
3	PEG	L	301	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (28) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	H	305	PG4	O2-C3-C4-O3
3	I	304	PEG	O1-C1-C2-O2
3	I	301	PEG	O1-C1-C2-O2
4	H	305	PG4	O1-C1-C2-O2
3	I	305	PEG	O2-C3-C4-O4
3	H	303	PEG	O2-C3-C4-O4
3	M	302	PEG	O1-C1-C2-O2
3	I	303	PEG	O1-C1-C2-O2
3	M	302	PEG	C4-C3-O2-C2
4	H	305	PG4	C3-C4-O3-C5
3	H	303	PEG	O1-C1-C2-O2
3	I	301	PEG	O2-C3-C4-O4
3	H	304	PEG	C4-C3-O2-C2
3	L	301	PEG	C1-C2-O2-C3
3	L	301	PEG	C4-C3-O2-C2
3	H	302	PEG	O1-C1-C2-O2
3	L	301	PEG	O2-C3-C4-O4
3	I	303	PEG	O2-C3-C4-O4
3	I	304	PEG	C4-C3-O2-C2
3	H	303	PEG	C1-C2-O2-C3
3	I	302	PEG	O2-C3-C4-O4
3	I	302	PEG	C1-C2-O2-C3
3	H	301	PEG	O1-C1-C2-O2
4	H	305	PG4	C8-C7-O4-C6
4	H	305	PG4	C5-C6-O4-C7
3	H	303	PEG	C4-C3-O2-C2
4	H	305	PG4	O3-C5-C6-O4
3	H	302	PEG	O2-C3-C4-O4

There are no ring outliers.

10 monomers are involved in 34 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	M	302	PEG	2	0
4	H	305	PG4	8	0
3	I	302	PEG	6	0
3	H	302	PEG	3	0
3	I	304	PEG	1	0
3	H	304	PEG	2	0
3	I	301	PEG	3	0
3	H	301	PEG	2	0
3	I	303	PEG	5	0
3	H	303	PEG	2	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	H	217/233 (93%)	0.34	28 (12%) 7 9	7, 20, 63, 78	9 (4%)
1	I	224/233 (96%)	-0.05	9 (4%) 42 50	9, 19, 48, 61	10 (4%)
2	L	211/214 (98%)	0.35	16 (7%) 20 23	10, 27, 47, 61	4 (1%)
2	M	211/214 (98%)	0.19	6 (2%) 55 63	10, 26, 42, 57	5 (2%)
All	All	863/894 (96%)	0.20	59 (6%) 23 27	7, 23, 51, 78	28 (3%)

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	M	209	THR	7.3
2	L	209	THR	6.6
1	H	135	THR	5.8
1	H	213	PRO	5.0
1	I	130	SER	4.5
2	L	168	SER	4.1
1	H	187	SER	4.1
1	H	1	GLU	4.0
1	H	127	SER	4.0
1	H	186	SER	3.9
1	H	185	PRO	3.7
1	H	158	ALA	3.7
2	L	40	PRO	3.6
2	M	156	LYS	3.6
1	H	189	LEU	3.6
2	L	155	VAL	3.6
2	M	155	VAL	3.6
1	H	136	ALA	3.5
1	H	206	LYS	3.5
1	H	184	VAL	3.3
1	H	188	SER	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	I	131	THR	3.2
1	H	193	THR	3.1
1	I	129	LYS	2.9
1	I	132	SER	2.9
1	H	195	ILE	2.8
2	L	41	GLY	2.8
1	I	128	SER	2.7
2	L	166	LYS	2.7
2	L	156	LYS	2.6
1	I	1	GLU	2.6
1	H	161	SER	2.6
1	H	140	CYS	2.6
1	H	162	GLY	2.6
2	L	1	SER	2.5
1	H	191	THR	2.5
2	L	170	ASN	2.5
1	H	137	ALA	2.4
1	I	158	ALA	2.4
1	H	204	ASN	2.4
2	L	107	GLY	2.4
2	L	143	ALA	2.4
1	H	194	TYR	2.3
1	H	192	GLN	2.3
1	H	163	VAL	2.2
1	H	210	ARG	2.2
2	L	172	TYR	2.2
1	I	213	PRO	2.2
2	L	109	PRO	2.1
1	H	159	LEU	2.1
2	L	169	ASN	2.1
1	H	126	PRO	2.1
2	M	40	PRO	2.1
2	M	1	SER	2.0
2	M	56	SER	2.0
1	I	204	ASN	2.0
1	H	154	TRP	2.0
2	L	106(A)	LEU	2.0
2	L	56	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	PEG	H	304	7/7	0.69	0.21	49,53,56,60	0
3	PEG	M	301	7/7	0.72	0.22	46,50,55,55	0
3	PEG	H	301	7/7	0.80	0.18	42,46,53,60	0
3	PEG	L	301	7/7	0.80	0.19	35,49,58,59	0
3	PEG	H	302	7/7	0.80	0.17	37,46,52,58	0
3	PEG	I	302	7/7	0.81	0.21	40,50,59,65	0
4	PG4	H	305	13/13	0.81	0.19	25,49,61,61	0
3	PEG	I	304	7/7	0.83	0.16	30,40,52,55	0
3	PEG	I	305	7/7	0.83	0.18	44,53,60,60	0
3	PEG	H	303	7/7	0.85	0.19	25,48,58,59	0
3	PEG	M	302	7/7	0.86	0.16	29,49,58,63	0
3	PEG	I	301	7/7	0.89	0.13	31,37,43,43	0
3	PEG	I	303	7/7	0.93	0.12	21,30,39,40	0

6.5 Other polymers [i](#)

There are no such residues in this entry.