



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:57 PM UTC

PDB ID : 6M53 / pdb_00006m53
Title : Crystal structure of 2, 3-dihydroxybenzoic acid decarboxylase from *Fusarium oxysporum*
Authors : Song, M.K.; Feng, J.H.; Liu, W.D.; Wu, Q.Q.; Zhu, D.M.
Deposited on : 2020-03-09
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2145 (1.56-1.56)
Clashscore	190562	2189 (1.56-1.56)
Ramachandran outliers	187476	2153 (1.56-1.56)
Sidechain outliers	187428	2150 (1.56-1.56)
RSRZ outliers	180081	2146 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	343	0% 93% 7% 0%
1	B	343	3% 92% 7% 0%
1	C	343	4% 89% 7% 0%
1	D	343	19% 90% 7% 0%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 12387 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2,3-dihydroxybenzoate decarboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	334	Total	C	N	O	S	0	0	0
			2692	1737	460	486	9			
1	B	339	Total	C	N	O	S	0	0	0
			2739	1764	471	495	9			
1	C	328	Total	C	N	O	S	0	0	0
			2638	1701	452	476	9			
1	D	332	Total	C	N	O	S	0	0	0
			2677	1729	457	482	9			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	HIS	-	expression tag	UNP A0A420U2F4
A	338	HIS	-	expression tag	UNP A0A420U2F4
A	339	HIS	-	expression tag	UNP A0A420U2F4
A	340	HIS	-	expression tag	UNP A0A420U2F4
A	341	HIS	-	expression tag	UNP A0A420U2F4
A	342	HIS	-	expression tag	UNP A0A420U2F4
A	343	HIS	-	expression tag	UNP A0A420U2F4
B	1	HIS	-	expression tag	UNP A0A420U2F4
B	338	HIS	-	expression tag	UNP A0A420U2F4
B	339	HIS	-	expression tag	UNP A0A420U2F4
B	340	HIS	-	expression tag	UNP A0A420U2F4
B	341	HIS	-	expression tag	UNP A0A420U2F4
B	342	HIS	-	expression tag	UNP A0A420U2F4
B	343	HIS	-	expression tag	UNP A0A420U2F4
C	1	HIS	-	expression tag	UNP A0A420U2F4
C	338	HIS	-	expression tag	UNP A0A420U2F4
C	339	HIS	-	expression tag	UNP A0A420U2F4
C	340	HIS	-	expression tag	UNP A0A420U2F4
C	341	HIS	-	expression tag	UNP A0A420U2F4
C	342	HIS	-	expression tag	UNP A0A420U2F4
C	343	HIS	-	expression tag	UNP A0A420U2F4

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	1	HIS	-	expression tag	UNP A0A420U2F4
D	338	HIS	-	expression tag	UNP A0A420U2F4
D	339	HIS	-	expression tag	UNP A0A420U2F4
D	340	HIS	-	expression tag	UNP A0A420U2F4
D	341	HIS	-	expression tag	UNP A0A420U2F4
D	342	HIS	-	expression tag	UNP A0A420U2F4
D	343	HIS	-	expression tag	UNP A0A420U2F4

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0
2	B	1	Total Zn 1 1	0	0
2	C	1	Total Zn 1 1	0	0
2	D	1	Total Zn 1 1	0	0

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total C O 6 3 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	450	Total	O	0	0
			450	450		
4	B	477	Total	O	0	0
			477	477		
4	C	412	Total	O	0	0
			412	412		
4	D	280	Total	O	0	0
			280	280		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 2,3-dihydroxybenzoate decarboxylase

Chain A:  93% . . .





- Molecule 1: 2,3-dihydroxybenzoate decarboxylase

Chain B:  92% 7% .



- Molecule 1: 2,3-dihydroxybenzoate decarboxylase


Chain C:  89% 7% .



- Molecule 1: 2,3-dihydroxybenzoate decarboxylase

Chain D:  90% 7% .





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	77.62Å 132.25Å 140.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.77 – 1.55 24.77 – 1.55	Depositor EDS
% Data completeness (in resolution range)	99.0 (24.77-1.55) 98.9 (24.77-1.55)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.73 (at 1.55Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.186 , 0.199 0.190 , 0.202	Depositor DCC
R_{free} test set	10355 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	20.3	Xtriage
Anisotropy	0.180	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12387	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.33	0/2773	0.59	1/3767 (0.0%)
1	B	0.36	0/2823	0.62	3/3835 (0.1%)
1	C	0.34	0/2716	0.56	1/3689 (0.0%)
1	D	0.32	0/2758	0.55	1/3747 (0.0%)
All	All	0.33	0/11070	0.58	6/15038 (0.0%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	190	PRO	N-CA-C	6.15	118.21	110.70
1	B	291	ILE	O-C-N	6.14	127.45	121.98
1	B	190	PRO	N-CA-C	5.60	117.53	110.70
1	C	190	PRO	N-CA-C	5.44	117.34	110.70
1	B	291	ILE	CA-C-O	-5.33	115.61	120.73
1	D	190	PRO	N-CA-C	5.22	117.07	110.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2692	0	2611	8	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2739	0	2642	14	0
1	C	2638	0	2565	13	0
1	D	2677	0	2598	18	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	B	12	0	15	2	0
3	C	6	0	7	0	0
4	A	450	0	0	3	0
4	B	477	0	0	3	0
4	C	412	0	0	4	0
4	D	280	0	0	2	0
All	All	12387	0	10438	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:2:MET:HE1	1:D:316:ARG:HA	1.48	0.94
1:A:92:LYS:HE3	4:A:816:HOH:O	1.98	0.62
1:B:267:SER:HB2	1:B:292:ASP:HB2	1.84	0.60
1:D:2:MET:HE1	1:D:316:ARG:CA	2.29	0.58
1:B:52:GLU:HG2	1:B:340:HIS:NE2	2.24	0.53
1:D:74:LYS:H	1:D:74:LYS:HE2	1.73	0.53
1:B:170:ARG:HG3	4:B:713:HOH:O	2.08	0.53
1:B:216:LYS:NZ	4:B:507:HOH:O	2.41	0.53
1:B:324:LYS:HA	1:B:329:LEU:HG	1.92	0.52
1:C:263:TRP:HH2	1:C:325:LYS:HZ1	1.58	0.51
1:C:277:PHE:CZ	1:C:281:GLU:HG3	2.45	0.50
1:D:332:PHE:C	1:D:332:PHE:CD1	2.89	0.50
1:B:196:GLN:NE2	3:B:401:GOL:O2	2.38	0.50
1:D:170:ARG:HD2	4:D:612:HOH:O	2.11	0.49
1:C:179:LYS:HG3	4:C:512:HOH:O	2.12	0.48
1:D:252:LEU:HB2	1:D:257:TYR:CZ	2.49	0.48
1:A:74:LYS:NZ	4:A:506:HOH:O	2.46	0.48
1:A:9:GLU:O	1:A:292:ASP:HA	2.14	0.47
1:D:9:GLU:O	1:D:292:ASP:HA	2.14	0.47
1:D:189:GLY:HA2	1:D:194:PHE:CE1	2.51	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:183:LYS:HB2	1:C:183:LYS:HE2	1.76	0.46
1:C:9:GLU:O	1:C:292:ASP:HA	2.17	0.45
1:D:216:LYS:N	1:D:216:LYS:HD3	2.32	0.45
1:C:87:ILE:O	1:C:91:ILE:HG12	2.16	0.45
1:C:65:PRO:HB2	1:C:69:ASP:HB2	1.99	0.44
1:B:276:GLN:HG2	4:B:959:HOH:O	2.17	0.43
1:B:179:LYS:NZ	1:D:141:ASP:OD2	2.47	0.43
1:C:265:THR:HA	1:C:288:LEU:O	2.17	0.43
1:D:123:TYR:O	4:D:501:HOH:O	2.21	0.43
1:D:134:GLN:HB3	1:D:144:ILE:HB	1.99	0.43
1:D:9:GLU:OE1	1:D:166:TYR:OH	2.36	0.43
1:B:313:VAL:HG12	1:B:317:LYS:HE3	2.00	0.42
1:B:334:GLN:HG2	3:B:402:GOL:O2	2.19	0.42
1:D:324:LYS:HG2	1:D:332:PHE:HE2	1.85	0.42
1:C:316:ARG:HD3	4:C:827:HOH:O	2.18	0.42
1:C:322:ASN:O	1:C:325:LYS:HG3	2.20	0.42
1:D:165:PHE:HB3	1:D:219:ILE:HD13	2.02	0.42
1:D:46:ARG:O	1:D:50:MET:HG3	2.19	0.42
1:A:87:ILE:O	1:A:91:ILE:HG12	2.20	0.41
1:B:238:TRP:CE2	1:D:190:PRO:HG2	2.55	0.41
1:C:94:HIS:HD2	4:C:837:HOH:O	2.03	0.41
1:A:110:LYS:O	1:A:114:GLU:HG3	2.21	0.41
1:C:251:LYS:HE3	4:C:790:HOH:O	2.20	0.41
1:A:126:LYS:NZ	4:A:507:HOH:O	2.48	0.41
1:B:242:ILE:O	1:B:246:LEU:HG	2.21	0.41
1:A:167:LEU:HD21	1:A:201:HIS:HB3	2.03	0.41
1:B:238:TRP:CZ2	1:D:190:PRO:HG2	2.56	0.40
1:A:324:LYS:HG2	1:A:332:PHE:HE2	1.86	0.40
1:C:167:LEU:HD21	1:C:201:HIS:HB3	2.04	0.40
1:B:65:PRO:HB2	1:B:69:ASP:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	332/343 (97%)	320 (96%)	11 (3%)	1 (0%)	36	18
1	B	337/343 (98%)	326 (97%)	10 (3%)	1 (0%)	36	18
1	C	326/343 (95%)	315 (97%)	10 (3%)	1 (0%)	36	18
1	D	330/343 (96%)	315 (96%)	14 (4%)	1 (0%)	36	18
All	All	1325/1372 (97%)	1276 (96%)	45 (3%)	4 (0%)	36	18

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	192	LEU
1	B	192	LEU
1	C	192	LEU
1	D	192	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	281/290 (97%)	279 (99%)	2 (1%)	76	59
1	B	286/290 (99%)	285 (100%)	1 (0%)	86	77
1	C	275/290 (95%)	274 (100%)	1 (0%)	84	73
1	D	279/290 (96%)	278 (100%)	1 (0%)	84	73
All	All	1121/1160 (97%)	1116 (100%)	5 (0%)	84	73

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	92	LYS
1	A	110	LYS
1	B	325	LYS
1	C	139	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	333	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	236	ASN
1	A	261	ASN
1	B	236	ASN
1	B	261	ASN
1	C	45	GLN
1	C	236	ASN
1	D	17	HIS
1	D	45	GLN
1	D	236	ASN
1	D	261	ASN
1	D	331	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 4 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	B	401	-	5,5,5	1.44	1 (20%)	5,5,5	1.02	0
3	GOL	B	402	-	5,5,5	1.11	0	5,5,5	1.19	1 (20%)
3	GOL	C	401	-	5,5,5	1.79	2 (40%)	5,5,5	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	401	-	-	2/4/4/4	-
3	GOL	B	402	-	-	0/4/4/4	-
3	GOL	C	401	-	-	0/4/4/4	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	401	GOL	O2-C2	-2.78	1.35	1.43
3	B	401	GOL	O2-C2	-2.40	1.36	1.43
3	C	401	GOL	C1-C2	2.24	1.60	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	402	GOL	C3-C2-C1	-2.13	103.99	111.80

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	401	GOL	O1-C1-C2-C3
3	B	401	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	401	GOL	1	0
3	B	402	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	334/343 (97%)	-0.06	4 (1%) 76 83	13, 19, 30, 54	0
1	B	339/343 (98%)	0.26	12 (3%) 47 55	14, 22, 35, 47	0
1	C	328/343 (95%)	0.48	14 (4%) 40 48	14, 24, 36, 47	0
1	D	332/343 (96%)	1.18	65 (19%) 3 3	16, 31, 46, 63	0
All	All	1333/1372 (97%)	0.46	95 (7%) 22 25	13, 23, 40, 63	0

All (95) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	333	TYR	6.1
1	C	24	TRP	5.3
1	B	27	LEU	5.2
1	C	138	ALA	5.2
1	D	23	TRP	4.6
1	D	138	ALA	4.3
1	B	24	TRP	4.2
1	B	340	HIS	4.1
1	D	332	PHE	4.1
1	D	13	ALA	4.0
1	D	80	ALA	4.0
1	D	21	THR	3.8
1	D	120	VAL	3.7
1	C	27	LEU	3.6
1	D	148	GLY	3.6
1	D	119	VAL	3.5
1	C	28	PHE	3.5
1	A	331	GLN	3.5
1	B	138	ALA	3.4
1	B	28	PHE	3.3
1	D	76	ALA	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	121	THR	3.2
1	C	329	LEU	3.2
1	D	14	LEU	3.1
1	A	330	PRO	3.1
1	D	331	GLN	3.1
1	A	333	TYR	3.0
1	B	247	GLY	3.0
1	C	23	TRP	3.0
1	D	87	ILE	3.0
1	D	330	PRO	3.0
1	D	153	VAL	3.0
1	C	119	VAL	2.9
1	C	26	GLY	2.9
1	D	81	VAL	2.9
1	D	24	TRP	2.9
1	D	114	GLU	2.9
1	D	47	ILE	2.9
1	D	83	VAL	2.8
1	B	136	ALA	2.8
1	D	22	ARG	2.8
1	D	12	PHE	2.7
1	C	325	LYS	2.7
1	D	113	ALA	2.7
1	D	247	GLY	2.7
1	B	246	LEU	2.7
1	D	79	LEU	2.7
1	D	70	VAL	2.7
1	D	90	ALA	2.6
1	D	112	ALA	2.6
1	D	161	LEU	2.6
1	D	109	PRO	2.6
1	D	139	ASP	2.6
1	D	25	ALA	2.5
1	D	93	ALA	2.5
1	D	72	ASP	2.5
1	D	71	TRP	2.5
1	D	140	GLY	2.5
1	D	249	SER	2.5
1	D	37	ALA	2.4
1	D	91	ILE	2.4
1	D	157	THR	2.4
1	D	124	GLY	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	176	ILE	2.4
1	D	123	TYR	2.4
1	D	125	PHE	2.4
1	B	26	GLY	2.4
1	D	64	ALA	2.4
1	D	154	PHE	2.4
1	D	163	VAL	2.3
1	B	174	GLY	2.3
1	D	78	ALA	2.3
1	D	107	HIS	2.3
1	D	329	LEU	2.3
1	D	136	ALA	2.2
1	D	246	LEU	2.2
1	D	248	LEU	2.2
1	D	42	ILE	2.2
1	D	98	LEU	2.2
1	D	44	GLU	2.2
1	D	86	TYR	2.2
1	D	149	PRO	2.2
1	C	110	LYS	2.2
1	A	335	SER	2.1
1	B	186	TRP	2.1
1	D	67	VAL	2.1
1	D	28	PHE	2.1
1	D	295	PHE	2.1
1	C	25	ALA	2.1
1	D	11	ALA	2.1
1	D	55	VAL	2.0
1	C	30	ILE	2.0
1	C	139	ASP	2.0
1	C	312	ASP	2.0
1	D	49	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	B	401	6/6	0.89	0.10	26,29,29,33	0
3	GOL	B	402	6/6	0.89	0.09	24,24,30,32	0
2	ZN	A	401	1/1	0.90	0.53	30,30,30,30	0
2	ZN	B	403	1/1	0.90	0.43	30,30,30,30	0
2	ZN	C	402	1/1	0.94	0.41	30,30,30,30	0
3	GOL	C	401	6/6	0.96	0.10	16,19,22,24	0
2	ZN	D	401	1/1	0.97	0.52	30,30,30,30	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ZN A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN B 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN C 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ZN D 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.