



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 03:38 AM UTC

PDB ID : 6M5I / pdb_00006m5i
Title : Crystal structure of 2019-nCoV nsp7-nsp8c complex
Authors : Yan, L.M.; Ge, J.; Zhao, Y.; Lou, Z.Y.; Rao, Z.H.
Deposited on : 2020-03-10
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	198	
2	A	82	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1547 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Non-structural protein 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	115	Total	C	N	O	S	0	0	0
			891	562	149	173	7			

- Molecule 2 is a protein called Non-structural protein 7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	81	Total	C	N	O	S	0	0	0
			623	389	104	123	7			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	21	Total	O	0	0
			21	21		
3	A	12	Total	O	0	0
			12	12		

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	50.94Å 69.38Å 101.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.06 – 2.50 41.06 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.8 (41.06-2.50) 99.8 (41.06-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.17 (at 2.51Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.208 , 0.292 0.210 , 0.292	Depositor DCC
R_{free} test set	654 reflections (9.99%)	wwPDB-VP
Wilson B-factor (Å ²)	71.9	Xtriage
Anisotropy	0.402	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 55.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	1547	wwPDB-VP
Average B, all atoms (Å ²)	81.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.61	0/904	0.90	0/1233
2	A	0.64	0/626	0.92	2/843 (0.2%)
All	All	0.62	0/1530	0.90	2/2076 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2	LYS	N-CA-C	-6.29	105.61	113.28
2	A	40	LEU	N-CA-C	-5.08	105.93	111.82

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	891	0	906	33	3
2	A	623	0	654	22	3
3	A	12	0	0	2	0
3	B	21	0	0	4	0
All	All	1547	0	1560	52	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:119:CYS:SG	3:B:321:HOH:O	2.30	0.90
1:B:95:MET:SD	3:B:320:HOH:O	2.45	0.75
2:A:62:MET:O	3:A:101:HOH:O	2.06	0.73
1:B:173:GLN:OE1	3:B:301:HOH:O	2.08	0.72
1:B:192:THR:OG1	3:B:302:HOH:O	2.09	0.69
2:A:25:SER:O	2:A:27:LYS:N	2.28	0.67
2:A:76:LEU:O	2:A:79:ARG:NH1	2.30	0.65
2:A:18:GLN:NE2	2:A:23:GLU:OE2	2.30	0.65
1:B:111:ILE:HD13	2:A:53:VAL:HG23	1.82	0.61
2:A:50:GLU:O	2:A:53:VAL:HG12	2.01	0.60
1:B:89:THR:HG22	1:B:93:GLN:OE1	2.02	0.60
1:B:83:ASP:OD2	1:B:83:ASP:N	2.36	0.57
2:A:3:MET:HB2	2:A:45:THR:HG22	1.85	0.57
1:B:91:ALA:O	1:B:95:MET:HE2	2.05	0.57
2:A:3:MET:HE1	2:A:44:ASP:N	2.21	0.56
1:B:178:SER:OG	1:B:179:MET:O	2.19	0.56
2:A:13:LEU:HD23	2:A:55:LEU:HD23	1.88	0.56
2:A:1:SER:OG	2:A:2:LYS:N	2.40	0.55
2:A:2:LYS:HG2	2:A:5:ASP:HB2	1.89	0.54
1:B:165:VAL:HG12	1:B:171:ILE:HD13	1.89	0.54
1:B:166:ASP:HB2	1:B:186:ALA:HB3	1.89	0.54
1:B:182:SER:HA	1:B:185:LEU:HD12	1.89	0.54
2:A:3:MET:SD	2:A:45:THR:HA	2.50	0.51
1:B:139:ASP:HB2	1:B:142:THR:OG1	2.10	0.51
1:B:97:PHE:HA	1:B:100:LEU:HB2	1.93	0.50
1:B:179:MET:O	1:B:180:ASP:HB2	2.11	0.50
1:B:108:LEU:O	1:B:111:ILE:HG12	2.12	0.50
2:A:25:SER:C	2:A:27:LYS:H	2.18	0.49
2:A:54:SER:O	2:A:57:SER:OG	2.27	0.49
1:B:87:LYS:HA	1:B:90:SER:HB3	1.94	0.49
1:B:96:LEU:HD22	2:A:16:VAL:HG21	1.94	0.48
2:A:12:VAL:O	2:A:16:VAL:HG23	2.14	0.48
1:B:83:ASP:O	1:B:87:LYS:HG3	2.13	0.47
1:B:97:PHE:HA	1:B:100:LEU:HD12	1.95	0.47
2:A:21:ARG:NH1	3:A:102:HOH:O	2.39	0.47
1:B:144:LYS:HD3	1:B:144:LYS:HA	1.48	0.46
2:A:71:LEU:O	2:A:75:MET:HG2	2.15	0.46
1:B:174:LEU:HA	1:B:177:ILE:HD12	1.97	0.45
1:B:137:ILE:HD13	1:B:191:VAL:HG23	1.98	0.45
1:B:187:TRP:CE3	1:B:189:LEU:HG	2.52	0.44
1:B:90:SER:O	1:B:94:THR:OG1	2.27	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:85:ARG:NH2	1:B:88:VAL:HG21	2.34	0.43
2:A:43:LYS:HD2	2:A:43:LYS:HA	1.71	0.43
1:B:125:ILE:HG13	2:A:57:SER:CB	2.48	0.43
1:B:158:LEU:HD23	1:B:158:LEU:HA	1.87	0.42
1:B:132:LYS:NZ	1:B:134:MET:SD	2.90	0.42
2:A:59:LEU:HD11	2:A:66:VAL:HG21	2.01	0.42
1:B:152:PHE:HB3	1:B:159:TRP:HB2	2.02	0.41
1:B:164:VAL:HG22	1:B:191:VAL:HG22	2.02	0.41
2:A:35:LEU:O	2:A:39:ILE:HG23	2.20	0.41
1:B:159:TRP:HA	1:B:194:LEU:O	2.20	0.40
1:B:194:LEU:HD23	1:B:194:LEU:HA	1.82	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:85:ARG:NH2	2:A:37:ASN:CG[4_545]	1.91	0.29
1:B:85:ARG:NH2	2:A:37:ASN:OD1[4_545]	1.95	0.25
1:B:85:ARG:NH2	2:A:37:ASN:ND2[4_545]	1.97	0.23

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	113/198 (57%)	101 (89%)	7 (6%)	5 (4%)	2	2
2	A	79/82 (96%)	67 (85%)	9 (11%)	3 (4%)	2	3
All	All	192/280 (69%)	168 (88%)	16 (8%)	8 (4%)	2	2

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	26	SER
1	B	104	ASP
1	B	128	THR
1	B	174	LEU
1	B	177	ILE
2	A	65	ALA
2	A	80	ALA
1	B	161	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	101/167 (60%)	101 (100%)	0	100	100
2	A	75/76 (99%)	74 (99%)	1 (1%)	61	82
All	All	176/243 (72%)	175 (99%)	1 (1%)	78	91

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	39	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	115/198 (58%)	-0.16	0 100 100	65, 79, 98, 112	0
2	A	81/82 (98%)	-0.10	0 100 100	62, 80, 109, 119	0
All	All	196/280 (70%)	-0.14	0 100 100	62, 79, 105, 119	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.