



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 15, 2026 – 01:29 AM UTC

PDB ID : 9M5O / pdb_00009m5o
Title : Plasmodium vivax aspartyl-tRNA synthetase in complex with aspartyl sulfamoyl adenosine (Asp-AMS) Complex
Authors : Sharma, V.K.; Manickam, Y.; Sharma, A.; Bagale, S.; Pradeepkumar, P.I.
Deposited on : 2025-03-06
Resolution : 1.84 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1296 (1.84-1.84)
Clashscore	190562	1329 (1.84-1.84)
Ramachandran outliers	187476	1318 (1.84-1.84)
Sidechain outliers	187428	1318 (1.84-1.84)
RSRZ outliers	180081	1296 (1.84-1.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	536	5% 82% 11% 6%
1	B	536	3% 81% 11% 7%

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 9167 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called aspartate-tRNA ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	502	Total	C	N	O	S	0	10	0
			4118	2640	683	766	29			
1	B	499	Total	C	N	O	S	0	3	0
			4053	2602	675	749	27			

- Molecule 2 is 5'-O-(L-alpha-aspartylsulfamoyl)adenosine (CCD ID: DSZ) (formula: C₁₄H₁₉N₇O₉S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			31	14	7	9	1		
2	B	1	Total	C	N	O	S	0	0
			31	14	7	9	1		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total 2	Mg 2	0	0
3	B	2	Total 2	Mg 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	429	Total 452	O 452	0	23
4	B	459	Total 478	O 478	0	20


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: aspartate-tRNA ligase



- Molecule 1: aspartate-tRNA ligase



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	140.25Å 140.25Å 267.61Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.20 – 1.84 50.20 – 1.84	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.20-1.84) 99.9 (50.20-1.84)	Depositor EDS
R_{merge}	0.22	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.46 (at 1.84Å)	Xtriage
Refinement program	PHENIX (1.15rc1_3423: ???)	Depositor
R, R_{free}	0.168 , 0.196 0.169 , 0.195	Depositor DCC
R_{free} test set	6603 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	30.0	Xtriage
Anisotropy	0.112	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 48.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9167	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, DSZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.77	24/4237 (0.6%)	0.68	6/5713 (0.1%)
1	B	0.78	22/4151 (0.5%)	0.68	5/5599 (0.1%)
All	All	0.77	46/8388 (0.5%)	0.68	11/11312 (0.1%)

All (46) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	195	ARG	C-O	-8.77	1.14	1.24
1	B	276	GLU	C-O	-7.79	1.14	1.23
1	A	573	LYS	C-O	-7.63	1.15	1.24
1	A	303	LEU	C-O	-7.48	1.15	1.24
1	A	580	ASN	C-O	-7.38	1.15	1.24
1	A	304	ASN	C-O	-7.35	1.14	1.24
1	A	419	LYS	C-O	-7.27	1.16	1.23
1	B	127	SER	C-O	-7.21	1.15	1.24
1	B	396	ARG	C-O	-7.20	1.16	1.24
1	B	274	ALA	C-O	-7.15	1.14	1.24
1	B	395	PHE	C-O	-7.14	1.16	1.24
1	A	396	ARG	C-O	-7.11	1.16	1.24
1	B	445	LYS	C-O	-7.11	1.16	1.24
1	A	579	ALA	C-O	-7.05	1.15	1.24
1	B	238	SER	C-O	-6.69	1.15	1.23
1	A	539	GLU	C-O	-6.64	1.16	1.24
1	A	307[A]	CYS	CA-C	6.64	1.62	1.52
1	A	307[B]	CYS	CA-C	6.64	1.62	1.52
1	B	235	PRO	C-O	-6.59	1.16	1.23
1	A	197	LYS	C-O	-6.51	1.14	1.23
1	A	421	ASP	C-O	-6.39	1.16	1.23
1	A	541	ALA	C-O	-6.10	1.16	1.24
1	B	126	ILE	C-O	-6.08	1.17	1.24
1	B	442	LYS	C-O	-6.06	1.15	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	296	LYS	C-O	-6.05	1.16	1.23
1	B	504	LYS	C-O	-5.97	1.17	1.24
1	A	306	ARG	C-O	-5.92	1.17	1.24
1	A	305	ASN	C-O	-5.85	1.16	1.24
1	A	183	SER	C-O	-5.80	1.16	1.24
1	A	537	GLU	C-O	-5.74	1.17	1.24
1	B	112	VAL	C-O	-5.68	1.17	1.24
1	A	198	GLY	C-O	-5.67	1.16	1.23
1	B	236	LEU	C-O	-5.63	1.16	1.23
1	B	537	GLU	C-O	-5.63	1.17	1.24
1	B	114	GLU	C-O	-5.55	1.16	1.23
1	B	419	LYS	C-O	-5.43	1.17	1.24
1	A	417	THR	C-O	-5.30	1.17	1.23
1	B	275	LYS	C-O	-5.30	1.17	1.23
1	A	482	GLN	C-O	-5.26	1.18	1.24
1	A	356	ASN	C-O	-5.25	1.16	1.23
1	A	196	SER	C-O	-5.20	1.17	1.24
1	B	130	LYS	C-O	-5.15	1.17	1.24
1	B	417	THR	C-O	-5.03	1.17	1.23
1	B	128	LYS	C-O	-5.03	1.17	1.24
1	A	181	LEU	C-O	-5.01	1.17	1.24
1	B	356	ASN	C-O	-5.01	1.17	1.23

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	307[A]	CYS	CA-C-O	8.68	129.99	119.49
1	A	307[B]	CYS	CA-C-O	8.68	129.99	119.49
1	A	182	GLN	N-CA-C	8.30	120.41	111.36
1	B	295	ILE	N-CA-C	7.77	119.47	109.30
1	B	295	ILE	CB-CA-C	-6.06	102.13	111.71
1	A	399	ASN	N-CA-CB	6.05	120.08	110.22
1	A	307[A]	CYS	N-CA-C	6.03	119.74	112.38
1	A	307[B]	CYS	N-CA-C	6.03	119.74	112.38
1	B	354	GLY	N-CA-C	5.63	121.19	113.99
1	B	397	ALA	CA-C-N	5.04	128.01	120.95
1	B	397	ALA	C-N-CA	5.04	128.01	120.95

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4118	0	4028	29	0
1	B	4053	0	3953	30	0
2	A	31	0	18	0	0
2	B	31	0	18	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	452	0	0	3	0
4	B	478	0	0	3	0
All	All	9167	0	8017	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:106:THR:O	1:B:110:GLU:HG3	2.04	0.57
1:A:591:TYR:HA	1:B:273:THR:HG23	1.86	0.57
1:A:460:GLU:OE2	1:A:463:GLN:NE2	2.43	0.51
1:B:330:LYS:HG2	1:B:389:PHE:CE1	2.46	0.51
1:A:116:ILE:HD11	1:A:132:ASN:HD22	1.75	0.51
1:A:330:LYS:HZ2	1:B:330:LYS:HD2	1.76	0.51
1:A:360:ILE:HD11	1:B:360:ILE:HD12	1.92	0.51
1:B:398:GLU:OE1	4:B:801:HOH:O	2.19	0.50
1:A:209:LEU:HD21	1:A:317:SER:OG	2.11	0.49
1:B:489:LYS:NZ	4:B:811:HOH:O	2.45	0.49
1:A:209:LEU:HD12	1:A:448:MET:HE3	1.93	0.49
1:B:480:LEU:HD22	1:B:502:MET:CE	2.44	0.48
1:B:465:LEU:HG	1:B:516:THR:HB	1.95	0.47
1:B:496:TYR:CE2	1:B:498:MET:HA	2.49	0.47
1:A:330:LYS:HZ2	1:B:330:LYS:CD	2.27	0.47
1:A:525:PRO:HG2	1:A:528:LEU:HD12	1.97	0.47
1:A:535:TYR:HA	1:A:543:SER:HA	1.96	0.47
1:A:149:MET:HE1	1:A:259:LYS:HG2	1.97	0.47
1:A:537:GLU:OE2	1:B:125:LYS:HE2	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:141:LEU:HB2	1:A:207:HIS:HB3	1.96	0.46
1:B:113:CYS:HB3	1:B:124:VAL:HG12	1.98	0.46
1:A:228[A]:MET:HG3	4:A:901:HOH:O	2.16	0.46
1:A:205:LEU:O	1:A:211:SER:HA	2.17	0.45
1:A:630:ILE:HD13	4:A:930:HOH:O	2.15	0.45
1:A:330:LYS:HG2	1:A:389:PHE:CE1	2.51	0.45
1:B:205:LEU:O	1:B:211:SER:HA	2.16	0.44
1:A:273:THR:HG23	1:B:591:TYR:HA	1.99	0.44
1:B:583:ASN:OD1	1:B:584:PHE:N	2.50	0.44
1:A:465:LEU:HG	1:A:516:THR:HB	2.00	0.44
1:A:116:ILE:CD1	1:A:132:ASN:HD22	2.31	0.44
1:B:523:ASN:HA	1:B:543:SER:O	2.18	0.44
1:B:126:ILE:HD13	1:B:126:ILE:HA	1.75	0.43
1:A:121:TYR:HA	1:A:267:ILE:O	2.19	0.43
1:B:604:GLU:HG2	1:B:620:SER:HB2	1.99	0.43
1:A:100:ASN:O	1:A:104:GLU:HG2	2.19	0.43
1:B:535:TYR:HA	1:B:543:SER:HA	2.01	0.42
1:B:141:LEU:HB2	1:B:207:HIS:HB3	2.01	0.42
1:B:145:TYR:OH	1:B:448:MET:HE1	2.19	0.42
1:A:630:ILE:HD11	1:B:348:GLY:HA3	2.02	0.42
1:A:358:PHE:HB3	1:B:360:ILE:HD11	2.02	0.42
1:B:211:SER:OG	1:B:256:THR:HA	2.20	0.42
1:B:412:LEU:HB2	1:B:601:ILE:HB	2.02	0.41
1:A:199:SER:HB3	1:A:218:ILE:HG22	2.00	0.41
1:B:480:LEU:HD13	1:B:502:MET:HE2	2.02	0.41
1:A:287:LYS:HB3	1:A:287:LYS:HE2	1.96	0.41
1:B:480:LEU:HD13	1:B:502:MET:CE	2.50	0.41
1:A:474:GLU:O	1:A:478:LYS:HG3	2.20	0.41
1:A:537:GLU:HG2	4:A:1116[B]:HOH:O	2.20	0.41
1:B:630:ILE:HD13	4:B:936:HOH:O	2.21	0.41
1:A:523:ASN:HA	1:A:543:SER:O	2.21	0.41
1:B:566:ASN:O	1:B:570:GLU:HG3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	506/536 (94%)	494 (98%)	12 (2%)	0	100	100
1	B	496/536 (92%)	485 (98%)	11 (2%)	0	100	100
All	All	1002/1072 (94%)	979 (98%)	23 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	455/485 (94%)	455 (100%)	0	100	100
1	B	442/485 (91%)	441 (100%)	1 (0%)	87	84
All	All	897/970 (92%)	896 (100%)	1 (0%)	88	85

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	125	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	132	ASN
1	A	487	HIS
1	B	132	ASN
1	B	182	GLN
1	B	207	HIS
1	B	213	GLN
1	B	298	ASN
1	B	356	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	483	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	DSZ	A	701	3	33,33,33	0.24	0	46,49,49	0.37	0
2	DSZ	B	701	3	33,33,33	0.20	0	46,49,49	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DSZ	A	701	3	-	4/22/39/39	0/3/3/3
2	DSZ	B	701	3	-	4/22/39/39	0/3/3/3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	701	DSZ	O-C-CA-N
2	B	701	DSZ	O-C-CA-N
2	B	701	DSZ	NAT-C-CA-N
2	A	701	DSZ	O-C-CA-CB
2	B	701	DSZ	O-C-CA-CB
2	B	701	DSZ	NAT-C-CA-CB
2	A	701	DSZ	NAT-C-CA-N
2	A	701	DSZ	NAT-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	502/536 (93%)	-0.06	28 (5%)	30 32	15, 31, 69, 98	10 (1%)
1	B	499/536 (93%)	-0.14	15 (3%)	52 57	15, 31, 64, 95	3 (0%)
All	All	1001/1072 (93%)	-0.10	43 (4%)	40 42	15, 31, 66, 98	13 (1%)

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	289	THR	7.9
1	B	295	ILE	6.9
1	B	294	SER	6.3
1	A	253	ILE	4.6
1	A	130	LYS	4.0
1	A	251	VAL	3.8
1	A	295	ILE	3.8
1	A	178	ALA	3.7
1	A	101	LEU	3.7
1	A	494	LEU	3.7
1	A	351	SER	3.7
1	A	100	ASN	3.6
1	A	255	SER	3.6
1	A	353	GLY	3.4
1	B	254	ASP	3.3
1	B	135	GLU	3.2
1	B	102	LYS	3.0
1	A	350	SER	3.0
1	B	103	ASN	3.0
1	A	250	GLU	3.0
1	B	399	ASN	2.9
1	A	132	ASN	2.8
1	B	150	LYS	2.8
1	A	252	PRO	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	178	ALA	2.6
1	B	134	LYS	2.6
1	A	443	GLY	2.5
1	A	135	GLU	2.5
1	A	219	LYS	2.4
1	B	355	ALA	2.4
1	A	403	TYR	2.3
1	A	99	GLU	2.2
1	B	182	GLN	2.2
1	A	126	ILE	2.2
1	B	218	ILE	2.2
1	B	179	HIS	2.2
1	A	218	ILE	2.2
1	A	116	ILE	2.1
1	A	289	THR	2.1
1	A	133	GLU	2.1
1	A	360	ILE	2.0
1	A	209	LEU	2.0
1	A	254	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MG	A	702	1/1	0.90	0.08	37,37,37,37	1
3	MG	A	703	1/1	0.97	0.05	30,30,30,30	1
2	DSZ	A	701	31/31	0.99	0.04	18,22,25,25	0
2	DSZ	B	701	31/31	0.99	0.03	18,23,25,25	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MG	B	702	1/1	0.99	0.02	28,28,28,28	1
3	MG	B	703	1/1	0.99	0.02	32,32,32,32	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around MG A 702:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MG A 703:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around DSZ A 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around DSZ B 701:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MG B 702:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MG B 703:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.