



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 07:14 PM UTC

PDB ID : 3M6S / pdb_00003m6s
Title : Crystal structure of H1N1pdm Hemagglutinin
Authors : Yang, H.; Carney, P.J.; Stevens, J.
Deposited on : 2010-03-16
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	331	2% 78% 16% . .
1	C	331	2% 73% 19% 5% .
1	E	331	3% 75% 17% 5% .
1	G	331	3% 76% 16% 5% .
1	I	331	6% 77% 16% . .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	K	331	
2	B	181	
2	D	181	
2	F	181	
2	H	181	
2	J	181	
2	L	181	
3	M	2	
3	O	2	
3	P	2	
3	Q	2	
4	N	3	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	MAN	N	3	X	-	-	-
5	NAG	G	331	X	-	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 23587 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	325	Total	C	N	O	S	0	0	0
			2533	1602	435	485	11			
1	C	323	Total	C	N	O	S	0	0	0
			2518	1593	433	481	11			
1	E	323	Total	C	N	O	S	0	0	0
			2518	1593	433	481	11			
1	G	322	Total	C	N	O	S	0	0	0
			2514	1591	432	480	11			
1	I	322	Total	C	N	O	S	0	0	0
			2511	1588	432	480	11			
1	K	321	Total	C	N	O	S	0	0	0
			2506	1587	431	477	11			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	ALA	-	expression tag	UNP C5MV42
A	-2	ASP	-	expression tag	UNP C5MV42
A	-1	PRO	-	expression tag	UNP C5MV42
A	0	GLY	-	expression tag	UNP C5MV42
C	-3	ALA	-	expression tag	UNP C5MV42
C	-2	ASP	-	expression tag	UNP C5MV42
C	-1	PRO	-	expression tag	UNP C5MV42
C	0	GLY	-	expression tag	UNP C5MV42
E	-3	ALA	-	expression tag	UNP C5MV42
E	-2	ASP	-	expression tag	UNP C5MV42
E	-1	PRO	-	expression tag	UNP C5MV42
E	0	GLY	-	expression tag	UNP C5MV42
G	-3	ALA	-	expression tag	UNP C5MV42
G	-2	ASP	-	expression tag	UNP C5MV42
G	-1	PRO	-	expression tag	UNP C5MV42
G	0	GLY	-	expression tag	UNP C5MV42
I	-3	ALA	-	expression tag	UNP C5MV42

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
I	-2	ASP	-	expression tag	UNP C5MV42
I	-1	PRO	-	expression tag	UNP C5MV42
I	0	GLY	-	expression tag	UNP C5MV42
K	-3	ALA	-	expression tag	UNP C5MV42
K	-2	ASP	-	expression tag	UNP C5MV42
K	-1	PRO	-	expression tag	UNP C5MV42
K	0	GLY	-	expression tag	UNP C5MV42

- Molecule 2 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	169	Total	C	N	O	S	5	0	0
			1368	860	232	270	6			
2	D	171	Total	C	N	O	S	0	0	0
			1386	870	234	276	6			
2	F	171	Total	C	N	O	S	0	0	0
			1386	870	234	276	6			
2	H	170	Total	C	N	O	S	0	0	0
			1377	865	233	273	6			
2	J	170	Total	C	N	O	S	0	0	0
			1377	865	233	273	6			
2	L	171	Total	C	N	O	S	0	0	0
			1386	870	234	276	6			

There are 36 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	175	SER	GLY	engineered mutation	UNP C5MV42
B	176	GLY	VAL	engineered mutation	UNP C5MV42
B	177	ARG	LYS	engineered mutation	UNP C5MV42
B	179	VAL	-	expression tag	UNP C5MV42
B	180	PRO	-	expression tag	UNP C5MV42
B	181	ARG	-	expression tag	UNP C5MV42
D	175	SER	GLY	engineered mutation	UNP C5MV42
D	176	GLY	VAL	engineered mutation	UNP C5MV42
D	177	ARG	LYS	engineered mutation	UNP C5MV42
D	179	VAL	-	expression tag	UNP C5MV42
D	180	PRO	-	expression tag	UNP C5MV42
D	181	ARG	-	expression tag	UNP C5MV42
F	175	SER	GLY	engineered mutation	UNP C5MV42
F	176	GLY	VAL	engineered mutation	UNP C5MV42
F	177	ARG	LYS	engineered mutation	UNP C5MV42

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	179	VAL	-	expression tag	UNP C5MV42
F	180	PRO	-	expression tag	UNP C5MV42
F	181	ARG	-	expression tag	UNP C5MV42
H	175	SER	GLY	engineered mutation	UNP C5MV42
H	176	GLY	VAL	engineered mutation	UNP C5MV42
H	177	ARG	LYS	engineered mutation	UNP C5MV42
H	179	VAL	-	expression tag	UNP C5MV42
H	180	PRO	-	expression tag	UNP C5MV42
H	181	ARG	-	expression tag	UNP C5MV42
J	175	SER	GLY	engineered mutation	UNP C5MV42
J	176	GLY	VAL	engineered mutation	UNP C5MV42
J	177	ARG	LYS	engineered mutation	UNP C5MV42
J	179	VAL	-	expression tag	UNP C5MV42
J	180	PRO	-	expression tag	UNP C5MV42
J	181	ARG	-	expression tag	UNP C5MV42
L	175	SER	GLY	engineered mutation	UNP C5MV42
L	176	GLY	VAL	engineered mutation	UNP C5MV42
L	177	ARG	LYS	engineered mutation	UNP C5MV42
L	179	VAL	-	expression tag	UNP C5MV42
L	180	PRO	-	expression tag	UNP C5MV42
L	181	ARG	-	expression tag	UNP C5MV42

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	M	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	O	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	P	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	Q	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	N	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	E	1	Total	C	N	O	0	0
			14	8	1	5		
5	G	1	Total	C	N	O	0	0
			14	8	1	5		
5	G	1	Total	C	N	O	0	0
			14	8	1	5		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Hemagglutinin



• Molecule 1: Hemagglutinin





• Molecule 1: Hemagglutinin








• Molecule 1: Hemagglutinin



• Molecule 1: Hemagglutinin




• Molecule 1: Hemagglutinin




• Molecule 2: Hemagglutinin







• Molecule 2: Hemagglutinin




• Molecule 2: Hemagglutinin




• Molecule 2: Hemagglutinin



• Molecule 2: Hemagglutinin




- Molecule 2: Hemagglutinin



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	73.98Å 109.71Å 129.90Å 86.25° 74.68° 75.10°	Depositor
Resolution (Å)	50.00 – 2.80 50.00 – 2.80	Depositor EDS
% Data completeness (in resolution range)	98.1 (50.00-2.80) 98.1 (50.00-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.32 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.230 , 0.256 0.227 , 0.251	Depositor DCC
R_{free} test set	4608 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	50.1	Xtriage
Anisotropy	0.013	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 47.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.016 for -h,-k,-h+l	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	23587	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.91	1/2597 (0.0%)	1.12	9/3530 (0.3%)
1	C	0.97	2/2581 (0.1%)	1.15	7/3507 (0.2%)
1	E	0.89	2/2581 (0.1%)	1.10	9/3507 (0.3%)
1	G	1.02	2/2577 (0.1%)	1.18	17/3502 (0.5%)
1	I	0.83	3/2573 (0.1%)	1.06	8/3495 (0.2%)
1	K	0.92	3/2569 (0.1%)	1.10	7/3491 (0.2%)
2	B	0.82	0/1396	1.07	8/1881 (0.4%)
2	D	0.90	0/1414	1.03	2/1905 (0.1%)
2	F	0.79	0/1414	1.01	2/1905 (0.1%)
2	H	0.84	0/1405	1.00	4/1893 (0.2%)
2	J	0.92	0/1405	1.05	5/1893 (0.3%)
2	L	0.83	0/1414	1.01	5/1905 (0.3%)
All	All	0.90	13/23926 (0.1%)	1.09	83/32414 (0.3%)

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	I	298	ILE	CA-CB	6.46	1.61	1.53
1	K	298	ILE	CA-CB	6.22	1.61	1.53
1	G	267	ILE	CA-CB	6.02	1.61	1.53
1	C	72	THR	CA-CB	5.74	1.60	1.53
1	A	267	ILE	CA-CB	5.61	1.60	1.54

The worst 5 of 83 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	143	LYS	CB-CA-C	12.37	129.19	109.90
1	A	116	ILE	CB-CA-C	-8.35	100.87	112.14
1	G	116	ILE	CB-CA-C	-8.14	100.99	112.22
1	I	10	ASN	N-CA-C	8.04	118.57	108.45
1	I	2	THR	N-CA-C	7.96	120.48	108.46

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2533	0	2483	41	1
1	C	2518	0	2472	53	0
1	E	2518	0	2472	44	1
1	G	2514	0	2466	37	2
1	I	2511	0	2464	63	0
1	K	2506	0	2460	60	1
2	B	1368	0	1293	37	3
2	D	1386	0	1305	42	0
2	F	1386	0	1305	38	0
2	H	1377	0	1299	20	0
2	J	1377	0	1299	56	0
2	L	1386	0	1305	46	0
3	M	28	0	25	2	0
3	O	28	0	24	2	0
3	P	28	0	25	1	0
3	Q	28	0	25	0	0
4	N	39	0	34	1	0
5	A	14	0	13	1	0
5	E	14	0	13	1	0
5	G	28	0	26	0	0
All	All	23587	0	22808	401	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 401 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:45:ARG:HG3	1:I:45:ARG:HH11	1.07	1.10
1:E:45:ARG:HG3	1:E:45:ARG:HH11	1.09	1.10
1:I:5:ILE:HD13	2:J:24:TYR:CD1	1.87	1.09

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:4:CYS:O	2:J:24:TYR:HA	1.52	1.09
1:A:45:ARG:HG3	1:A:45:ARG:HH11	1.14	1.08

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:143:LYS:CE	1:G:224:GLU:OE1[1_466]	1.91	0.29
1:A:168:ASP:O	1:K:273:HIS:CE1[1_655]	1.98	0.22
2:B:169:ASN:OD1	1:G:222:ASP:OD2[1_466]	2.13	0.07
2:B:164:GLU:OE1	1:E:120:THR:O[1_556]	2.19	0.01

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	323/331 (98%)	305 (94%)	16 (5%)	2 (1%)	21	51
1	C	321/331 (97%)	303 (94%)	16 (5%)	2 (1%)	21	51
1	E	321/331 (97%)	303 (94%)	16 (5%)	2 (1%)	21	51
1	G	320/331 (97%)	303 (95%)	16 (5%)	1 (0%)	36	66
1	I	320/331 (97%)	299 (93%)	19 (6%)	2 (1%)	21	51
1	K	319/331 (96%)	300 (94%)	18 (6%)	1 (0%)	36	66
2	B	167/181 (92%)	163 (98%)	4 (2%)	0	100	100
2	D	169/181 (93%)	163 (96%)	6 (4%)	0	100	100
2	F	169/181 (93%)	164 (97%)	5 (3%)	0	100	100
2	H	168/181 (93%)	164 (98%)	4 (2%)	0	100	100
2	J	168/181 (93%)	162 (96%)	6 (4%)	0	100	100
2	L	169/181 (93%)	162 (96%)	7 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	2934/3072 (96%)	2791 (95%)	133 (4%)	10 (0%)	36	66

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	261	ALA
1	G	261	ALA
1	I	261	ALA
1	K	261	ALA
1	C	261	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	284/289 (98%)	254 (89%)	30 (11%)	6	22
1	C	282/289 (98%)	245 (87%)	37 (13%)	4	14
1	E	282/289 (98%)	246 (87%)	36 (13%)	4	15
1	G	282/289 (98%)	252 (89%)	30 (11%)	6	22
1	I	281/289 (97%)	249 (89%)	32 (11%)	5	19
1	K	281/289 (97%)	243 (86%)	38 (14%)	4	13
2	B	146/156 (94%)	129 (88%)	17 (12%)	5	18
2	D	148/156 (95%)	132 (89%)	16 (11%)	6	21
2	F	148/156 (95%)	134 (90%)	14 (10%)	8	26
2	H	147/156 (94%)	129 (88%)	18 (12%)	5	16
2	J	147/156 (94%)	132 (90%)	15 (10%)	7	23
2	L	148/156 (95%)	133 (90%)	15 (10%)	7	23
All	All	2576/2670 (96%)	2278 (88%)	298 (12%)	5	18

5 of 298 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	J	38	LEU
2	L	64	THR
2	J	91	ILE
1	K	152	VAL
2	D	121	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 27 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	51	HIS
1	I	31	ASN
1	K	223	GLN
1	G	223	GLN
1	I	51	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

11 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	M	1	3,1	14,14,15	0.77	0	17,19,21	1.98	3 (17%)
3	NAG	M	2	3	14,14,15	0.94	1 (7%)	17,19,21	2.04	6 (35%)
4	NAG	N	1	1,4	14,14,15	0.81	0	17,19,21	1.59	3 (17%)
4	NAG	N	2	4	14,14,15	0.77	0	17,19,21	2.08	5 (29%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	MAN	N	3	4	11,11,12	0.87	0	15,15,17	2.67	6 (40%)
3	NAG	O	1	3,1	14,14,15	0.55	0	17,19,21	2.23	5 (29%)
3	NAG	O	2	3	14,14,15	1.06	1 (7%)	17,19,21	1.51	4 (23%)
3	NAG	P	1	3,1	14,14,15	1.19	1 (7%)	17,19,21	2.41	5 (29%)
3	NAG	P	2	3	14,14,15	0.62	0	17,19,21	1.30	3 (17%)
3	NAG	Q	1	3,1	14,14,15	1.84	2 (14%)	17,19,21	1.81	4 (23%)
3	NAG	Q	2	3	14,14,15	1.02	1 (7%)	17,19,21	2.06	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	M	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	M	2	3	-	0/6/23/26	0/1/1/1
4	NAG	N	1	1,4	-	1/6/23/26	0/1/1/1
4	NAG	N	2	4	-	2/6/23/26	0/1/1/1
4	MAN	N	3	4	1/1/4/5	2/2/19/22	0/1/1/1
3	NAG	O	1	3,1	-	1/6/23/26	0/1/1/1
3	NAG	O	2	3	-	0/6/23/26	0/1/1/1
3	NAG	P	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	P	2	3	-	0/6/23/26	0/1/1/1
3	NAG	Q	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	Q	2	3	-	0/6/23/26	0/1/1/1

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	Q	1	NAG	C1-C2	5.59	1.60	1.52
3	P	1	NAG	C1-C2	4.16	1.58	1.52
3	O	2	NAG	C1-C2	2.80	1.56	1.52
3	Q	2	NAG	C1-C2	2.38	1.55	1.52
3	M	2	NAG	C1-C2	2.34	1.55	1.52

The worst 5 of 48 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	M	1	NAG	O5-C1-C2	-6.13	101.81	111.29
3	O	1	NAG	C1-O5-C5	5.99	120.22	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	N	3	MAN	C1-O5-C5	5.50	119.56	112.19
3	P	1	NAG	C1-O5-C5	5.11	119.03	112.19
4	N	2	NAG	C1-O5-C5	4.96	118.83	112.19

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	N	3	MAN	C1

5 of 8 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	N	3	MAN	C4-C5-C6-O6
3	M	1	NAG	O5-C5-C6-O6
4	N	3	MAN	O5-C5-C6-O6
3	M	1	NAG	C4-C5-C6-O6
4	N	2	NAG	C4-C5-C6-O6


There are no ring outliers.


5 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	O	2	NAG	1	0
4	N	1	NAG	1	0
3	M	1	NAG	2	0
3	O	1	NAG	2	0
3	P	1	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	G	331	1	14,14,15	1.60	2 (14%)	17,19,21	2.34	7 (41%)
5	NAG	E	328	1	14,14,15	0.54	0	17,19,21	1.37	2 (11%)
5	NAG	A	328	1	14,14,15	0.88	1 (7%)	17,19,21	1.59	3 (17%)
5	NAG	G	332	1	14,14,15	1.03	1 (7%)	17,19,21	2.73	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	G	331	1	1/1/5/7	2/6/23/26	0/1/1/1
5	NAG	E	328	1	-	2/6/23/26	0/1/1/1
5	NAG	A	328	1	-	2/6/23/26	0/1/1/1
5	NAG	G	332	1	-	0/6/23/26	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	G	331	NAG	C1-C2	4.88	1.59	1.52
5	G	332	NAG	C1-C2	2.65	1.56	1.52
5	G	331	NAG	C3-C2	2.39	1.57	1.52
5	A	328	NAG	C1-C2	2.29	1.55	1.52

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	G	332	NAG	C1-O5-C5	10.26	125.93	112.19
5	G	331	NAG	C1-O5-C5	6.58	121.01	112.19
5	A	328	NAG	C1-O5-C5	3.99	117.53	112.19
5	G	331	NAG	C3-C4-C5	-3.09	104.63	110.23
5	A	328	NAG	C2-N2-C7	-2.98	118.91	122.90

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	G	331	NAG	C1

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	328	NAG	C4-C5-C6-O6
5	A	328	NAG	O5-C5-C6-O6
5	E	328	NAG	C4-C5-C6-O6
5	E	328	NAG	O5-C5-C6-O6
5	G	331	NAG	O5-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	E	328	NAG	1	0
5	A	328	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	325/331 (98%)	0.13	8 (2%) 58 48	7, 20, 33, 49	0
1	C	323/331 (97%)	-0.07	8 (2%) 58 48	7, 20, 33, 49	0
1	E	323/331 (97%)	-0.03	10 (3%) 51 41	8, 20, 33, 48	0
1	G	322/331 (97%)	-0.04	11 (3%) 48 39	8, 20, 33, 49	0
1	I	322/331 (97%)	0.57	19 (5%) 28 21	8, 20, 33, 48	0
1	K	321/331 (96%)	0.26	12 (3%) 45 36	8, 20, 33, 49	0
2	B	169/181 (93%)	0.34	9 (5%) 32 24	4, 31, 50, 69	1 (0%)
2	D	171/181 (94%)	0.26	4 (2%) 61 51	4, 32, 53, 69	0
2	F	171/181 (94%)	0.59	12 (7%) 22 16	4, 32, 50, 69	0
2	H	170/181 (93%)	0.90	20 (11%) 9 7	4, 32, 50, 69	0
2	J	170/181 (93%)	0.68	8 (4%) 36 29	4, 32, 50, 69	0
2	L	171/181 (94%)	1.57	49 (28%) 1 1	4, 32, 50, 69	0
All	All	2958/3072 (96%)	0.34	170 (5%) 29 22	4, 22, 44, 69	1 (0%)

The worst 5 of 170 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	L	71	ASN	6.7
1	I	71	SER	5.7
2	L	73	LEU	5.6
1	G	72	THR	5.5
2	L	67	GLY	5.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MAN	N	3	11/12	0.46	0.16	77,83,85,86	0
3	NAG	Q	2	14/15	0.47	0.19	81,84,88,88	0
3	NAG	Q	1	14/15	0.63	0.15	56,67,74,78	0
3	NAG	O	2	14/15	0.71	0.17	76,80,82,82	0
3	NAG	M	2	14/15	0.75	0.13	73,77,78,79	0
3	NAG	P	2	14/15	0.78	0.14	89,94,101,101	0
3	NAG	O	1	14/15	0.83	0.11	52,60,68,69	0
3	NAG	P	1	14/15	0.89	0.11	61,68,73,82	0
4	NAG	N	2	14/15	0.92	0.08	45,58,63,71	0
3	NAG	M	1	14/15	0.92	0.09	47,52,60,66	0
4	NAG	N	1	14/15	0.93	0.08	48,53,55,57	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain O:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain P:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain Q:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain N:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	G	331	14/15	0.55	0.19	59,72,74,76	0
5	NAG	G	332	14/15	0.68	0.15	64,74,80,81	0
5	NAG	A	328	14/15	0.83	0.12	49,55,59,59	0
5	NAG	E	328	14/15	0.88	0.10	46,51,53,55	0

6.5 Other polymers [i](#)

There are no such residues in this entry.